

NUMBER THEORETICAL LOCALLY RECOVERABLE CODES

ANDREA FERRAGUTI, DORIAN GOLDFELD, AND GIACOMO MICHELI

ABSTRACT. In this paper we give constructions for infinite sequences of finite non-linear locally recoverable codes $\mathcal{C} \subseteq \prod_{i=1}^N \mathbb{F}_{q_i}$ over a product of finite fields arising from basis expansions in algebraic number fields. The codes in our sequences have increasing length and size, constant rate, fixed locality, and minimum distance going to infinity.

1. INTRODUCTION

There has been a lot of interest recently in Locally Recoverable Codes (LRC) [1, 2, 3, 4, 7, 8, 9, 10, 11], which is a special class of linear codes that allow local recovery of erasures. More specifically, they allow recovery of simultaneous erasures exactly as commonly used codes do (e.g. Reed-Solomon codes) by looking at the entries of a codeword that correspond to an information set (i.e. k other components where no erasure happened), but they also allow recovery of a single erasure by looking at fewer nodes than k .

The applications in which LRC thrive are related to distributed storage and cloud storage systems because they easily allow the recovery of the data in a single failed server or hard-drive (which in this context we simply call *node*), but they also allow recovery of more serious failures, such as simultaneous failures of multiple nodes in the system.

In this paper we construct Locally Recoverable Codes using Number Fields. The technique to construct these codes is purely number theoretical, but the constructed codes are subsets of a product of finite fields (so they are practical to implement). We believe that having LRCs over products of different finite fields is an interesting feature as it provides more flexibility in the design of a system: for example, we might want to increase the length of these codes without having to enlarge the base field for all the components of every codeword, as one would be forced to do in the context of Reed-Solomon codes when one wants to extend the length of the code beyond the size of the finite field. In fact, with our construction, we can simply take additional reductions of elements of \mathcal{O}_K modulo other prime ideals of \mathcal{O}_K . Of course, our codes are not linear because of the nature of a product of finite fields and the

2020 *Mathematics Subject Classification.* 11T06.

Dorian Goldfeld is partially supported by Simons Collaboration Grant 567168.

Giacomo Micheli was supported in part by the National Science Foundation under Grant No 2127742.

fact that $(\mathcal{O}_K, +)$ does not have an \mathbb{F}_q linear structure for any q . Nevertheless, they allow efficient local recovery and have good minimum distance.

In addition, we can construct a family of codes such that the distance grows linearly with length and dimension, and the asymptotic rate (see Definition 2.1) can be made constant. From the methodology standpoint, our construction builds up new interactions between analytic number theory and coding theory, which stems from the idea that one can look at an integer as a constant function from the set of primes $\{\mathfrak{p}_i\}_{i \in \{1, \dots, n\}}$ of the ring of integers lying over a totally split rational prime p , to \mathbb{F}_p .

2. BACKGROUND ON CODING THEORY

Let n be a positive integer and F_1, \dots, F_n be finite fields ordered by increasing size. We define a code \mathcal{C} in $R_m = \prod_{i=1}^m F_i$ as a subset of R_m . The distance between two codewords $x, y \in \mathcal{C}$ is the number of indexes $i \in \{1, \dots, n\}$ such that $x_i \neq y_i$. The minimum distance of \mathcal{C} is the minimal positive integer d such that there exist two elements $x, y \in \mathcal{C}$ such that $d(x, y) = d$. If X is a set, let us denote the powerset of X as 2^X .

For every $m \in \mathbb{N}$, let $\mathcal{C}_m \in 2^{R_m}$ be a choice of a code of R_m . For any set A , let us denote by $\#A$ the cardinality of A . We say that a sequence of codes $\{\mathcal{C}_m\}_{m \in \mathbb{N}}$ is *almost good* if

$$(2.1) \quad \liminf_{m \rightarrow \infty} \frac{\log(\#\mathcal{C}_m)}{\log(\#R_m)} = \gamma > 0$$

and

$$(2.2) \quad \liminf_{m \rightarrow \infty} d(\mathcal{C}_m) = +\infty$$

Notice that the choice of the basis of the logarithm clearly does not affect the definition.

Also, notice the difference between our definition and the usual definition of a good family of codes, where the ratio between distance and length is required to converge to a constant. In our case, since the distance and locality are not weighted by how large the finite fields we are using are, the standard definition of good codes carries an inherent disadvantage that is essentially unavoidable. For this reason, we do keep track of the size of the code vs the size of the full space but for simplicity we avoid weighting the distance d depending on the finite fields where the components belong, and only require $d \rightarrow \infty$ (even though in our case the growth is linear, which is what happens with optimal codes).

We say that a code $\mathcal{C} \subseteq R_m$ has locality r if for any $i \in \{1, \dots, n\}$ there is a method to reconstruct the i -th component of a codeword $c \in \mathcal{C}$ by knowing at most r other components of c .

3. BACKGROUND ON NUMBER FIELDS

Let K/\mathbb{Q} be a number field of degree δ . Recall that if $\beta \in K$, the *norm* of β , denoted by $N(\beta)$, is the determinant of the \mathbb{Q} -linear map $K \rightarrow K$ defined by $x \mapsto \beta x$. Let \mathcal{O}_K be the ring of integers of K , and let $\alpha \in \mathcal{O}_K$ be an element that generates K , i.e. such that $\mathbb{Q}(\alpha) = K$. Let

$$m_\alpha(x) = b_0 + b_1x + \dots + b_{\delta-1}x^{\delta-1} + x^\delta \in \mathbb{Z}[x]$$

be the minimal polynomial of α over \mathbb{Q} , and let $S := \max\{|b_i| : i \in \{0, \dots, \delta-1\}\}$, where $|\cdot|$ denotes the usual archimedean absolute value. For any prime ideal \mathfrak{p} of \mathcal{O}_K , let $\mathbb{F}_\mathfrak{p}$ be the field $\mathcal{O}_K/\mathfrak{p}$.

Recall that a prime $p \in \mathbb{N}$ is *totally split* in K/\mathbb{Q} if $p\mathcal{O}_K$ factors as $\prod_{i=1}^{[K:\mathbb{Q}]} \mathfrak{p}_i$, where the \mathfrak{p}_i 's are pairwise distinct prime ideals of \mathcal{O}_K and $[\mathbb{F}_{\mathfrak{p}_i} : \mathbb{F}_p] = 1$ for all $i \in \{1, \dots, \delta\}$.

Lemma 3.1. *With the notation above, let $y = \sum_{i=0}^{\delta-1} z_i \alpha^i \in \mathcal{O}_K$, with $z_i \in \mathbb{Z}$ and $|z_i| < M$ for every i . Then $|N(y)| \leq \delta^{\delta/2} (1+S)^{(\delta-1)\delta/2} (M-1)^\delta$.*

Proof. Clearly we can assume that $M > 1$, as otherwise the claim is trivial. The set $\mathcal{B} := \{1, \alpha, \dots, \alpha^{\delta-1}\}$ is a \mathbb{Q} -basis of K by assumption. Notice that if $w := \sum_{i=0}^{\delta-1} w_i \alpha^i \in \mathcal{O}_K$, with $w_i \in \mathbb{Z}$ for every i , then

$$\begin{aligned} w \cdot \alpha &= \sum_{i=0}^{\delta-1} w_i \alpha^{i+1} \\ &= \sum_{i=1}^{\delta-1} w_{i-1} \alpha^i + w_\delta \alpha^\delta \end{aligned}$$

Using now that $m_\alpha(\alpha) = 0$ we get that

$$(3.1) \quad w \cdot \alpha = -b_0 w_{\delta-1} + \sum_{i=1}^{\delta-1} (w_{i-1} - b_i w_{\delta-1}) \alpha^i.$$

Let A_y be the multiplication-by- y matrix with respect to the basis \mathcal{B} , where the elements of K , expressed in the basis \mathcal{B} are considered as column vectors. More precisely, A_y is the matrix that makes the following diagram commutative

$$\begin{array}{ccc} K & \xrightarrow{y \cdot} & K \\ \downarrow \iota & & \downarrow \iota \\ \mathbb{Q}^\delta & \xrightarrow{A_y \cdot} & \mathbb{Q}^\delta \end{array}$$

where ι is the usual isomorphism of vector spaces that sends an element of K into its expression in the basis \mathcal{B} . We claim that entries in the j -th column of A_y are bounded, in absolute value, by $(M-1)(1+S)^{j-1}$. For $j = 1$ this is obvious since

entries in the first column are the coefficients of $y \cdot 1 = y$ with respect to \mathcal{B} . Now suppose that the claim is true for the j -th column and let us prove it for the $(j+1)$ -th. The j -th column is given by the result of the multiplication

$$y \cdot \alpha^{j-1} = \sum_{i=0}^{\delta-1} c_i \alpha^i,$$

with $|c_i| \leq (M-1)(1+S)^{j-1}$ for every $i \in \{0, \dots, \delta-1\}$ by the inductive hypothesis. Now let us consider the $(j+1)$ -th columns, given by the multiplication

$$y \cdot \alpha^j = \sum_{i=0}^{\delta-1} d_i \alpha^i$$

with $d_0, \dots, d_{\delta-1} \in \mathbb{Z}$. Since $y \cdot \alpha^j = (y \cdot \alpha^{j-1}) \cdot \alpha$, equation (3.1) and the inductive hypothesis show that $|d_0| \leq |b_0 c_{\delta-1}| \leq S(1+S)^{j-1}$ and $|d_i| \leq |c_{i-1}| + |b_i c_{\delta-1}| \leq (M-1)(1+S)^j$. The claim follows since $S(1+S)^{j-1} \leq (M-1)(1+S)^j$ for every $j \geq 0$.

Now the bound on $|N(y)| = |\det A_y|$ follows from Hadamard's inequality, which states that the determinant of a complex matrix is bounded, in absolute value by the product of the norms of the column vectors c_j of A_y . In fact,

$$|\det A_y| \leq \prod_{j=1}^{\delta} |c_j| \leq \prod_{j=1}^{\delta} \sqrt{\delta(M-1)^2(1+S)^{2j}} = \delta^{\delta/2} (M-1)^{\delta} (1+S)^{\delta(\delta-1)/2}.$$

□

4. CONSTRUCTION OF NUMBER THEORETICAL LOCALLY RECOVERABLE CODES

4.1. Overview of the construction. First, we construct an ambient code \mathcal{D} (that is essentially a Chinese remainder code), for which we can prove nice distance properties. After that, we will extract a subcode of \mathcal{D} that verifies the locality property we are seeking for. Finally we show how to construct almost good families of locally recoverable codes in the sense of equations (2.1) and (2.2).

4.2. Construction of the ambient code \mathcal{D} . First, we need to construct Chinese remainder codes that are similar to the Reed-Solomon Codes (à-la Guruswami, see [5]).

Let K be a number field of degree δ with ring of integers \mathcal{O}_K , and let $\alpha \in \mathcal{O}_K$ be such that $\mathbb{Q}(\alpha) = K$. For $M \geq 1$ we define

$$\mathcal{R}[M] := \left\{ \sum_{i=0}^{\delta-1} z_i \alpha^i \mid 0 \leq z_i < M, \quad \forall i \in \{0, \dots, \delta-1\} \right\}.$$

Let $m_\alpha(x) = b_0 + b_1 x + \dots + x^\delta \in \mathbb{Z}[x]$ be the minimal polynomial of α over \mathbb{Q} , and let $S := \max\{|b_i| : i = 0, \dots, \delta-1\}$. Let

$$(4.1) \quad C_\alpha := \delta^{\delta/2} (1+S)^{(\delta-1)\delta/2}$$

so that by Lemma 3.1 we have that $|N(y)| \leq C_\alpha \cdot (M-1)^\delta$ for every $y \in \mathcal{R}[M]$.

Let $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ be distinct prime ideals of \mathcal{O}_K , ordered by increasing norm size, and for every i let $\mathbb{F}_{\mathfrak{p}_i} := \mathcal{O}_K/\mathfrak{p}_i$. Assume that $\prod_{i=1}^n N(\mathfrak{p}_i) > C_\alpha \cdot (M-1)^\delta$ (this is needed to achieve injectivity of the encoding map ϕ defined below).

The number theoretical Reed-Solomon code $\mathcal{D} = \mathcal{D}(K, M, \{\mathfrak{p}_i\}_{i \in \{1, \dots, n\}})$ is defined as the image $\phi(\mathcal{R}[M])$ of the map

$$\begin{aligned} \phi : \mathcal{R}[M] &\longrightarrow \prod_{i=1}^n \mathbb{F}_{\mathfrak{p}_i} \\ y &\mapsto (y \bmod \mathfrak{p}_1, \dots, y \bmod \mathfrak{p}_n) \end{aligned}$$

See [6] for more on this. In the rest of the paper we will refer to ϕ as the *encoding map*.

Theorem 4.1. *Let \mathcal{D} be the code defined above, and let $\mathcal{P} := \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$. Let $d(\mathcal{D})$ be the minimal distance of \mathcal{D} and let*

$$m := \min_{T \subseteq \mathcal{P}} \left\{ \#T : \prod_{\mathfrak{p} \in T} N(\mathfrak{p}) > C_\alpha \cdot (M-1)^\delta \right\}.$$

Then the following hold.

- (1) *The map ϕ is injective.*
- (2) *$d(\mathcal{D}) \geq n - m + 1$.*
- (3) *If $\prod_{\mathfrak{p} \in U} N(\mathfrak{p}) < M^\delta$ for some $U \subseteq \mathcal{P}$ with $\#U = m - 1$, then equality holds in (2).*

Remark 4.2. Since ϕ is thought as the encoding map, its injectivity is fundamental because we want that different messages are mapped to different codewords. This is achieved by adding just enough redundancy by considering at least n distinct reductions, where n is chosen such that $\prod_{i=1}^n N(\mathfrak{p}_i) > C_\alpha \cdot (M-1)^\delta$, as we will explain.

Item (2) provides the code with a lower bound for the minimum distance: in fact the more redundancy is added (and therefore n grows because we provide reductions at many prime ideals) the more the minimum distance grows.

Item (3) ensures that, as far as the product of the norms is not too large then there are indeed two codewords at distance $m - 1$.

Proof. Let $T \subseteq \mathcal{P}$ be a subset of cardinality m such that $\prod_{\mathfrak{p} \in T} N(\mathfrak{p}) > C_\alpha \cdot (M-1)^\delta$, and let $y_1, y_2 \in \mathcal{R}[M]$ be such that $\phi(y_1) = \phi(y_2)$. In particular, we have that $y_1 \equiv y_2 \pmod{\mathfrak{p}}$ for every $\mathfrak{p} \in T$. It follows that $\prod_{\mathfrak{p} \in T} N(\mathfrak{p}) \mid N(y_1 - y_2)$. By Lemma 3.1 we have that $N(y_1 - y_2) \leq C_\alpha \cdot (M-1)^\delta$, and hence by the definition of m we must have that $y_1 = y_2$. This proves (1) and (2) at the same time.

To prove (3), notice that if $\prod_{\mathfrak{p} \in U} N(\mathfrak{p}) < M^\delta$ then the map $\mathcal{R}[M] \rightarrow \prod_{\mathfrak{p} \in U} \mathbb{F}_{\mathfrak{p}}$ is not injective for cardinality reasons. It follows that there are $y_1 \neq y_2 \in \mathcal{R}[M]$ such

that $y_1 \equiv y_2 \pmod{\mathfrak{p}}$ for all $\mathfrak{p} \in U$. On the other hand $\phi(y_1) \neq \phi(y_2)$ by (1), and hence $\phi(y_1)$ and $\phi(y_2)$ have distance $n - m + 1$. \square

4.3. Construction of the locally recoverable code \mathcal{C} as a subset of \mathcal{D} . Let K be a number field of degree $r + 1$ with ring of integers \mathcal{O}_K , and let $M \in \mathbb{N}$. Let $\alpha \in \mathcal{O}_K$ be such that $K = \mathbb{Q}(\alpha)$. Let

$$\mathcal{R}[M]^- = \left\{ \sum_{i=0}^{r-1} a_i \alpha^i \mid 0 \leq a_i < M \quad \forall i \in \{0, \dots, r-1\} \right\} \subsetneq \mathcal{R}[M].$$

Notice that this differs from the set $\mathcal{R}[M]$ previously defined, as we are forcing the coefficient of α^r to be 0 (this is a strictly smaller set of elements than $\mathcal{R}[M]$ since the minimal polynomial of α has degree $r + 1$). Let now s be a positive integer and define

$$\mathcal{A}[M] = \left\{ \sum_{j=0}^s f_j M^j : f_j \in \mathcal{R}[M]^- \quad \forall j \in \{0, \dots, s\} \right\}.$$

Lemma 4.3.

$$\#\mathcal{A}[M] = M^{r(s+1)}$$

Proof. This follows from the fact that elements of $\mathcal{R}[M]$ are a complete set of representatives for the quotient $\mathcal{O}_K/(M)$. Hence if $\sum_{j=0}^{r-1} f_j M^j = \sum_{j=0}^{r-1} g_j M^j$ then $f_0 \equiv g_0 \pmod{M}$, but this implies that $f_0 = g_0$. The claim follows then by an easy induction. \square

Let ℓ be a positive integer, and let $p_1 < p_2 < \dots < p_\ell$ be rational primes that are totally split in $K : \mathbb{Q}$. Suppose moreover that no p_i divides the discriminant of the minimal polynomial of α . For every $i \in \{1, \dots, \ell\}$ let $\mathfrak{p}_1^{(p_i)}, \dots, \mathfrak{p}_{r+1}^{(p_i)}$ be the prime ideals of \mathcal{O}_K that lie above p_i . Notice that $\mathbb{F}_{\mathfrak{p}_j^{(p_i)}} = \mathbb{F}_{p_i}$ for all i, j .

The number theoretical locally recoverable code $\mathcal{C} = \mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}})$ is defined as the image $\phi(\mathcal{A}[M])$ of the map

$$\begin{aligned} \phi : \mathcal{A}[M] &\longrightarrow \prod_{i=1}^{\ell} \prod_{j=1}^{r+1} \mathbb{F}_{\mathfrak{p}_j^{(p_i)}} = \prod_{i=1}^{\ell} \mathbb{F}_{p_i}^{r+1} \\ x &\mapsto (x \bmod \mathfrak{p}_j^{(p_i)})_{(i,j) \in U \times V} \end{aligned}$$

where $U = \{1, \dots, \ell\}$ and $V = \{1, \dots, r+1\}$. For simplicity of notation, let us define $c_j^{(i)} = x \bmod \mathfrak{p}_j^{(p_i)}$. Notice that

$$\mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}}) \subseteq \mathcal{D}\left(K, M^{s+1}, \{\mathfrak{p}_j^{(p_i)}\}_{i,j}\right).$$

Lemma 4.4. *Assume that*

$$\prod_{i \in \{1, \dots, \ell\}, j \in \{1, \dots, r+1\}} N(\mathfrak{p}_j^{(p_i)}) > C_\alpha \cdot (M^{s+1} - 1)^{r+1},$$

where C_α is the constant (4.1). Then the code $\mathcal{C} = \phi(\mathcal{A}[M])$ has size $\#\mathcal{A}[M] = M^{r(s+1)}$, i.e. ϕ is injective.

Proof. To see this, simply notice that $\mathcal{A}[M] \subseteq \mathcal{R}[M^{s+1}]$ and then apply Theorem 4.1. \square

Definition 4.5. Whenever the hypothesis of Lemma 4.4 are verified, we say that the code $\mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}})$ is a *good split code* of length $n = (r+1)\ell$ and size $M^{r(s+1)}$ over the number field K .

Proposition 4.6. Let $\mathcal{C} = \mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}})$ be a good split code. Then \mathcal{C} has locality r .

Proof. Suppose that the component $c_k^{(h)}$ of the codeword $c = (c_j^{(i)})_{j=1, \dots, r+1}^{i=1, \dots, \ell}$ has to be retrieved. Such codeword c arises from a message $m = \sum_{j=0}^s f_j M^j \in \mathcal{A}[M]$, where $f_j = \sum_{i=0}^{r-1} a_{i,j} \alpha^i \in \mathcal{R}[M]^-$. Now consider the components $c_1^{(h)}, \dots, c_{k-1}^{(h)}, c_{k+1}^{(h)}, \dots, c_{r+1}^{(h)}$ of the codeword c . Each of them arises as the reduction of m modulo $\mathfrak{p}_j^{(p_h)}$, for some $j \in \{1, \dots, k-1, k+1, \dots, r+1\}$. The key point is now the following: since p_h is totally split in K and it does not divide the discriminant of the minimal polynomial $m_\alpha(x)$ of α , by the Dedekind criterion we have that $m_\alpha(x) \bmod p_h = \prod_{i=1}^{r+1} (x - \beta_i) \in \mathbb{F}_{p_h}[x]$ where $\beta_1, \dots, \beta_{r+1} \in \mathbb{F}_{p_h}$ are pairwise distinct elements and β_j is the image of α via the reduction map $\mathcal{O}_K \rightarrow \mathcal{O}_K/\mathfrak{p}_j^{(p_h)} \cong \mathbb{F}_{p_h}$. Since $m = \sum_{i=0}^{r-1} u_i \alpha^i$ for some integers u_0, \dots, u_{r-1} , the component $c_j^{(h)}$ of c can be written as $\sum_{i=0}^{r-1} u_i \beta_j^i$. This gives us a system of linear equations in \mathbb{F}_{p_h} , whose indeterminates are the reductions $\tilde{u}_0, \dots, \tilde{u}_{r-1}$ of the u_i 's modulo p_h :

$$\begin{cases} \tilde{u}_0 + \tilde{u}_1 \beta_1 + \dots + \tilde{u}_{r-1} \beta_1^{r-1} = c_1^{(h)} \\ \tilde{u}_0 + \tilde{u}_1 \beta_2 + \dots + \tilde{u}_{r-1} \beta_2^{r-1} = c_2^{(h)} \\ \dots \\ \tilde{u}_0 + \tilde{u}_1 \beta_{r+1} + \dots + \tilde{u}_{r-1} \beta_{r+1}^{r-1} = c_{r+1}^{(h)} \end{cases}.$$

Notice that the k -th row is deleted, since $c_k^{(h)}$ is missing. This is a system of r equations in r indeterminates, and its determinant is non-zero because the β_i 's are pairwise distinct and the matrix representing the system is a Vandermonde matrix. If $(v_0, \dots, v_{r-1}) \in \mathbb{F}_{p_h}^r$ is its unique solution, then $c_k^{(h)} = \sum_{i=0}^{r-1} v_i \beta_k^i$. \square

Proposition 4.7. *Let $\mathcal{C} = \mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}})$ be a good split code, and let $\mathcal{P} = \{\mathfrak{p}_j^{(p_i)} : i \in \{1, \dots, \ell\}, j \in \{1, \dots, r+1\}\}$. Let*

$$m := \min_{T \subseteq \mathcal{P}} \left\{ \#T : \prod_{\mathfrak{p} \in T} N(\mathfrak{p}) > C_\alpha \cdot (M^{s+1} - 1)^{r+1} \right\}.$$

Then \mathcal{C} has minimum distance $d \geq (r+1)\ell - m + 1$.

Proof. The code \mathcal{C} is a subcode of the number theoretical Reed-Solomon Code with parameter M^{s+1} , so we simply apply again Theorem 4.1 with $n = (r+1)\ell$ (as our set of primes consists of $r+1$ primes on \mathcal{O}_K lying above each of the ℓ rational primes). \square

The following theorem summarizes what we proved until now

Theorem 4.8. *Let K be a number field of degree $r+1$, let s be a positive integer, let $M \in \mathbb{N}$, and let $\{p_i\}_{i \in \{1, \dots, \ell\}}$ be a set of rational primes that are totally split in K/\mathbb{Q} . Let $\mathcal{C} = \mathcal{C}(r, s, K, M, \{p_i\}_{i \in \{1, \dots, \ell\}})$ be a good split code over K . Then \mathcal{C} has length $\ell(r+1)$, size $M^{r(s+1)}$, minimum distance at least $(r+1)\ell - m + 1$, and locality r .*

Proof. Simply combine Lemma 4.4 and Propositions 4.6, and 4.7. \square

Example 4.9. Let us illustrate our construction with a working example. Let $K = \mathbb{Q}(\alpha)$, where $\alpha := \zeta_{16} + \zeta_{16}^{-1}$ and ζ_{16} is a primitive 16-th root of 1. The field K is the largest totally real subfield of $\mathbb{Q}(\zeta_{16})$, and the extension K/\mathbb{Q} is cyclic of degree 4 and has discriminant 2^{11} . The minimal polynomial of α is $x^4 - 4x^2 + 2$. The constant (4.1) is given by $C_\alpha = 4^2(1+4)^6 = 16 \cdot 5^6 = 250000$. Primes that split completely in K are exactly those congruent to ± 1 modulo 16. Let $p_1 = 17$, $p_2 = 31$ and $p_3 = 47$, so that $\ell = 3$. Let $\mathfrak{p}_1^{(p_1)}, \dots, \mathfrak{p}_4^{(p_i)}$ be the primes of K lying above p_i , for every i . Let $M = 2$ and $s = 3$. One computes that $17^4 \cdot 31^4 > C_\alpha(M^4 - 1)^4$, while $17^4 \cdot 31^3 < C_\alpha(M^4 - 1)^4$, so that $\mathcal{C} = \mathcal{C}(4, 3, K, 2, \{17, 31, 47\})$ is a good split code of size 2^{12} and minimum distance at least $12 - 8 + 1 = 5$ according to Theorem 4.1. We have that $\mathcal{R}[2]^- = \{a_0 + a_1\alpha + a_2\alpha^2 : a_i \in \{0, 1\}\}$ while $\mathcal{A}[2] = \{f_0 + f_1 \cdot 2 + f_2 \cdot 2^2 + f_3 \cdot 2^3 : f_i \in \mathcal{R}[2]^-\}$. To find the encoding map, notice that

$$x^4 - 4x^2 + 2 \equiv \begin{cases} (x+5)(x+8)(x+9)(x+12) & \text{mod } 17 \\ (x+5)(x+14)(x+17)(x+26) & \text{mod } 31, \\ (x+3)(x+18)(x+29)(x+44) & \text{mod } 47 \end{cases}$$

and therefore such map is given by

$$\begin{aligned} \phi: \mathcal{A}[M] &\rightarrow \mathbb{F}_{17}^4 \times \mathbb{F}_{31}^4 \times \mathbb{F}_{47}^4 \\ \alpha &\mapsto ([5, 8, 9, 12], [5, 14, 17, 26], [3, 18, 29, 44]). \end{aligned}$$

One can compute that the actual minimum distance of \mathcal{C} is 6, strictly better than the bound that comes from Theorem 4.1.

4.4. Almost good families of good split codes. We will now show how to construct an almost good family of good split codes, in the sense of equations (2.1) and (2.2). To do so, we first need the following analytic number theoretical lemma.

Lemma 4.10. *Let K be a Galois extension of \mathbb{Q} . For every $\ell \geq 1$, let p_1, \dots, p_ℓ be the first ℓ totally split primes of K/\mathbb{Q} . Then*

$$\log \left(\prod_{i=1}^{\ell} p_i \right) \sim \ell \log \ell$$

as ℓ tends to infinity.

Proof. First, let us recall that, if X is a positive integer, the totally split primes of K/\mathbb{Q} verify

$$(4.2) \quad \log \left(\prod_{\substack{p \leq X \\ \text{totally split}}} p \right) \sim \frac{X}{[K : \mathbb{Q}]}$$

In addition, the Chebotarev Density Theorem ensures that the asymptotic formula for the ℓ -th totally split prime is

$$p_\ell \sim \ell \log(\ell) [K : \mathbb{Q}].$$

By setting $X = p_\ell$ in (4.2) and relabeling the product we get

$$\log \left(\prod_{i=1}^{\ell} p_i \right) \sim \frac{\ell \log(\ell) [K : \mathbb{Q}]}{[K : \mathbb{Q}]} = \ell \log(\ell)$$

□

Remark 4.11. There are secondary terms in the asymptotic formula for p_ℓ , i.e. the ℓ -th (rational) totally split prime, and these secondary terms are larger than the secondary terms in the asymptotic formula for the ℓ -th rational prime.

Let now K be a number field of degree $r + 1$ that is Galois over \mathbb{Q} (this restriction allows for easier computations in Theorem 4.12), let $\alpha \in \mathcal{O}_K$ be a generator, let C_α be the constant (4.1) and let $s \in \mathbb{N}$. Let $\{p_j\}_{j \in \mathbb{N}}$ be the strictly increasing sequence of primes that are totally split in K/\mathbb{Q} , and for every $\ell \geq 1$ let $P_\ell := \prod_{i=1}^{\ell} p_i$.

Theorem 4.12. *Let $0 < c < 1$, let $k \in \mathbb{R}^+$ be such that $k < 1/\sqrt[r+1]{C_\alpha}$, and let $M_\ell := \left\lfloor s^{+1} \sqrt[k \cdot P_\ell / P_{\lfloor c\ell \rfloor}] \right\rfloor$. Let $\mathcal{C}_\ell := \mathcal{C}(r, s, K, M_\ell, \{p_i\}_{i \in \{1, \dots, \ell\}})$. Then $\{\mathcal{C}_\ell\}_{\ell \geq 1}$ is an almost good family of good split codes.*

Proof. To prove that \mathcal{C}_ℓ is a good split code, it is enough to show that

$$N \left(\prod_{p \in \mathcal{A}} \prod_{j=1}^{r+1} \mathfrak{p}_j^{(p_i)} \right) = P_\ell^{r+1} > C_\alpha (M_\ell^{s+1} - 1)^{r+1}$$

for some set of primes $\mathcal{A} \subseteq \{p_1, \dots, p_\ell\}$. This also shows that the code has distance at least $n - \#\mathcal{A} + 1$. We now show that if we choose \mathcal{A} to be the set of all primes of \mathcal{O}_K lying above $p_{\lfloor c\ell \rfloor + 1}, \dots, p_\ell$, the hypothesis of Proposition 4.7 are satisfied. By multiplying both sides of the inequality $1 > k \sqrt[r+1]{C_\alpha}$ by $P_\ell/P_{\lfloor c\ell \rfloor}$ we obtain that:

$$P_\ell/P_{\lfloor c\ell \rfloor} = \prod_{i=\lfloor c\ell \rfloor + 1}^{\ell} p_i > \sqrt[r+1]{C_\alpha} \frac{kP_\ell}{P_{\lfloor c\ell \rfloor}}.$$

By raising both sides to the $(r+1)$ -th power we get that, since $N(\mathfrak{p}_j^{(p_i)}) = p_i$ for all j 's,

$$\begin{aligned} N\left(\prod_{i=\lfloor c\ell \rfloor + 1}^{\ell} \prod_{j=1}^{r+1} \mathfrak{p}_j^{(p_i)}\right) &= \prod_{i=\lfloor c\ell \rfloor + 1}^{\ell} p_i^{r+1} > C_\alpha \left(\frac{kP_\ell}{P_{\lfloor c\ell \rfloor}}\right)^{r+1} \\ &\geq C_\alpha M_\ell^{(s+1)(r+1)} \\ &> C_\alpha (M_\ell^{s+1} - 1)^{r+1}, \end{aligned}$$

proving both that the code is good, that the distance tends to infinity as ℓ grows.

Next, we need to prove that the rate of \mathcal{C}_ℓ tends to a constant greater than zero, i.e. (2.2). Let $R_\ell := \prod_{i=1}^{\ell} \mathbb{F}_{p_i}^{r+1}$. Then

$$\begin{aligned} \liminf_{\ell \rightarrow +\infty} \frac{\log \#\mathcal{C}_\ell}{\log \#R_\ell} &= \liminf_{\ell \rightarrow +\infty} \frac{r(s+1) \log \left[\sqrt[s+1]{k \cdot P_\ell/P_{\lfloor c\ell \rfloor}} \right]}{(r+1) \log P_\ell} \\ &\geq \liminf_{\ell \rightarrow +\infty} \frac{r(s+1) \log \left(\sqrt[s+1]{k \cdot P_\ell/P_{\lfloor c\ell \rfloor}} - 1 \right)}{(r+1) \log P_\ell} \\ &\geq \liminf_{\ell \rightarrow +\infty} \frac{r \log (k \cdot P_\ell/P_{\lfloor c\ell \rfloor})}{(r+1) \log P_\ell} \end{aligned}$$

Now, using the properties of logarithms and the fact that $\log(P_\ell) \sim \ell \log \ell$ thanks to Lemma 4.10 we get

$$\begin{aligned} \liminf_{\ell \rightarrow +\infty} \frac{r \log (k \cdot P_\ell/P_{\lfloor c\ell \rfloor})}{(r+1) \log P_\ell} &= \liminf_{\ell \rightarrow +\infty} \frac{r(\log k + \ell \log \ell - c\ell \log c\ell)}{(r+1)\ell \log \ell} \\ &= \frac{r(1-c)}{r+1}, \end{aligned}$$

satisfying (2.1). \square

Remark 4.13. Notice that the distance grows linearly in ℓ (which is itself proportional to length and dimension), that is a desirable code property.

5. REALIZATION OF THE CONSTRUCTION

By the Kronecker-Weber theorem, one can always construct a Galois extension K/\mathbb{Q} of degree $r+1$ such that $\text{Gal}(K/\mathbb{Q})$ is cyclic of order $r+1$. This guarantees

that the construction is always feasible and the number of totally split places is “large” (as their density will be roughly asymptotic to $1/(r + 1)$).

The lemma that follows provides a constructive proof for the following curious (but expected) fact, for which we could not find reference in the literature: given a positive integer δ and n rational primes p_1, \dots, p_n larger than δ it is always possible to construct explicitly a number field of degree δ where p_1, \dots, p_n are all totally split. This shows that if one desires to construct a locally recoverable code over a certain fixed product of finite fields, this is in theory possible.

Lemma 5.1. *Let $\delta \in \mathbb{Z}_{>1}$ and let p_1, \dots, p_n be distinct rational primes all larger than δ . Then it is possible to explicitly construct a monic, irreducible polynomial $f(x) \in \mathbb{Z}[x]$ of degree δ such that if α is a root of f then all the p_i 's are totally split in the number field $\mathbb{Q}(\alpha)$.*

Proof. For each $i \in \{1, \dots, n\}$, choose $\alpha_1^i, \dots, \alpha_\delta^i \in \mathbb{Z}$ such that $\alpha_j^i \not\equiv \alpha_k^i \pmod{p_i}$ for every $j \neq k$ (this is possible because $p_i > \delta$). Next, choose a new prime p_{n+1} , different from p_1, \dots, p_n , and for every $i \in \{1, \dots, n + 1\}$ let $q_i := \prod_{j \neq i} p_j$. Notice that $q_1 + \dots + q_n$ is coprime with q_{n+1} , as if a prime p divides q_{n+1} then $p = p_i$ for some $i \in \{1, \dots, n\}$ and hence p divides q_j for every $j \in \{1, \dots, n\} \setminus \{i\}$; it follows that p does not divide $q_1 + \dots + q_n$. Hence there exist $u_1, u_2 \in \mathbb{Z}$ with $u_1 u_2 \neq 0$ such that $u_1(q_1 + \dots + q_n) + u_2 q_{n+1} = 1$. Notice that u_1 is coprime with $p_1 \cdot \dots \cdot p_n$ and u_2 is coprime with p_{n+1} . Now let $g(x) \in \mathbb{Z}[x]$ be a monic degree δ polynomial that is irreducible modulo p_{n+1} .

Consider then the polynomial

$$f(x) = u_1 \sum_{i=1}^n q_i \prod_{j=1}^{\delta} (x - \alpha_j^i) + u_2 q_{n+1} g(x) \in \mathbb{Z}[x].$$

By construction, $f(x)$ is monic. Moreover $f(x)$ is irreducible in $\mathbb{Z}[x]$ because $f(x)$ is irreducible modulo p_{n+1} (because its reduction is $g(x) \pmod{p_{n+1}}$, which is irreducible by construction), and hence the number field K generated by a root α of f has degree δ . It remains to show that p_1, \dots, p_n are totally split in K . But this follows immediately from Dedekind criterion, that can be applied because none of the p_i 's divide the discriminant of f since f has no multiple roots modulo any p_i . It follows that the factorization pattern of p_i in \mathcal{O}_K coincides with that of f modulo p_i ; by construction this is a product of δ distinct linear terms. \square

6. ACKNOWLEDGEMENTS

This work was supported in part by the National Science Foundation under Grant No 2127742.

REFERENCES

- [1] Alexander Barg, Kathryn Haymaker, Everett W Howe, Gretchen L Matthews, and Anthony Várilly-Alvarado. Locally recoverable codes from algebraic curves and surfaces. In *Algebraic Geometry for Coding Theory and Cryptography*, pages 95–127. Springer, 2017.
- [2] Daniele Bartoli, Maria Montanucci, and Luciane Quoos. Locally recoverable codes from automorphism group of function fields of genus $g \geq 1$. *IEEE Transactions on Information Theory*, 66(11):6799–6808, 2020.
- [3] Austin Dukes, Andrea Ferraguti, and Giacomo Micheli. Optimal selection for good polynomials of degree up to five. *Designs, Codes and Cryptography*, 90(6):1427–1436, 2022.
- [4] Ragnar Freij-Hollanti, Thomas Westerbäck, and Camilla Hollanti. Locally repairable codes with availability and hierarchy: matroid theory via examples. In *International Zurich Seminar on Communications-Proceedings*, pages 45–49. ETH Zurich, 2016.
- [5] Venkatesan Guruswami. Constructions of codes from number fields. *IEEE Trans. Inform. Theory*, 49(3):594–603, 2003.
- [6] H. W. Lenstra, Jr. Codes from algebraic number fields. In *Mathematics and computer science, II (Amsterdam, 1986)*, volume 4 of *CWI Monogr.*, pages 95–104. North-Holland, Amsterdam, 1986.
- [7] Jian Liu, Sihem Mesnager, and Lusheng Chen. New constructions of optimal locally recoverable codes via good polynomials. *IEEE Transactions on Information Theory*, 64(2):889–899, 2017.
- [8] Jian Liu, Sihem Mesnager, and Lusheng Chen. New constructions of optimal locally recoverable codes via good polynomials. *IEEE Transactions on Information Theory*, 64(2):889–899, 2018.
- [9] Giacomo Micheli. Construction of locally recoverable codes which are optimal. *IEEE transactions on information theory*, 66(1):167–175, 2020.
- [10] Natalia Silberstein, Ankit Singh Rawat, O Ozan Koyluoglu, and Sriram Vishwanath. Optimal locally repairable codes via rank-metric codes. In *2013 IEEE International Symposium on Information Theory*, pages 1819–1823. IEEE, 2013.
- [11] Itzhak Tamo and Alexander Barg. A family of optimal locally recoverable codes. *IEEE Transactions on Information Theory*, 60(8):4661–4676, 2014.

DICATAM, UNIVERSITY OF BRESCIA, VIA BRANZE 43, I-25123 BRESCIA
Email address: `andrea.ferraguti@unibs.it`

COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027 USA
Email address: `goldfeld@columbia.edu`

UNIVERSITY OF SOUTH FLORIDA, 4202 E FOWLER AVE, 33620 TAMPA, US.
Email address: `gmicheli@usf.edu`