

67. Let  $P_1$  have normal vector  $\mathbf{n}_1$ . Then  $\mathbf{n}_1 = \langle 3, 6, -3 \rangle$ ,  $\mathbf{n}_2 = \langle 4, -12, 8 \rangle$ ,  $\mathbf{n}_3 = \langle 3, -9, 6 \rangle$ ,  $\mathbf{n}_4 = \langle 1, 2, -1 \rangle$ . Now  $\mathbf{n}_1 = 3\mathbf{n}_4$ , so  $\mathbf{n}_1$  and  $\mathbf{n}_4$  are parallel, and hence  $P_1$  and  $P_4$  are parallel; similarly  $P_2$  and  $P_3$  are parallel because  $\mathbf{n}_2 = \frac{4}{3}\mathbf{n}_3$ . However,  $\mathbf{n}_1$  and  $\mathbf{n}_2$  are not parallel (so not all four planes are parallel). Notice that the point  $(2, 0, 0)$  lies on both  $P_1$  and  $P_4$ , so these two planes are identical. The point  $(\frac{5}{4}, 0, 0)$  lies on  $P_2$  but not on  $P_3$ , so these are different planes.
68. Let  $L_1$  have direction vector  $\mathbf{v}_1$ . Rewrite the symmetric equations for  $L_3$  as  $\frac{x-1}{1/2} = \frac{y-1}{-1/4} = \frac{z+1}{1}$ ; then  $\mathbf{v}_1 = \langle 6, -3, 12 \rangle$ ,  $\mathbf{v}_2 = \langle 2, 1, 4 \rangle$ ,  $\mathbf{v}_3 = \langle \frac{1}{2}, -\frac{1}{4}, 1 \rangle$ , and  $\mathbf{v}_4 = \langle 4, 2, 8 \rangle$ .  $\mathbf{v}_1 = 12\mathbf{v}_3$ , so  $L_1$  and  $L_3$  are parallel.  $\mathbf{v}_4 = 2\mathbf{v}_2$ , so  $L_2$  and  $L_4$  are parallel. (Note that  $L_1$  and  $L_2$  are not parallel.)  $L_1$  contains the point  $(1, 1, 5)$ , but this point does not lie on  $L_3$ , so they're not identical.  $(3, 1, 5)$  lies on  $L_4$  and also on  $L_2$  (for  $t = 1$ ), so  $L_2$  and  $L_4$  are the same line.
69. Let  $Q = (1, 3, 4)$  and  $R = (2, 1, 1)$ , points on the line corresponding to  $t = 0$  and  $t = 1$ . Let  $P = (4, 1, -2)$ . Then  $\mathbf{a} = \overrightarrow{QR} = \langle 1, -2, -3 \rangle$ ,  $\mathbf{b} = \overrightarrow{QP} = \langle 3, -2, -6 \rangle$ . The distance is
- $$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 1, -2, -3 \rangle \times \langle 3, -2, -6 \rangle|}{|\langle 1, -2, -3 \rangle|} = \frac{|\langle 6, -3, 4 \rangle|}{|\langle 1, -2, -3 \rangle|} = \frac{\sqrt{6^2 + (-3)^2 + 4^2}}{\sqrt{1^2 + (-2)^2 + (-3)^2}} = \frac{\sqrt{61}}{\sqrt{14}} = \sqrt{\frac{61}{14}}.$$
70. Let  $Q = (0, 6, 3)$  and  $R = (2, 4, 4)$ , points on the line corresponding to  $t = 0$  and  $t = 1$ . Let  $P = (0, 1, 3)$ . Then  $\mathbf{a} = \overrightarrow{QR} = \langle 2, -2, 1 \rangle$  and  $\mathbf{b} = \overrightarrow{QP} = \langle 0, -5, 0 \rangle$ . The distance is
- $$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 2, -2, 1 \rangle \times \langle 0, -5, 0 \rangle|}{|\langle 2, -2, 1 \rangle|} = \frac{|\langle 5, 0, -10 \rangle|}{|\langle 2, -2, 1 \rangle|} = \frac{\sqrt{5^2 + 0^2 + (-10)^2}}{\sqrt{2^2 + (-2)^2 + 1^2}} = \frac{\sqrt{125}}{\sqrt{9}} = \frac{5\sqrt{5}}{3}.$$
71. By Equation 9, the distance is  $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|3(1) + 2(-2) + 6(4) - 5|}{\sqrt{3^2 + 2^2 + 6^2}} = \frac{|18|}{\sqrt{49}} = \frac{18}{7}.$
72. By Equation 9, the distance is  $D = \frac{|1(-6) - 2(3) - 4(5) - 8|}{\sqrt{1^2 + (-2)^2 + (-4)^2}} = \frac{|-40|}{\sqrt{21}} = \frac{40}{\sqrt{21}}.$
73. Put  $y = z = 0$  in the equation of the first plane to get the point  $(2, 0, 0)$  on the plane. Because the planes are parallel, the distance  $D$  between them is the distance from  $(2, 0, 0)$  to the second plane. By Equation 9,
- $$D = \frac{|4(2) - 6(0) + 2(0) - 3|}{\sqrt{4^2 + (-6)^2 + (2)^2}} = \frac{5}{\sqrt{56}} = \frac{5}{2\sqrt{14}} \text{ or } \frac{5\sqrt{14}}{28}.$$
74. Put  $x = y = 0$  in the equation of the first plane to get the point  $(0, 0, 0)$  on the plane. Because the planes are parallel the distance  $D$  between them is the distance from  $(0, 0, 0)$  to the second plane  $3x - 6y + 9z - 1 = 0$ . By Equation 9,
- $$D = \frac{|3(0) - 6(0) + 9(0) - 1|}{\sqrt{3^2 + (-6)^2 + 9^2}} = \frac{1}{\sqrt{126}} = \frac{1}{3\sqrt{14}}.$$
75. The distance between two parallel planes is the same as the distance between a point on one of the planes and the other plane. Let  $P_0 = (x_0, y_0, z_0)$  be a point on the plane given by  $ax + by + cz + d_1 = 0$ . Then  $ax_0 + by_0 + cz_0 + d_1 = 0$  and the

67. Let  $P_i$  have normal vector  $\mathbf{n}_i$ . Then  $\mathbf{n}_1 = \langle 3, 6, -3 \rangle$ ,  $\mathbf{n}_2 = \langle 4, -12, 8 \rangle$ ,  $\mathbf{n}_3 = \langle 3, -9, 6 \rangle$ ,  $\mathbf{n}_4 = \langle 1, 2, -1 \rangle$ . Now  $\mathbf{n}_1 = 3\mathbf{n}_4$ , so  $\mathbf{n}_1$  and  $\mathbf{n}_4$  are parallel, and hence  $P_1$  and  $P_4$  are parallel; similarly  $P_2$  and  $P_3$  are parallel because  $\mathbf{n}_2 = \frac{4}{3}\mathbf{n}_3$ . However,  $\mathbf{n}_1$  and  $\mathbf{n}_2$  are not parallel (so not all four planes are parallel). Notice that the point  $(2, 0, 0)$  lies on both  $P_1$  and  $P_4$ , so these two planes are identical. The point  $(\frac{5}{4}, 0, 0)$  lies on  $P_2$  but not on  $P_3$ , so these are different planes.
68. Let  $L_i$  have direction vector  $\mathbf{v}_i$ . Rewrite the symmetric equations for  $L_3$  as  $\frac{x-1}{1/2} = \frac{y-1}{-1/4} = \frac{z+1}{1}$ ; then  $\mathbf{v}_1 = \langle 6, -3, 12 \rangle$ ,  $\mathbf{v}_2 = \langle 2, 1, 4 \rangle$ ,  $\mathbf{v}_3 = \langle \frac{1}{2}, -\frac{1}{4}, 1 \rangle$ , and  $\mathbf{v}_4 = \langle 4, 2, 8 \rangle$ .  $\mathbf{v}_1 = 12\mathbf{v}_3$ , so  $L_1$  and  $L_3$  are parallel.  $\mathbf{v}_4 = 2\mathbf{v}_2$ , so  $L_2$  and  $L_4$  are parallel. (Note that  $L_1$  and  $L_2$  are not parallel.)  $L_1$  contains the point  $(1, 1, 5)$ , but this point does not lie on  $L_3$ , so they're not identical.  $(3, 1, 5)$  lies on  $L_4$  and also on  $L_2$  (for  $t = 1$ ), so  $L_2$  and  $L_4$  are the same line.
69. Let  $Q = (1, 3, 4)$  and  $R = (2, 1, 1)$ , points on the line corresponding to  $t = 0$  and  $t = 1$ . Let  $P = (4, 1, -2)$ . Then  $\mathbf{a} = \overrightarrow{QR} = \langle 1, -2, -3 \rangle$ ,  $\mathbf{b} = \overrightarrow{QP} = \langle 3, -2, -6 \rangle$ . The distance is
- $$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 1, -2, -3 \rangle \times \langle 3, -2, -6 \rangle|}{|\langle 1, -2, -3 \rangle|} = \frac{|\langle 6, -3, 4 \rangle|}{|\langle 1, -2, -3 \rangle|} = \frac{\sqrt{6^2 + (-3)^2 + 4^2}}{\sqrt{1^2 + (-2)^2 + (-3)^2}} = \frac{\sqrt{61}}{\sqrt{14}} = \sqrt{\frac{61}{14}}.$$
70. Let  $Q = (0, 6, 3)$  and  $R = (2, 4, 4)$ , points on the line corresponding to  $t = 0$  and  $t = 1$ . Let  $P = (0, 1, 3)$ . Then  $\mathbf{a} = \overrightarrow{QR} = \langle 2, -2, 1 \rangle$  and  $\mathbf{b} = \overrightarrow{QP} = \langle 0, -5, 0 \rangle$ . The distance is
- $$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 2, -2, 1 \rangle \times \langle 0, -5, 0 \rangle|}{|\langle 2, -2, 1 \rangle|} = \frac{|\langle 5, 0, -10 \rangle|}{|\langle 2, -2, 1 \rangle|} = \frac{\sqrt{5^2 + 0^2 + (-10)^2}}{\sqrt{2^2 + (-2)^2 + 1^2}} = \frac{\sqrt{125}}{\sqrt{9}} = \frac{5\sqrt{5}}{3}.$$
71. By Equation 9, the distance is  $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|3(1) + 2(-2) + 6(4) - 5|}{\sqrt{3^2 + 2^2 + 6^2}} = \frac{|18|}{\sqrt{49}} = \frac{18}{7}.$
72. By Equation 9, the distance is  $D = \frac{|1(-6) - 2(3) - 4(5) - 8|}{\sqrt{1^2 + (-2)^2 + (-4)^2}} = \frac{|-40|}{\sqrt{21}} = \frac{40}{\sqrt{21}}.$
73. Put  $y = z = 0$  in the equation of the first plane to get the point  $(2, 0, 0)$  on the plane. Because the planes are parallel, the distance  $D$  between them is the distance from  $(2, 0, 0)$  to the second plane. By Equation 9,
- $$D = \frac{|4(2) - 6(0) + 2(0) - 3|}{\sqrt{4^2 + (-6)^2 + (2)^2}} = \frac{5}{\sqrt{56}} = \frac{5}{2\sqrt{14}} \text{ or } \frac{5\sqrt{14}}{28}.$$
74. Put  $x = y = 0$  in the equation of the first plane to get the point  $(0, 0, 0)$  on the plane. Because the planes are parallel the distance  $D$  between them is the distance from  $(0, 0, 0)$  to the second plane  $3x - 6y + 9z - 1 = 0$ . By Equation 9,
- $$D = \frac{|3(0) - 6(0) + 9(0) - 1|}{\sqrt{3^2 + (-6)^2 + 9^2}} = \frac{1}{\sqrt{126}} = \frac{1}{3\sqrt{14}}.$$
75. The distance between two parallel planes is the same as the distance between a point on one of the planes and the other plane. Let  $P_0 = (x_0, y_0, z_0)$  be a point on the plane given by  $ax + by + cz + d_1 = 0$ . Then  $ax_0 + by_0 + cz_0 + d_1 = 0$  and the



distance between  $P_0$  and the plane given by  $ax + by + cz + d_2 = 0$  is, from Equation 9,

$$D = \frac{|ax_0 + by_0 + cz_0 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|-d_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}.$$

76. The planes must have parallel normal vectors, so if  $ax + by + cz + d = 0$  is such a plane, then for some  $t \neq 0$ ,

$\langle a, b, c \rangle = t\langle 1, 2, -2 \rangle = \langle t, 2t, -2t \rangle$ . So this plane is given by the equation  $x + 2y - 2z + k = 0$ , where  $k = d/t$ . By

Exercise 75, the distance between the planes is  $2 = \frac{|1 - k|}{\sqrt{1^2 + 2^2 + (-2)^2}} \Leftrightarrow 6 = |1 - k| \Leftrightarrow k = 7 \text{ or } -5$ . So the desired planes have equations  $x + 2y - 2z = 7$  and  $x + 2y - 2z = -5$ .

77.  $L_1: x = y = z \Rightarrow x = y$  (1).  $L_2: x + 1 = y/2 = z/3 \Rightarrow x + 1 = y/2$  (2). The solution of (1) and (2) is  $x = y = -2$ . However, when  $x = -2$ ,  $x = z \Rightarrow z = -2$ , but  $x + 1 = z/3 \Rightarrow z = -3$ , a contradiction. Hence the lines do not intersect. For  $L_1$ ,  $\mathbf{v}_1 = \langle 1, 1, 1 \rangle$ , and for  $L_2$ ,  $\mathbf{v}_2 = \langle 1, 2, 3 \rangle$ , so the lines are not parallel. Thus the lines are skew lines. If two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both  $\langle 1, 1, 1 \rangle$  and  $\langle 1, 2, 3 \rangle$ , the direction vectors of the two lines. So set  $\mathbf{n} = \langle 1, 1, 1 \rangle \times \langle 1, 2, 3 \rangle = \langle 3 - 2, -3 + 1, 2 - 1 \rangle = \langle 1, -2, 1 \rangle$ . From above, we know that  $(-2, -2, -2)$  and  $(-2, -2, -3)$  are points of  $L_1$  and  $L_2$  respectively. So in the notation of Equation 8,  $1(-2) - 2(-2) + 1(-2) + d_1 = 0 \Rightarrow d_1 = 0$  and  $1(-2) - 2(-2) + 1(-3) + d_2 = 0 \Rightarrow d_2 = 1$ .

By Exercise 75, the distance between these two skew lines is  $D = \frac{|0 - 1|}{\sqrt{1 + 4 + 1}} = \frac{1}{\sqrt{6}}$ .

*Alternate solution (without reference to planes):* A vector which is perpendicular to both of the lines is

$\mathbf{n} = \langle 1, 1, 1 \rangle \times \langle 1, 2, 3 \rangle = \langle 1, -2, 1 \rangle$ . Pick any point on each of the lines, say  $(-2, -2, -2)$  and  $(-2, -2, -3)$ , and form the vector  $\mathbf{b} = \langle 0, 0, 1 \rangle$  connecting the two points. The distance between the two skew lines is the absolute value of the scalar

projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,  $D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{|1 \cdot 0 - 2 \cdot 0 + 1 \cdot 1|}{\sqrt{1 + 4 + 1}} = \frac{1}{\sqrt{6}}$ .

78. First notice that if two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both  $\mathbf{v}_1 = \langle 1, 6, 2 \rangle$  and  $\mathbf{v}_2 = \langle 2, 15, 6 \rangle$ , the direction vectors of the two lines respectively. Thus set  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 36 - 30, 4 - 6, 15 - 12 \rangle = \langle 6, -2, 3 \rangle$ . Setting  $t = 0$  and  $s = 0$  gives the points  $(1, 1, 0)$  and  $(1, 5, -2)$ . So in the notation of Equation 8,  $6 - 2 + 0 + d_1 = 0 \Rightarrow d_1 = -4$  and  $6 - 10 - 6 + d_2 = 0 \Rightarrow d_2 = 10$ .

Then by Exercise 75, the distance between the two skew lines is given by  $D = \frac{|-4 - 10|}{\sqrt{36 + 4 + 9}} = \frac{14}{7} = 2$ .

*Alternate solution (without reference to planes):* We already know that the direction vectors of the two lines are

$\mathbf{v}_1 = \langle 1, 6, 2 \rangle$  and  $\mathbf{v}_2 = \langle 2, 15, 6 \rangle$ . Then  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 6, -2, 3 \rangle$  is perpendicular to both lines. Pick any point on each of the lines, say  $(1, 1, 0)$  and  $(1, 5, -2)$ , and form the vector  $\mathbf{b} = \langle 0, 4, -2 \rangle$  connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,

$$D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{1}{\sqrt{36 + 4 + 9}} |0 - 8 - 6| = \frac{14}{7} = 2.$$

distance between  $P_0$  and the plane given by  $ax + by + cz + d_2 = 0$  is, from Equation 9,

$$D = \frac{|ax_0 + by_0 + cz_0 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|-d_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}.$$

76. The planes must have parallel normal vectors, so if  $ax + by + cz + d = 0$  is such a plane, then for some  $t \neq 0$ ,  $\langle a, b, c \rangle = t\langle 1, 2, -2 \rangle = \langle t, 2t, -2t \rangle$ . So this plane is given by the equation  $x + 2y - 2z + k = 0$ , where  $k = d/t$ . By

Exercise 75, the distance between the planes is  $2 = \frac{|1 - k|}{\sqrt{1^2 + 2^2 + (-2)^2}} \Leftrightarrow 6 = |1 - k| \Leftrightarrow k = 7 \text{ or } -5$ . So the desired planes have equations  $x + 2y - 2z = 7$  and  $x + 2y - 2z = -5$ .

77.  $L_1: x = y = z \Rightarrow x = y$  (1).  $L_2: x + 1 = y/2 = z/3 \Rightarrow x + 1 = y/2$  (2). The solution of (1) and (2) is  $x = y = -2$ . However, when  $x = -2$ ,  $x = z \Rightarrow z = -2$ , but  $x + 1 = z/3 \Rightarrow z = -3$ , a contradiction. Hence the lines do not intersect. For  $L_1$ ,  $\mathbf{v}_1 = \langle 1, 1, 1 \rangle$ , and for  $L_2$ ,  $\mathbf{v}_2 = \langle 1, 2, 3 \rangle$ , so the lines are not parallel. Thus the lines are skew lines. If two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both  $\langle 1, 1, 1 \rangle$  and  $\langle 1, 2, 3 \rangle$ , the direction vectors of the two lines. So set  $\mathbf{n} = \langle 1, 1, 1 \rangle \times \langle 1, 2, 3 \rangle = \langle 3 - 2, -3 + 1, 2 - 1 \rangle = \langle 1, -2, 1 \rangle$ . From above, we know that  $(-2, -2, -2)$  and  $(-2, -2, -3)$  are points of  $L_1$  and  $L_2$  respectively. So in the notation of Equation 8,  $1(-2) - 2(-2) + 1(-2) + d_1 = 0 \Rightarrow d_1 = 0$  and  $1(-2) - 2(-2) + 1(-3) + d_2 = 0 \Rightarrow d_2 = 1$ .

By Exercise 75, the distance between these two skew lines is  $D = \frac{|0 - 1|}{\sqrt{1 + 4 + 1}} = \frac{1}{\sqrt{6}}$ .

*Alternate solution (without reference to planes):* A vector which is perpendicular to both of the lines is

$\mathbf{n} = \langle 1, 1, 1 \rangle \times \langle 1, 2, 3 \rangle = \langle 1, -2, 1 \rangle$ . Pick any point on each of the lines, say  $(-2, -2, -2)$  and  $(-2, -2, -3)$ , and form the vector  $\mathbf{b} = \langle 0, 0, 1 \rangle$  connecting the two points. The distance between the two skew lines is the absolute value of the scalar

projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,  $D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{|1 \cdot 0 - 2 \cdot 0 + 1 \cdot 1|}{\sqrt{1 + 4 + 1}} = \frac{1}{\sqrt{6}}$ .

78. First notice that if two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both  $\mathbf{v}_1 = \langle 1, 6, 2 \rangle$  and  $\mathbf{v}_2 = \langle 2, 15, 6 \rangle$ , the direction vectors of the two lines respectively. Thus set  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 36 - 30, 4 - 6, 15 - 12 \rangle = \langle 6, -2, 3 \rangle$ . Setting  $t = 0$  and  $s = 0$  gives the points  $(1, 1, 0)$  and  $(1, 5, -2)$ . So in the notation of Equation 8,  $6 - 2 + 0 + d_1 = 0 \Rightarrow d_1 = -4$  and  $6 - 10 - 6 + d_2 = 0 \Rightarrow d_2 = 10$ .

Then by Exercise 75, the distance between the two skew lines is given by  $D = \frac{|-4 - 10|}{\sqrt{36 + 4 + 9}} = \frac{14}{7} = 2$ .

*Alternate solution (without reference to planes):* We already know that the direction vectors of the two lines are

$\mathbf{v}_1 = \langle 1, 6, 2 \rangle$  and  $\mathbf{v}_2 = \langle 2, 15, 6 \rangle$ . Then  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 6, -2, 3 \rangle$  is perpendicular to both lines. Pick any point on each of the lines, say  $(1, 1, 0)$  and  $(1, 5, -2)$ , and form the vector  $\mathbf{b} = \langle 0, 4, -2 \rangle$  connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,

$$D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{1}{\sqrt{36 + 4 + 9}} |0 - 8 - 6| = \frac{14}{7} = 2.$$



79. A direction vector for  $L_1$  is  $\mathbf{v}_1 = \langle 2, 0, -1 \rangle$  and a direction vector for  $L_2$  is  $\mathbf{v}_2 = \langle 3, 2, 2 \rangle$ . These vectors are not parallel so neither are the lines. Parametric equations for the lines are  $L_1: x = 2t, y = 0, z = -t$ , and  $L_2: x = 1 + 3s, y = -1 + 2s, z = 1 + 2s$ . No values of  $t$  and  $s$  satisfy these equations simultaneously, so the lines don't intersect and hence are skew. We can view the lines as lying in two parallel planes; a common normal vector to the planes is  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4 \rangle$ . Line  $L_1$  passes through the origin, so  $(0, 0, 0)$  lies on one of the planes, and  $(1, -1, 1)$  is a point on  $L_2$  and therefore on the other plane. Equations of the planes then are  $2x - 7y + 4z = 0$  and  $2x - 7y + 4z - 13 = 0$ , and by Exercise 75, the distance between the two skew lines is  $D = \frac{|0 - (-13)|}{\sqrt{4 + 49 + 16}} = \frac{13}{\sqrt{69}}$ .

*Alternate solution (without reference to planes):* Direction vectors of the two lines are  $\mathbf{v}_1 = \langle 2, 0, -1 \rangle$  and  $\mathbf{v}_2 = \langle 3, 2, 2 \rangle$ . Then  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4 \rangle$  is perpendicular to both lines. Pick any point on each of the lines, say  $(0, 0, 0)$  and  $(1, -1, 1)$ , and form the vector  $\mathbf{b} = \langle 1, -1, 1 \rangle$  connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,  $D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{|2 + 7 + 4|}{\sqrt{4 + 49 + 16}} = \frac{13}{\sqrt{69}}$ .

80. A direction vector for the line  $L_1$  is  $\mathbf{v}_1 = \langle 1, 2, 2 \rangle$ . A normal vector for the plane  $P_1$  is  $\mathbf{n}_1 = \langle 1, -1, 2 \rangle$ . The vector from the point  $(0, 0, 1)$  to  $(3, 2, -1)$ ,  $\langle 3, 2, -2 \rangle$ , is parallel to the plane  $P_2$ , as is the vector from  $(0, 0, 1)$  to  $(1, 2, 1)$ , namely  $\langle 1, 2, 0 \rangle$ . Thus a normal vector for  $P_2$  is  $\langle 3, 2, -2 \rangle \times \langle 1, 2, 0 \rangle = \langle 4, -2, 4 \rangle$ , or we can use  $\mathbf{n}_2 = \langle 2, -1, 2 \rangle$ , and a direction vector for the line  $L_2$  of intersection of these planes is  $\mathbf{v}_2 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -1, 2 \rangle \times \langle 2, -1, 2 \rangle = \langle 0, 2, 1 \rangle$ . Notice that the point  $(3, 2, -1)$  lies on both planes, so it also lies on  $L_2$ . The lines are skew, so we can view them as lying in two parallel planes; a common normal vector to the planes is  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2 \rangle$ . Line  $L_1$  passes through the point  $(1, 2, 6)$ , so  $(1, 2, 6)$  lies on one of the planes, and  $(3, 2, -1)$  is a point on  $L_2$  and therefore on the other plane. Equations of the planes then are  $-2x - y + 2z - 8 = 0$  and  $-2x - y + 2z + 10 = 0$ , and by Exercise 75, the distance between the lines is

$$D = \frac{|-8 - 10|}{\sqrt{4 + 1 + 4}} = \frac{18}{3} = 6.$$

Alternatively, direction vectors for the lines are  $\mathbf{v}_1 = \langle 1, 2, 2 \rangle$  and  $\mathbf{v}_2 = \langle 0, 2, 1 \rangle$ , so  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2 \rangle$  is perpendicular to both lines. Pick any point on each of the lines, say  $(1, 2, 6)$  and  $(3, 2, -1)$ , and form the vector  $\mathbf{b} = \langle 2, 0, -7 \rangle$  connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of  $\mathbf{b}$  along  $\mathbf{n}$ , that is,  $D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{|-4 + 0 - 14|}{\sqrt{4 + 1 + 4}} = \frac{18}{3} = 6$ .

81. (a) A direction vector from tank A to tank B is  $\langle 765 - 325, 675 - 810, 599 - 561 \rangle = \langle 440, -135, 38 \rangle$ . Taking tank A's position  $(325, 810, 561)$  as the initial point, parametric equations for the line of sight are  $x = 325 + 440t$ ,  $y = 810 - 135t$ ,  $z = 561 + 38t$  for  $0 \leq t \leq 1$ .

56. (a) Rotation around  $\theta = \frac{\pi}{2}$  is the same as rotation around the  $y$ -axis, that is,  $S = \int_a^b 2\pi x \, ds$  where

$ds = \sqrt{(dx/dt)^2 + (dy/dt)^2} \, dt$  for a parametric equation, and for the special case of a polar equation,  $x = r \cos \theta$  and

$ds = \sqrt{(dx/d\theta)^2 + (dy/d\theta)^2} \, d\theta = \sqrt{r^2 + (dr/d\theta)^2} \, d\theta$  [see the derivation of Equation 10.4.5]. Therefore, for a polar

equation rotated around  $\theta = \frac{\pi}{2}$ ,  $S = \int_a^b 2\pi r \cos \theta \sqrt{r^2 + (dr/d\theta)^2} \, d\theta$ .


- (b) As in the solution for Exercise 55(b), we can double the surface area generated by rotating the curve from  $\theta = 0$  to  $\theta = \frac{\pi}{4}$  to obtain the total surface area.

$$\begin{aligned} S &= 2 \int_0^{\pi/4} 2\pi \sqrt{\cos 2\theta} \cos \theta \sqrt{\cos^2 2\theta + (\sin^2 2\theta)/\cos 2\theta} \, d\theta = 4\pi \int_0^{\pi/4} \sqrt{\cos 2\theta} \cos \theta \sqrt{\frac{\cos^2 2\theta + \sin^2 2\theta}{\cos 2\theta}} \, d\theta \\ &= 4\pi \int_0^{\pi/4} \sqrt{\cos 2\theta} \cos \theta \frac{1}{\sqrt{\cos 2\theta}} \, d\theta = 4\pi \int_0^{\pi/4} \cos \theta \, d\theta = 4\pi [\sin \theta]_0^{\pi/4} = 4\pi \left( \frac{\sqrt{2}}{2} - 0 \right) = 2\sqrt{2}\pi \end{aligned}$$

## 10.5 Conic Sections


1.  $x^2 = 6y$  and  $x^2 = 4py \Rightarrow 4p = 6 \Rightarrow p = \frac{3}{2}$ .

The vertex is  $(0, 0)$ , the focus is  $(0, \frac{3}{2})$ , and the directrix is  $y = -\frac{3}{2}$ .




2.  $2y^2 = 5x \Rightarrow y^2 = \frac{5}{2}x$ .  $4p = \frac{5}{2} \Rightarrow p = \frac{5}{8}$ .

The vertex is  $(0, 0)$ , the focus is  $(\frac{5}{8}, 0)$ , and the directrix is  $x = -\frac{5}{8}$ .




3.  $2x = -y^2 \Rightarrow y^2 = -2x$ .  $4p = -2 \Rightarrow p = -\frac{1}{2}$ .

The vertex is  $(0, 0)$ , the focus is  $(-\frac{1}{2}, 0)$ , and the directrix is  $x = \frac{1}{2}$ .




4.  $3x^2 + 8y = 0 \Rightarrow 3x^2 = -8y \Rightarrow x^2 = -\frac{8}{3}y$ .

$4p = -\frac{8}{3} \Rightarrow p = -\frac{2}{3}$ . The vertex is  $(0, 0)$ , the focus is  $(0, -\frac{2}{3})$ , and the directrix is  $y = \frac{2}{3}$ .







5.  $(x + 2)^2 = 8(y - 3)$ .  $4p = 8$ , so  $p = 2$ . The vertex is  $(-2, 3)$ , the focus is  $(-2, 5)$ , and the directrix is  $y = 1$ .




6.  $(y - 2)^2 = 2x + 1 = 2(x + \frac{1}{2})$ .  $4p = 2$ , so  $p = \frac{1}{2}$ . The vertex is  $(-\frac{1}{2}, 2)$ , the focus is  $(0, 2)$ , and the directrix is  $x = -1$ .





7.  $y^2 + 6y + 2x + 1 = 0 \Leftrightarrow y^2 + 6y = -2x - 1$   
 $\Leftrightarrow y^2 + 6y + 9 = -2x + 8 \Leftrightarrow$   
 $(y + 3)^2 = -2(x - 4)$ .  $4p = -2$ , so  $p = -\frac{1}{2}$ .  
 The vertex is  $(4, -3)$ , the focus is  $(\frac{7}{2}, -3)$ , and the directrix is  $x = \frac{9}{2}$ .



8.  $2x^2 - 16x - 3y + 38 = 0 \Leftrightarrow 2x^2 - 16x = 3y - 38$   
 $\Leftrightarrow 2(x^2 - 8x + 16) = 3y - 38 + 32 \Leftrightarrow$   
 $2(x - 4)^2 = 3y - 6 \Leftrightarrow (x - 4)^2 = \frac{3}{2}(y - 2)$ .  
 $4p = \frac{3}{2}$ , so  $p = \frac{3}{8}$ . The vertex is  $(4, 2)$ , the focus is  $(4, \frac{19}{8})$ , and the directrix is  $y = \frac{13}{8}$ .



9. The equation has the form  $y^2 = 4px$ , where  $p < 0$ . Since the parabola passes through  $(-1, 1)$ , we have  $1^2 = 4p(-1)$ , so  $4p = -1$  and an equation is  $y^2 = -x$  or  $x = -y^2$ .  $4p = -1$ , so  $p = -\frac{1}{4}$  and the focus is  $(-\frac{1}{4}, 0)$  while the directrix is  $x = \frac{1}{4}$ .
10. The vertex is  $(2, -2)$ , so the equation is of the form  $(x - 2)^2 = 4p(y + 2)$ , where  $p > 0$ . The point  $(0, 0)$  is on the parabola, so  $4 = 4p(2)$  and  $4p = 2$ . Thus, an equation is  $(x - 2)^2 = 2(y + 2)$ .  $4p = 2$ , so  $p = \frac{1}{2}$  and the focus is  $(2, -\frac{3}{2})$  while the directrix is  $y = -\frac{5}{2}$ .
11.  $\frac{x^2}{2} + \frac{y^2}{4} = 1 \Rightarrow a = \sqrt{4} = 2, b = \sqrt{2}$ ,  
 $c = \sqrt{a^2 - b^2} = \sqrt{4 - 2} = \sqrt{2}$ . The ellipse is centered at  $(0, 0)$ , with vertices at  $(0, \pm 2)$ . The foci are  $(0, \pm\sqrt{2})$ .
12.  $\frac{x^2}{36} + \frac{y^2}{8} = 1 \Rightarrow a = \sqrt{36} = 6, b = \sqrt{8}$ ,  
 $c = \sqrt{a^2 - b^2} = \sqrt{36 - 8} = \sqrt{28} = 2\sqrt{7}$ . The ellipse is centered at  $(0, 0)$ , with vertices at  $(\pm 6, 0)$ . The foci are  $(\pm 2\sqrt{7}, 0)$ .




$$13. x^2 + 9y^2 = 9 \Leftrightarrow \frac{x^2}{9} + \frac{y^2}{1} = 1 \Rightarrow a = \sqrt{9} = 3,$$

$$b = \sqrt{1} = 1, c = \sqrt{a^2 - b^2} = \sqrt{9 - 1} = \sqrt{8} = 2\sqrt{2}.$$

The ellipse is centered at  $(0, 0)$ , with vertices  $(\pm 3, 0)$ .


The foci are  $(\pm 2\sqrt{2}, 0)$ .



$$14. 100x^2 + 36y^2 = 225 \Leftrightarrow \frac{x^2}{\frac{225}{100}} + \frac{y^2}{\frac{225}{36}} = 1 \Leftrightarrow$$

$$\frac{x^2}{\frac{9}{4}} + \frac{y^2}{\frac{25}{4}} = 1 \Rightarrow a = \sqrt{\frac{25}{4}} = \frac{5}{2}, b = \sqrt{\frac{9}{4}} = \frac{3}{2},$$

$c = \sqrt{a^2 - b^2} = \sqrt{\frac{25}{4} - \frac{9}{4}} = 2$ . The ellipse is centered at  $(0, 0)$ , with vertices  $(0, \pm \frac{5}{2})$ . The foci are  $(0, \pm 2)$ .




$$15. 9x^2 - 18x + 4y^2 = 27 \Leftrightarrow$$

$$9(x^2 - 2x + 1) + 4y^2 = 27 + 9 \Leftrightarrow$$

$$9(x - 1)^2 + 4y^2 = 36 \Leftrightarrow \frac{(x - 1)^2}{4} + \frac{y^2}{9} = 1 \Rightarrow$$

$$a = 3, b = 2, c = \sqrt{5} \Rightarrow \text{center } (1, 0),$$

vertices  $(1, \pm 3)$ , foci  $(1, \pm \sqrt{5})$



$$16. x^2 + 3y^2 + 2x - 12y + 10 = 0 \Leftrightarrow$$


$$x^2 + 2x + 1 + 3(y^2 - 4y + 4) = -10 + 1 + 12 \Leftrightarrow$$

$$(x + 1)^2 + 3(y - 2)^2 = 3 \Leftrightarrow$$

$$\frac{(x + 1)^2}{3} + \frac{(y - 2)^2}{1} = 1 \Rightarrow a = \sqrt{3}, b = 1,$$

$$c = \sqrt{2} \Rightarrow \text{center } (-1, 2), \text{ vertices } (-1 \pm \sqrt{3}, 2),$$

foci  $(-1 \pm \sqrt{2}, 2)$



$$17. \text{The center is } (0, 0), a = 3, \text{ and } b = 2, \text{ so an equation is } \frac{x^2}{4} + \frac{y^2}{9} = 1. c = \sqrt{a^2 - b^2} = \sqrt{5}, \text{ so the foci are } (0, \pm \sqrt{5}).$$


$$18. \text{The ellipse is centered at } (2, 1), \text{ with } a = 3 \text{ and } b = 2. \text{ An equation is } \frac{(x - 2)^2}{9} + \frac{(y - 1)^2}{4} = 1. c = \sqrt{a^2 - b^2} = \sqrt{5}, \text{ so the foci are } (2 \pm \sqrt{5}, 1).$$



$$19. \frac{y^2}{25} - \frac{x^2}{9} = 1 \Rightarrow a = 5, b = 3, c = \sqrt{25 + 9} = \sqrt{34} \Rightarrow$$


center  $(0, 0)$ , vertices  $(0, \pm 5)$ , foci  $(0, \pm \sqrt{34})$ , asymptotes  $y = \pm \frac{5}{3}x$ .

*Note:* It is helpful to draw a  $2a$ -by- $2b$  rectangle whose center is the center of the hyperbola. The asymptotes are the extended diagonals of the rectangle.



$$20. \frac{x^2}{36} - \frac{y^2}{64} = 1 \Rightarrow a = 6, b = 8, c = \sqrt{36 + 64} = 10 \Rightarrow$$


center  $(0, 0)$ , vertices  $(\pm 6, 0)$ , foci  $(\pm 10, 0)$ , asymptotes  $y = \pm \frac{8}{6}x = \pm \frac{4}{3}x$



$$21. x^2 - y^2 = 100 \Leftrightarrow \frac{x^2}{100} - \frac{y^2}{100} = 1 \Rightarrow a = b = 10,$$

$c = \sqrt{100 + 100} = 10\sqrt{2} \Rightarrow$  center  $(0, 0)$ , vertices  $(\pm 10, 0)$ ,


foci  $(\pm 10\sqrt{2}, 0)$ , asymptotes  $y = \pm \frac{10}{10}x = \pm x$



$$22. y^2 - 16x^2 = 16 \Leftrightarrow \frac{y^2}{16} - \frac{x^2}{1} = 1 \Rightarrow a = 4, b = 1,$$

$c = \sqrt{16 + 1} = \sqrt{17} \Rightarrow$  center  $(0, 0)$ , vertices  $(0, \pm 4)$ ,

foci  $(0, \pm \sqrt{17})$ , asymptotes  $y = \pm \frac{4}{1}x = \pm 4x$




$$23. x^2 - y^2 + 2y = 2 \Leftrightarrow x^2 - (y^2 - 2y + 1) = 2 - 1 \Leftrightarrow$$

$$\frac{x^2}{1} - \frac{(y-1)^2}{1} = 1 \Rightarrow a = b = 1, c = \sqrt{1 + 1} = \sqrt{2} \Rightarrow$$

center  $(0, 1)$ , vertices  $(\pm 1, 1)$ , foci  $(\pm \sqrt{2}, 1)$ ,

asymptotes  $y - 1 = \pm \frac{1}{1}x = \pm x$ .




24.  $9y^2 - 4x^2 - 36y - 8x = 4 \Leftrightarrow$

$$9(y^2 - 4y + 4) - 4(x^2 + 2x + 1) = 4 + 36 - 4 \Leftrightarrow$$

$$9(y - 2)^2 - 4(x + 1)^2 = 36 \Leftrightarrow \frac{(y - 2)^2}{4} - \frac{(x + 1)^2}{9} = 1 \Rightarrow$$

$$a = 2, b = 3, c = \sqrt{4 + 9} = \sqrt{13} \Rightarrow \text{center } (-1, 2), \text{ vertices}$$

$$(-1, 2 \pm 2), \text{ foci } (-1, 2 \pm \sqrt{13}), \text{ asymptotes } y - 2 = \pm \frac{2}{3}(x + 1).$$



25.  $4x^2 = y^2 + 4 \Leftrightarrow 4x^2 - y^2 = 4 \Leftrightarrow \frac{x^2}{1} - \frac{y^2}{4} = 1$ . This is an equation of a *hyperbola* with vertices  $(\pm 1, 0)$ .

The foci are at  $(\pm\sqrt{1+4}, 0) = (\pm\sqrt{5}, 0)$ .

26.  $4x^2 = y + 4 \Leftrightarrow x^2 = \frac{1}{4}(y + 4)$ . This is an equation of a *parabola* with  $4p = \frac{1}{4}$ , so  $p = \frac{1}{16}$ . The vertex is  $(0, -4)$  and the focus is  $(0, -4 + \frac{1}{16}) = (0, -\frac{63}{16})$ .

27.  $x^2 = 4y - 2y^2 \Leftrightarrow x^2 + 2y^2 - 4y = 0 \Leftrightarrow x^2 + 2(y^2 - 2y + 1) = 2 \Leftrightarrow x^2 + 2(y - 1)^2 = 2 \Leftrightarrow \frac{x^2}{2} + \frac{(y - 1)^2}{1} = 1$ . This is an equation of an *ellipse* with vertices at  $(\pm\sqrt{2}, 1)$ . The foci are at  $(\pm\sqrt{2-1}, 1) = (\pm 1, 1)$ .

28.  $y^2 - 2 = x^2 - 2x \Leftrightarrow y^2 - x^2 + 2x = 2 \Leftrightarrow y^2 - (x^2 - 2x + 1) = 2 - 1 \Leftrightarrow \frac{y^2}{1} - \frac{(x - 1)^2}{1} = 1$ . This is an equation of a *hyperbola* with vertices  $(1, \pm 1)$ . The foci are at  $(1, \pm\sqrt{1+1}) = (1, \pm\sqrt{2})$ .

29.  $3x^2 - 6x - 2y = 1 \Leftrightarrow 3x^2 - 6x = 2y + 1 \Leftrightarrow 3(x^2 - 2x + 1) = 2y + 1 + 3 \Leftrightarrow 3(x - 1)^2 = 2y + 4 \Leftrightarrow (x - 1)^2 = \frac{2}{3}(y + 2)$ . This is an equation of a *parabola* with  $4p = \frac{2}{3}$ , so  $p = \frac{1}{6}$ . The vertex is  $(1, -2)$  and the focus is  $(1, -2 + \frac{1}{6}) = (1, -\frac{11}{6})$ .

30.  $x^2 - 2x + 2y^2 - 8y + 7 = 0 \Leftrightarrow (x^2 - 2x + 1) + 2(y^2 - 4y + 4) = -7 + 1 + 8 \Leftrightarrow (x - 1)^2 + 2(y - 2)^2 = 2 \Leftrightarrow \frac{(x - 1)^2}{2} + \frac{(y - 2)^2}{1} = 1$ . This is an equation of an *ellipse* with vertices at  $(1 \pm \sqrt{2}, 2)$ . The foci are at  $(1 \pm \sqrt{2-1}, 2) = (1 \pm 1, 2)$ .

31. The parabola with vertex  $(0, 0)$  and focus  $(1, 0)$  opens to the right and has  $p = 1$ , so its equation is  $y^2 = 4px$ , or  $y^2 = 4x$ .

32. The parabola with focus  $(0, 0)$  and directrix  $y = 6$  has vertex  $(0, 3)$  and opens downward, so  $p = -3$  and its equation is  $(x - 0)^2 = 4p(y - 3)$ , or  $x^2 = -12(y - 3)$ .

33. The distance from the focus  $(-4, 0)$  to the directrix  $x = 2$  is  $2 - (-4) = 6$ , so the distance from the focus to the vertex is  $\frac{1}{2}(6) = 3$  and the vertex is  $(-1, 0)$ . Since the focus is to the left of the vertex,  $p = -3$ . An equation is  $y^2 = 4p(x + 1) \Rightarrow y^2 = -12(x + 1)$ .


34. The parabola with vertex  $(2, 3)$  and focus  $(2, -1)$  opens downward and has  $p = -1 - 3 = -4$ , so its equation is  $(x - 2)^2 = 4p(y - 3)$ , or  $(x - 2)^2 = -16(y - 3)$ .




4. The vector from  $(621, -147, 206)$  to  $(563, 31, 242)$ ,  $\mathbf{v}_1 = \langle -58, 178, 36 \rangle$ , lies in the plane of the rectangle, as does the vector from  $(621, -147, 206)$  to  $(657, -111, 86)$ ,  $\mathbf{v}_2 = \langle 36, 36, -120 \rangle$ . A normal vector for the plane is  $\mathbf{v}_1 \times \mathbf{v}_2 = \langle -1888, -142, -708 \rangle$  or  $\langle 8, 2, 3 \rangle$ , and an equation of the plane is  $8x + 2y + 3z = 5292$ . The line  $L$  intersects this plane when  $8(230 + 630t) + 2(-285 + 390t) + 3(102 + 162t) = 5292 \Rightarrow t = \frac{1858}{3153} \approx 0.589$ . The corresponding point is approximately  $(601.25, -55.18, 197.46)$ . Starting at this point, a portion of the line is hidden behind the rectangle. The line becomes visible again at the left edge of the rectangle, specifically the edge between the points  $(621, -147, 206)$  and  $(657, -111, 86)$ . (This is most easily determined by graphing the rectangle and the line.) A plane through these two points and the camera's location,  $(1000, 0, 0)$ , will clip the line at the point it becomes visible. Two vectors in this plane are  $\mathbf{v}_1 = \langle -379, -147, 206 \rangle$  and  $\mathbf{v}_2 = \langle -343, -111, 86 \rangle$ . A normal vector for the plane is  $\mathbf{v}_1 \times \mathbf{v}_2 = \langle 10224, -38064, -8352 \rangle$  and an equation of the plane is  $213x - 793y - 174z = 213,000$ .  $L$  intersects this plane when  $213(230 + 630t) - 793(-285 + 390t) - 174(102 + 162t) = 213,000 \Rightarrow t = \frac{44,247}{203,268} \approx 0.2177$ . The corresponding point is approximately  $(367.14, -200.11, 137.26)$ . Thus the portion of  $L$  that should be removed is the segment between the points  $(601.25, -55.18, 197.46)$  and  $(367.14, -200.11, 137.26)$ .

## 12.6 Cylinders and Quadric Surfaces


1. (a) In  $\mathbb{R}^2$ , the equation  $y = x^2$  represents a parabola.




- (b) In  $\mathbb{R}^3$ , the equation  $y = x^2$  doesn't involve  $z$ , so any horizontal plane with equation  $z = k$  intersects the graph in a curve with equation  $y = x^2$ . Thus, the surface is a parabolic cylinder, made up of infinitely many shifted copies of the same parabola. The rulings are parallel to the  $z$ -axis.




- (c) In  $\mathbb{R}^3$ , the equation  $z = y^2$  also represents a parabolic cylinder. Since  $x$  doesn't appear, the graph is formed by moving the parabola  $z = y^2$  in the direction of the  $x$ -axis. Thus, the rulings of the cylinder are parallel to the  $x$ -axis.




2. (a)




(b) Since the equation  $y = e^x$  doesn't involve  $z$ , horizontal traces are copies of the curve  $y = e^x$ . The rulings are parallel to the  $z$ -axis.




(c) The equation  $z = e^y$  doesn't involve  $x$ , so vertical traces in  $x = k$  (parallel to the  $yz$ -plane) are copies of the curve  $z = e^y$ . The rulings are parallel to the  $x$ -axis.




3. Since  $y$  is missing from the equation, the vertical traces  $x^2 + z^2 = 1$ ,  $y = k$ , are copies of the same circle in the plane  $y = k$ . Thus the surface  $x^2 + z^2 = 1$  is a circular cylinder with rulings parallel to the  $y$ -axis.




4. Since  $z$  is missing from the equation, the horizontal traces  $4x^2 + y^2 = 4$ ,  $z = k$ , are copies of the same ellipse in the plane  $z = k$ . Thus the surface  $4x^2 + y^2 = 4$  is an elliptic cylinder with rulings parallel to the  $z$ -axis.




5. Since  $x$  is missing, each vertical trace  $z = 1 - y^2$ ,  $x = k$ , is a copy of the same parabola in the plane  $x = k$ . Thus the surface  $z = 1 - y^2$  is a parabolic cylinder with rulings parallel to the  $x$ -axis.




6. Since  $x$  is missing, each vertical trace  $y = z^2$ ,  $x = k$ , is a copy of the same parabola in the plane  $x = k$ . Thus the surface  $y = z^2$  is a parabolic cylinder with rulings parallel to the  $x$ -axis.




2. (a)




(b) Since the equation  $y = e^x$  doesn't involve  $z$ , horizontal traces are copies of the curve  $y = e^x$ . The rulings are parallel to the  $z$ -axis.




(c) The equation  $z = e^y$  doesn't involve  $x$ , so vertical traces in  $x = k$  (parallel to the  $yz$ -plane) are copies of the curve  $z = e^y$ . The rulings are parallel to the  $x$ -axis.




3. Since  $y$  is missing from the equation, the vertical traces  $x^2 + z^2 = 1$ ,  $y = k$ , are copies of the same circle in the plane  $y = k$ . Thus the surface  $x^2 + z^2 = 1$  is a circular cylinder with rulings parallel to the  $y$ -axis.




4. Since  $z$  is missing from the equation, the horizontal traces  $4x^2 + y^2 = 4$ ,  $z = k$ , are copies of the same ellipse in the plane  $z = k$ . Thus the surface  $4x^2 + y^2 = 4$  is an elliptic cylinder with rulings parallel to the  $z$ -axis.



5. Since  $x$  is missing, each vertical trace  $z = 1 - y^2$ ,  $x = k$ , is a copy of the same parabola in the plane  $x = k$ . Thus the surface  $z = 1 - y^2$  is a parabolic cylinder with rulings parallel to the  $x$ -axis.




6. Since  $x$  is missing, each vertical trace  $y = z^2$ ,  $x = k$ , is a copy of the same parabola in the plane  $x = k$ . Thus the surface  $y = z^2$  is a parabolic cylinder with rulings parallel to the  $x$ -axis.






10. (a) The traces of  $-x^2 - y^2 + z^2 = 1$  in  $x = k$  are  $-y^2 + z^2 = 1 + k^2$ , a family of hyperbolas, as are the traces in  $y = k$ ,  $-x^2 + z^2 = 1 + k^2$ . The traces in  $z = k$  are  $x^2 + y^2 = k^2 - 1$ , a family of circles for  $|k| > 1$ . As  $|k|$  increases, the radii of the circles increase; the traces are empty for  $|k| < 1$ . This behavior, combined with the vertical traces, gives the graph of the hyperboloid of two sheets in Table 1.


- (b) The graph has the same shape as the hyperboloid in part (a) but is rotated so that its axis is the  $x$ -axis. Traces in  $x = k$ ,  $|k| > 1$ , are circles, while traces in  $y = k$  and  $z = k$  are hyperbolas.




11. For  $x = y^2 + 4z^2$ , the traces in  $x = k$  are  $y^2 + 4z^2 = k$ . When  $k > 0$  we have a family of ellipses. When  $k = 0$  we have just a point at the origin, and the trace is empty for  $k < 0$ . The traces in  $y = k$  are  $x = 4z^2 + k^2$ , a family of parabolas opening in the positive  $x$ -direction. Similarly, the traces in  $z = k$  are  $x = y^2 + 4k^2$ , a family of parabolas opening in the positive  $x$ -direction. We recognize the graph as an elliptic paraboloid with axis the  $x$ -axis and vertex the origin.




12.  $4x^2 + 9y^2 + 9z^2 = 36$ . The traces in  $x = k$  are  $9y^2 + 9z^2 = 36 - 4k^2 \Leftrightarrow y^2 + z^2 = 4 - \frac{4}{9}k^2$ , a family of circles for  $|k| < 3$ . (The traces are a single point for  $|k| = 3$  and are empty for  $|k| > 3$ .) The traces in  $y = k$  are  $4x^2 + 9z^2 = 36 - 9k^2$ , a family of ellipses for  $|k| < 2$ . Similarly, the traces in  $z = k$  are the ellipses  $4x^2 + 9y^2 = 36 - 9k^2$ ,  $|k| < 2$ . The graph is an ellipsoid centered at the origin with intercepts  $x = \pm 3$ ,  $y = \pm 2$ ,  $z = \pm 2$ .




13.  $x^2 = 4y^2 + z^2$ . The traces in  $x = k$  are the ellipses  $4y^2 + z^2 = k^2$ . The traces in  $y = k$  are  $x^2 - z^2 = 4k^2$ , hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . Similarly, the traces in  $z = k$  are  $x^2 - 4y^2 = k^2$ , hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . We recognize the graph as an elliptic cone with axis the  $x$ -axis and vertex the origin.




14.  $z^2 - 4x^2 - y^2 = 4$ . The traces in  $x = k$  are the hyperbolas  $z^2 - y^2 = 4 + 4k^2$ , and the traces in  $y = k$  are the hyperbolas  $z^2 - 4x^2 = 4 + k^2$ . The traces in  $z = k$  are  $4x^2 + y^2 = k^2 - 4$ , a family of ellipses for  $|k| > 2$ . (The traces are a single point for  $|k| = 2$  and are empty for  $|k| < 2$ .) The surface is a hyperboloid of two sheets with axis the  $z$ -axis.




15.  $9y^2 + 4z^2 = x^2 + 36$ . The traces in  $x = k$  are  $9y^2 + 4z^2 = k^2 + 36$ , a family of ellipses. The traces in  $y = k$  are  $4z^2 - x^2 = 9(4 - k^2)$ , a family of hyperbolas for  $|k| \neq 2$  and two intersecting lines when  $|k| = 2$ . (Note that the hyperbolas are oriented differently for  $|k| < 2$  than for  $|k| > 2$ .) The traces in  $z = k$  are  $9y^2 - x^2 = 4(9 - k^2)$ , a family of hyperbolas when  $|k| \neq 3$  (oriented differently for  $|k| < 3$  than for  $|k| > 3$ ) and two intersecting lines when  $|k| = 3$ . We recognize the graph as a hyperboloid of one sheet with axis the  $x$ -axis.




16.  $3x^2 + y + 3z^2 = 0$ . The traces in  $x = k$  are the parabolas  $y = -3z^2 - 3k^2$  which open to the left (in the negative  $y$ -direction). Traces in  $y = k$  are  $3x^2 + 3z^2 = -k \Leftrightarrow x^2 + z^2 = -\frac{k}{3}$ , a family of circles for  $k < 0$ . (Traces are empty for  $k > 0$  and a single point for  $k = 0$ .) Traces in  $z = k$  are the parabolas  $y = -3x^2 - 3k^2$  which open in the negative  $y$ -direction. The graph is a circular paraboloid with axis the  $y$ -axis, opening in the negative  $y$ -direction, and vertex the origin.




17.  $\frac{x^2}{9} + \frac{y^2}{25} + \frac{z^2}{4} = 1$ . The traces in  $x = k$  are  $\frac{y^2}{25} + \frac{z^2}{4} = 1 - \frac{k^2}{9}$ , a family of ellipses for  $|k| < 3$ . (The traces are a single point for  $|k| = 3$  and are empty for  $|k| > 3$ .) The traces in  $y = k$  are the ellipses  $\frac{x^2}{9} + \frac{z^2}{4} = 1 - \frac{k^2}{25}$ ,  $|k| < 5$ , and the traces in  $z = k$  are the ellipses  $\frac{x^2}{9} + \frac{y^2}{25} = 1 - \frac{k^2}{4}$ ,  $|k| < 2$ . The surface is an ellipsoid centered at the origin with intercepts  $x = \pm 3$ ,  $y = \pm 5$ ,  $z = \pm 2$ .




14.  $z^2 - 4x^2 - y^2 = 4$ . The traces in  $x = k$  are the hyperbolas  $z^2 - y^2 = 4 + 4k^2$ , and the traces in  $y = k$  are the hyperbolas  $z^2 - 4x^2 = 4 + k^2$ . The traces in  $z = k$  are  $4x^2 + y^2 = k^2 - 4$ , a family of ellipses for  $|k| > 2$ . (The traces are a single point for  $|k| = 2$  and are empty for  $|k| < 2$ .) The surface is a hyperboloid of two sheets with axis the  $z$ -axis.




15.  $9y^2 + 4z^2 = x^2 + 36$ . The traces in  $x = k$  are  $9y^2 + 4z^2 = k^2 + 36$ , a family of ellipses. The traces in  $y = k$  are  $4z^2 - x^2 = 9(4 - k^2)$ , a family of hyperbolas for  $|k| \neq 2$  and two intersecting lines when  $|k| = 2$ . (Note that the hyperbolas are oriented differently for  $|k| < 2$  than for  $|k| > 2$ .) The traces in  $z = k$  are  $9y^2 - x^2 = 4(9 - k^2)$ , a family of hyperbolas when  $|k| \neq 3$  (oriented differently for  $|k| < 3$  than for  $|k| > 3$ ) and two intersecting lines when  $|k| = 3$ . We recognize the graph as a hyperboloid of one sheet with axis the  $x$ -axis.



16.  $3x^2 + y + 3z^2 = 0$ . The traces in  $x = k$  are the parabolas  $y = -3z^2 - 3k^2$  which open to the left (in the negative  $y$ -direction). Traces in  $y = k$  are  $3x^2 + 3z^2 = -k \iff x^2 + z^2 = -\frac{k}{3}$ , a family of circles for  $k < 0$ . (Traces are empty for  $k > 0$  and a single point for  $k = 0$ .) Traces in  $z = k$  are the parabolas  $y = -3x^2 - 3k^2$  which open in the negative  $y$ -direction. The graph is a circular paraboloid with axis the  $y$ -axis, opening in the negative  $y$ -direction, and vertex the origin.




17.  $\frac{x^2}{9} + \frac{y^2}{25} + \frac{z^2}{4} = 1$ . The traces in  $x = k$  are  $\frac{y^2}{25} + \frac{z^2}{4} = 1 - \frac{k^2}{9}$ , a family of ellipses for  $|k| < 3$ . (The traces are a single point for  $|k| = 3$  and are empty for  $|k| > 3$ .) The traces in  $y = k$  are the ellipses  $\frac{x^2}{9} + \frac{z^2}{4} = 1 - \frac{k^2}{25}$ ,  $|k| < 5$ , and the traces in  $z = k$  are the ellipses  $\frac{x^2}{9} + \frac{y^2}{25} = 1 - \frac{k^2}{4}$ ,  $|k| < 2$ . The surface is an ellipsoid centered at the origin with intercepts  $x = \pm 3$ ,  $y = \pm 5$ ,  $z = \pm 2$ .







18.  $3x^2 - y^2 + 3z^2 = 0$ . The traces in  $x = k$  are  $y^2 - 3z^2 = 3k^2$ , a family of hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . Traces in  $y = k$  are the circles  $3x^2 + 3z^2 = k^2 \Leftrightarrow x^2 + z^2 = \frac{1}{3}k^2$ . The traces in  $z = k$  are  $y^2 - 3x^2 = 3k^2$ , hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . We recognize the surface as a circular cone with axis the  $y$ -axis and vertex the origin.



19.  $y = z^2 - x^2$ . The traces in  $x = k$  are the parabolas  $y = z^2 - k^2$ , opening in the positive  $y$ -direction. The traces in  $y = k$  are  $k = z^2 - x^2$ , two intersecting lines when  $k = 0$  and a family of hyperbolas for  $k \neq 0$  (note that the hyperbolas are oriented differently for  $k > 0$  than for  $k < 0$ ). The traces in  $z = k$  are the parabolas  $y = k^2 - x^2$  which open in the negative  $y$ -direction. Thus the surface is a hyperbolic paraboloid centered at  $(0, 0, 0)$ .



20.  $x = y^2 - z^2$ . The traces in  $x = k$  are  $y^2 - z^2 = k$ , two intersecting lines when  $k = 0$  and a family of hyperbolas for  $k \neq 0$  (oriented differently for  $k > 0$  than for  $k < 0$ ). The traces in  $y = k$  are the parabolas  $x = -z^2 + k^2$ , opening in the negative  $x$ -direction, and the traces in  $z = k$  are the parabolas  $x = y^2 - k^2$  which open in the positive  $x$ -direction. The graph is a hyperbolic paraboloid centered at  $(0, 0, 0)$ .



21. This is the equation of an ellipsoid:  $x^2 + 4y^2 + 9z^2 = x^2 + \frac{y^2}{(1/2)^2} + \frac{z^2}{(1/3)^2} = 1$ , with  $x$ -intercepts  $\pm 1$ ,  $y$ -intercepts  $\pm \frac{1}{2}$  and  $z$ -intercepts  $\pm \frac{1}{3}$ . So the major axis is the  $x$ -axis and the only possible graph is VII.

22. This is the equation of an ellipsoid:  $9x^2 + 4y^2 + z^2 = \frac{x^2}{(1/3)^2} + \frac{y^2}{(1/2)^2} + z^2 = 1$ , with  $x$ -intercepts  $\pm \frac{1}{3}$ ,  $y$ -intercepts  $\pm \frac{1}{2}$  and  $z$ -intercepts  $\pm 1$ . So the major axis is the  $z$ -axis and the only possible graph is IV.


23. This is the equation of a hyperboloid of one sheet, with  $a = b = c = 1$ . Since the coefficient of  $y^2$  is negative, the axis of the hyperboloid is the  $y$ -axis, hence the correct graph is II.

24. This is a hyperboloid of two sheets, with  $a = b = c = 1$ . This surface does not intersect the  $xz$ -plane at all, so the axis of the hyperboloid is the  $y$ -axis and the graph is III.


25. There are no real values of  $x$  and  $z$  that satisfy this equation for  $y < 0$ , so this surface does not extend to the left of the  $xz$ -plane. The surface intersects the plane  $y = k > 0$  in an ellipse. Notice that  $y$  occurs to the first power whereas  $x$  and  $z$  occur to the second power. So the surface is an elliptic paraboloid with axis the  $y$ -axis. Its graph is VI.

26. This is the equation of a cone with axis the  $y$ -axis, so the graph is I.


18.  $3x^2 - y^2 + 3z^2 = 0$ . The traces in  $x = k$  are  $y^2 - 3z^2 = 3k^2$ , a family of hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . Traces in  $y = k$  are the circles  $3x^2 + 3z^2 = k^2 \Leftrightarrow x^2 + z^2 = \frac{1}{3}k^2$ . The traces in  $z = k$  are  $y^2 - 3x^2 = 3k^2$ , hyperbolas for  $k \neq 0$  and two intersecting lines if  $k = 0$ . We recognize the surface as a circular cone with axis the  $y$ -axis and vertex the origin.



19.  $y = z^2 - x^2$ . The traces in  $x = k$  are the parabolas  $y = z^2 - k^2$ , opening in the positive  $y$ -direction. The traces in  $y = k$  are  $k = z^2 - x^2$ , two intersecting lines when  $k = 0$  and a family of hyperbolas for  $k \neq 0$  (note that the hyperbolas are oriented differently for  $k > 0$  than for  $k < 0$ ). The traces in  $z = k$  are the parabolas  $y = k^2 - x^2$  which open in the negative  $y$ -direction. Thus the surface is a hyperbolic paraboloid centered at  $(0, 0, 0)$ .



20.  $x = y^2 - z^2$ . The traces in  $x = k$  are  $y^2 - z^2 = k$ , two intersecting lines when  $k = 0$  and a family of hyperbolas for  $k \neq 0$  (oriented differently for  $k > 0$  than for  $k < 0$ ). The traces in  $y = k$  are the parabolas  $x = -z^2 + k^2$ , opening in the negative  $x$ -direction, and the traces in  $z = k$  are the parabolas  $x = y^2 - k^2$  which open in the positive  $x$ -direction. The graph is a hyperbolic paraboloid centered at  $(0, 0, 0)$ .



21. This is the equation of an ellipsoid:  $x^2 + 4y^2 + 9z^2 = x^2 + \frac{y^2}{(1/2)^2} + \frac{z^2}{(1/3)^2} = 1$ , with  $x$ -intercepts  $\pm 1$ ,  $y$ -intercepts  $\pm \frac{1}{2}$  and  $z$ -intercepts  $\pm \frac{1}{3}$ . So the major axis is the  $x$ -axis and the only possible graph is VII.

22. This is the equation of an ellipsoid:  $9x^2 + 4y^2 + z^2 = \frac{x^2}{(1/3)^2} + \frac{y^2}{(1/2)^2} + z^2 = 1$ , with  $x$ -intercepts  $\pm \frac{1}{3}$ ,  $y$ -intercepts  $\pm \frac{1}{2}$  and  $z$ -intercepts  $\pm 1$ . So the major axis is the  $z$ -axis and the only possible graph is IV.

23. This is the equation of a hyperboloid of one sheet, with  $a = b = c = 1$ . Since the coefficient of  $y^2$  is negative, the axis of the hyperboloid is the  $y$ -axis, hence the correct graph is II.


24. This is a hyperboloid of two sheets, with  $a = b = c = 1$ . This surface does not intersect the  $xz$ -plane at all, so the axis of the hyperboloid is the  $y$ -axis and the graph is III.

25. There are no real values of  $x$  and  $z$  that satisfy this equation for  $y < 0$ , so this surface does not extend to the left of the  $xz$ -plane. The surface intersects the plane  $y = k > 0$  in an ellipse. Notice that  $y$  occurs to the first power whereas  $x$  and  $z$  occur to the second power. So the surface is an elliptic paraboloid with axis the  $y$ -axis. Its graph is VI.


26. This is the equation of a cone with axis the  $y$ -axis, so the graph is I.




27. This surface is a cylinder because the variable  $y$  is missing from the equation. The intersection of the surface and the  $xz$ -plane is an ellipse. So the graph is VIII.
28. This is the equation of a hyperbolic paraboloid. The trace in the  $xy$ -plane is the parabola  $y = x^2$ . So the correct graph is V.
29. Vertical traces parallel to the  $xz$ -plane are circles centered at the origin whose radii increase as  $y$  decreases. (The trace in  $y = 1$  is just a single point and the graph suggests that traces in  $y = k$  are empty for  $k > 1$ .) The traces in vertical planes parallel to the  $yz$ -plane are parabolas opening to the left that shift to the left as  $|x|$  increases. One surface that fits this description is a circular paraboloid, opening to the left, with vertex  $(0, 1, 0)$ .




30. The vertical traces parallel to the  $yz$ -plane are ellipses that are smallest in the  $yz$ -plane and increase in size as  $|x|$  increases. One surface that fits this description is a hyperboloid of one sheet with axis the  $x$ -axis. The horizontal traces in  $z = k$  (hyperbolas and intersecting lines) also fit this surface, as shown in the figure.



31.  $y^2 = x^2 + \frac{1}{9}z^2$  or  $y^2 = x^2 + \frac{z^2}{9}$  represents an elliptic cone with vertex  $(0, 0, 0)$  and axis the  $y$ -axis.




32.  $4x^2 - y + 2z^2 = 0$  or  $y = \frac{x^2}{1/4} + \frac{z^2}{1/2}$  or  $\frac{y}{4} = x^2 + \frac{z^2}{2}$  represents an elliptic paraboloid with vertex  $(0, 0, 0)$  and axis the  $y$ -axis.






33.  $x^2 + 2y - 2z^2 = 0$  or  $2y = 2z^2 - x^2$  or  $y = z^2 - \frac{x^2}{2}$   
represents a hyperbolic paraboloid with center  $(0, 0, 0)$ .




34.  $y^2 = x^2 + 4z^2 + 4$  or  $-x^2 + y^2 - 4z^2 = 4$  or  $-\frac{x^2}{4} + \frac{y^2}{4} - z^2 = 1$  represents a hyperboloid of two sheets with axis the  $y$ -axis.




35. Completing squares in  $x$  and  $y$  gives

$$(x^2 - 2x + 1) + (y^2 - 6y + 9) - z = 0 \Leftrightarrow$$

$(x - 1)^2 + (y - 3)^2 - z = 0$  or  $z = (x - 1)^2 + (y - 3)^2$ , a circular paraboloid opening upward with vertex  $(1, 3, 0)$  and axis the vertical line  $x = 1, y = 3$ .



36. Completing squares in  $x$  and  $z$  gives  $(x^2 - 4x + 4) - y^2 - (z^2 + 2z + 1) + 3 = 0 + 4 - 1 \Leftrightarrow$   
 $(x - 2)^2 - y^2 - (z + 1)^2 = 0$  or  $(x - 2)^2 = y^2 + (z + 1)^2$ , a circular cone with vertex  $(2, 0, -1)$  and axis the horizontal line  $y = 0, z = -1$ .




37. Completing squares in  $x$  and  $z$  gives



$$(x^2 - 4x + 4) - y^2 + (z^2 - 2z + 1) = 0 + 4 + 1 \Leftrightarrow$$

$$(x - 2)^2 - y^2 + (z - 1)^2 = 5 \text{ or } \frac{(x - 2)^2}{5} - \frac{y^2}{5} + \frac{(z - 1)^2}{5} = 1, \text{ a}$$


hyperboloid of one sheet with center  $(2, 0, 1)$  and axis the horizontal line  $x = 2, z = 1$ .




42. We plot the surface  $z = x^2 - 6x + 4y^2$ .




43.




44.



45. The curve  $y = \sqrt{x}$  is equivalent to  $x = y^2$ ,  $y \geq 0$ . Rotating the curve about the  $x$ -axis creates a circular paraboloid with vertex at the origin, axis the  $x$ -axis, opening in the positive  $x$ -direction. The trace in the  $xy$ -plane is  $x = y^2$ ,  $z = 0$ , and the trace in the  $xz$ -plane is a parabola of the same shape:  $x = z^2$ ,  $y = 0$ . An equation for the surface is  $x = y^2 + z^2$ .



46. Rotating the line  $z = 2y$  about the  $z$ -axis creates a (right) circular cone with vertex at the origin and axis the  $z$ -axis. Traces in  $z = k$  ( $k \neq 0$ ) are circles with center  $(0, 0, k)$  and radius  $y = z/2 = k/2$ , so an equation for the trace is  $x^2 + y^2 = (k/2)^2$ ,  $z = k$ . Thus an equation for the surface is  $x^2 + y^2 = (z/2)^2$  or  $4x^2 + 4y^2 = z^2$ .



47. Let  $P = (x, y, z)$  be an arbitrary point equidistant from  $(-1, 0, 0)$  and the plane  $x = 1$ . Then the distance from  $P$  to  $(-1, 0, 0)$  is  $\sqrt{(x+1)^2 + y^2 + z^2}$  and the distance from  $P$  to the plane  $x = 1$  is  $|x - 1|/\sqrt{1^2} = |x - 1|$  (by Equation 12.5.9). So  $|x - 1| = \sqrt{(x+1)^2 + y^2 + z^2} \Leftrightarrow (x - 1)^2 = (x + 1)^2 + y^2 + z^2 \Leftrightarrow x^2 - 2x + 1 = x^2 + 2x + 1 + y^2 + z^2 \Leftrightarrow -4x = y^2 + z^2$ . Thus the collection of all such points  $P$  is a circular paraboloid with vertex at the origin, axis the  $x$ -axis, which opens in the negative  $x$ -direction.