



An Inverse Function Theorem for Free Groups

Joan S. Birman

Proceedings of the American Mathematical Society, Vol. 41, No. 2 (Dec., 1973),
634-638.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28197312%2941%3A2%3C634%3AAIFTF%3E2.0.CO%3B2-S>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

AN INVERSE FUNCTION THEOREM FOR FREE GROUPS

JOAN S. BIRMAN¹

ABSTRACT. Let F_n be a free group of rank n with free basis x_1, \dots, x_n . Let $\{y_1, \dots, y_k\}$ be a set of $k \leq n$ elements of F_n , where each y_i is represented by a word $Y_i(x_1, \dots, x_n)$ in the generators x_j . Let $\partial y_i / \partial x_j$ denote the free derivative of y_i with respect to x_j , and let $J_{kn} = \|\partial y_i / \partial x_j\|$ denote the $k \times n$ Jacobian matrix. **THEOREM.** *If $k=n$, the set $\{y_1, \dots, y_n\}$ generates F_n if and only if J_{nn} has a right inverse. If $k < n$, the set $\{y_1, \dots, y_k\}$ may be extended to a set of elements which generate F_n only if J_{kn} has a right inverse. Several applications are given.*

Let Z denote the ring of rational integers, and let ZF_n denote the integral group ring of a free group F_n which has the free basis x_1, \dots, x_n . Let $\partial / \partial x_j : ZF_n \rightarrow ZF_n$ denote the j th free partial derivative, in the sense of R. H. Fox [3]. The mapping $\partial / \partial x_j$ is defined as follows: If $w \in F_n$ is represented by the word $x_{\mu_1}^{\varepsilon_1} \cdots x_{\mu_r}^{\varepsilon_r}$, where $\varepsilon_i = \pm 1$ and $\mu_i = 1, \dots, n$, then $\partial w / \partial x_j \in ZF_n$ is defined by:

$$(1) \quad \frac{\partial w}{\partial x_j} = \sum_{k=1}^n \delta_{j, \mu_k} \varepsilon_k x_{\mu_1}^{\varepsilon_1} \cdots x_{\mu_{k-1}}^{\varepsilon_{k-1}} x_{\mu_{k+1}}^{\varepsilon_{k+1}} \cdots x_{\mu_r}^{\varepsilon_r}$$

where δ means the Kronecker symbol. More generally, if $u = \sum_{i=1}^t c_i w_i$, $c_i \in Z$, $w_i \in F_n$, we define

$$(2) \quad \frac{\partial u}{\partial x_j} = \sum_{i=1}^t c_i \frac{\partial w_i}{\partial x_j}$$

It is easy to show that this definition is independent of the choice representatives of the elements $w_i \in F_n$.

There are known analogues between the "free" calculus of polynomials in the noncommuting indeterminates x_1, \dots, x_n and the "ordinary" calculus of polynomials in commuting indeterminates, such as the existence of Taylor series [3]. However, it is worth noting that if α is the

Received by the editors November 30, 1972 and, in revised form, February 26, 1973.
 AMS (MOS) subject classifications (1970). Primary 55A05, 20E99; Secondary 20F55.
 Key words and phrases. Free calculus, primitive elements, free basis, invertible matrices.

¹ This work was supported in part by NSF Grant GP-34324X.

abelianizing homomorphism acting on F_n , and if α_* is the induced homomorphism from $ZF_n \rightarrow ZF_n^a$, then the image under α_* of the free partial derivative of u , as defined by (2), is not the ordinary partial derivative of $\alpha_*(u)$. Thus the free calculus and the ordinary calculus appear to be distinct theories. The purpose of this note is to point out a new analogue between the free calculus and the ordinary calculus, an "inverse function theorem" for free groups.

INVERSE FUNCTION THEOREM. *Let $\{y_1, \dots, y_k\}$ be a set of $k \leq n$ elements of F_n . Let J_{kn} denote the $k \times n$ "Jacobian" matrix $\|\partial y_i / \partial x_j\|$.*

(i) *If $k = n$, a necessary and sufficient condition for $\{y_1, \dots, y_n\}$ to be a generating set for F_n is that J_{nn} have a right inverse.*

(ii) *If $k < n$, a necessary condition for $\{y_1, \dots, y_k\}$ to extend to a generating set $\{y_1, \dots, y_n\}$ is that J_{kn} have a right inverse.*

PROOF. We first establish the sufficiency of the condition (i).² Suppose that $B = \|\beta_{ij}\|$ is a right inverse of J_{nn} . By a theorem of M. S. Montgomery [6], the matrix B is also a left inverse of J_{nn} . Hence

$$(3) \quad \sum_{s=1}^n \beta_{is} \left(\frac{\partial y_s}{\partial x_j} \right) = \delta_{ij} \quad (i, j = 1, \dots, n).$$

Multiplying both sides of (3) by $x_j - 1$, and summing over j , we obtain

$$(4) \quad \sum_{s=1}^n \beta_{is} \sum_{j=1}^n \frac{\partial y_s}{\partial x_j} (x_j - 1) = x_i - 1 \quad (i = 1, \dots, n).$$

By the "fundamental theorem" of free calculus [3]:

$$(5) \quad \sum_{j=1}^n \frac{\partial y_s}{\partial x_j} (x_j - 1) = y_s - 1 \quad (s = 1, \dots, n).$$

Hence

$$(6) \quad \sum_{s=1}^n \beta_{is} (y_s - 1) = x_i - 1 \quad (i = 1, \dots, n).$$

Now let H be the subgroup of F_n generated by y_1, \dots, y_n and let I_H be the ideal of ZF_n generated by $y_1 - 1, \dots, y_n - 1$. According to equation (6), the ring elements $x_i - 1$ belong to I_H for each $i = 1, \dots, n$. But then, by Lemma 4.1 of [2], it follows that $x_i \in H$ for each $i = 1, \dots, n$. Hence H coincides with F_n , and our result is established.

Necessity may be established by noting that if $\{y_1, \dots, y_k\}$ extends to a basis $\{y_1, \dots, y_n\}$, then we may write each x_i as a word $X_i(y_1, \dots, y_n)$,

² The very brief proof of sufficiency given here was suggested by the referee. It replaces a longer and more computational proof in an earlier version of this paper. The author wishes to thank the referee for his constructive suggestions.

in the generators y_j . Moreover, if y_i is represented by the word $Y_i(x_1, \dots, x_n)$, then we will have

$$(7) \quad Y_i(X_1(y_1, \dots, y_n), \dots, X_n(y_1, \dots, y_n)) = y_i \quad (i = 1, \dots, n).$$

The chain rule (see [3]) applied to (7) gives

$$(8) \quad \sum_{s=1}^n \left(\frac{\partial Y_i(x_1, \dots, x_n)}{\partial x_s} \right) \left(\frac{\partial X_s(y_1, \dots, y_n)}{\partial y_j} \right) = \delta_{ij} \quad (i, j = 1, \dots, n).$$

Hence the $n \times n$ matrix J_{nn} has a right inverse, $J_{nn}^* = \|\partial x_i / \partial y_j\|$. Thus condition (i) is seen to be sufficient. Also the $k \times n$ submatrix J_{kn} formed by the first k rows of J_{nn} has a right inverse, namely the submatrix J_{nk}^* of J_{nn}^* formed by the first k columns of J_{nn}^* . This proves (ii).

As an application, we will use the inverse function theorem to give a new proof of a classical theorem of J. Nielsen. This application was suggested by the referee. The proof below is a modification of his proof.

COROLLARY 1 (J. Nielsen, see [5]). *Any set of n elements which generate a free group of rank n are a set of free generators.*

PROOF. Suppose that $\{y_1, \dots, y_n\}$ generate F_n , and suppose also that $\{y_1, \dots, y_n\}$ satisfy the relation

$$(9) \quad r(y_1, \dots, y_n) = 1,$$

where we assume that $r(y_1, \dots, y_n)$ is freely reduced as a word in the y_j 's. By the chain rule it follows that

$$(10) \quad \sum_{\partial=1}^n \frac{\partial r}{\partial y_{\partial}} \frac{\partial y_{\partial}}{\partial x_i} = 0, \quad i = 1, \dots, n.$$

Setting $r = (\partial r / \partial y_1, \dots, \partial r / \partial y_n)$, we may rewrite (10) in the form

$$(11) \quad rJ_{nn} = o.$$

By the inverse function theorem, the matrix J_{nn} has a right inverse, say B . Multiplying both sides of (11) by B we obtain

$$(12) \quad rJ_{nn}B = rI = r = o,$$

hence $\partial r / \partial y_j = 0$ for each $j = 1, \dots, n$. But then it follows that r does not involve the letter y_j , since no cancellations are possible in (1) if a word is freely reduced. It then follows that r must be the trivial relator. This completes the proof of Corollary 1.

We observe that our theorem may be applied both ways. Consider first the case $k = n$. An algorithm for deciding whether a set of n elements

in a free group F_n a basis was discovered by J. Nielsen (see Chapter 2 of [5]), and Nielsen's algorithm translates into a straightforward algorithm for expressing a Jacobian matrix as a product of elementary invertible matrices over ZF_n .³ On the other hand, it is known [7] that the ring ZF_n can be embedded in a skew field K . Since a procedure exists for finding inverses of invertible matrices over skew fields (see Chapter IV of [1]), and since such inverses are unique, one may decide whether a Jacobian matrix is invertible over ZF_n by computing its inverse over K , and seeing whether in fact the entries are in ZF_n . This yields a new test to decide if y_1, \dots, y_n are a basis. This procedure is not however, a practical alternative to Nielsen's relatively simple algorithm; the elementary invertible matrices which are obtained by the method in [1] are almost certainly not in ZF_n , even when their product is in ZF_n , so that this procedure is unnecessarily complex. Our theorem does, however, yield a very simple necessary condition which a set $\{y_1, \dots, y_n\}$ must survive if it is a basis:

COROLLARY 2. *Let J_{nn}^α denote the image of J_{nn} under the abelianizing homomorphism α_* acting on ZF_n . Then $\{y_1, \dots, y_n\}$ is a basis for F_n only if $\det J_{nn}^\alpha$ is a unit in ZF_n^α .*

PROOF. By Theorem 1, p. 59, of [4], a square matrix over a commutative ring with 1 is invertible if and only if its determinant is a unit. \square

To see that the condition in Corollary 2 is not sufficient, let $n=2$ and consider the elements

$$(13) \quad y_1 = x_1,$$

$$(14) \quad y_2 = x_2 x_1 x_2 x_1^{-1} x_2^{-1} x_1^2 x_2^2 x_1^{-2} x_2^{-1} x_1 x_2^{-1} x_1^{-1} x_2^2 x_1^2 x_2^{-2} x_1^{-2}.$$

A simple calculation shows that $\det \|\partial y_i / \partial x_j\|^\alpha = 1$, yet y_1 and y_2 are not primitive, by the test given in Corollary N4, p. 169, of [5].

The more difficult question of deciding whether a set of $k < n$ elements in a free group are primitive was solved by J. H. C. Whitehead [9], [10] and by E. Rapaport [8], and the inverse function theorem may be applied to yield an analogous algorithm for deciding when a $k \times n$ Jacobian matrix over ZF_n has a right inverse. Once again, the Jacobian matrices corresponding to Whitehead transformations are a very pleasant set of elementary invertible matrices.

³ Note that the mapping which we have defined from $\text{Aut } F_n$ to the ring of invertible matrices over ZF_n is a crossed homomorphism. That is, if α and β are automorphisms of F_n which have Jacobian matrices $\|a_{i,j}\|$ and $\|b_{i,j}\|$ respectively, then the Jacobian matrix corresponding to $\alpha\beta$ is the product $\|a_{i,j}\| \|b_{i,j}\|_\alpha$, where $\|b_{i,j}\|_\alpha$ denotes the Jacobian matrix of β with respect to the transformed basis $\alpha(x_1), \dots, \alpha(x_n)$. Thus, in order to apply Nielsen's algorithm, one must repeatedly change basis.

REFERENCES

1. E. Artin, *Geometric algebra*, Interscience Tracts in Pure and Appl. Math., no. 3, Interscience, New York, 1957. MR 18, 553.
2. D. Cohen, *Groups of cohomological dimension one*, Lecture Notes in Math., vol. 245, Springer-Verlag, Berlin and New York, 1972.
3. R. H. Fox, *Free differential calculus. I. Derivation in the free group ring*, Ann. of Math. (2) 57 (1953), 547–560. MR 14, 843.
4. N. Jacobson, *Lectures in abstract algebra. I. Basic concepts*, Van Nostrand, Princeton, N.J., 1951. MR 12, 794.
5. W. Magnus, A. Karrass and D. Solitar, *Combinatorial group theory. Presentations of groups in terms of generators and relations*, Pure and Appl. Math. vol. 13, Interscience, New York, 1966. MR 34 #7617.
6. M. S. Montgomery, *Left and right inverses in group algebras*, Bull. Amer. Math. Soc. 75 (1969), 539–540. MR 39 #327.
7. B. Neumann, *On ordered division rings*, Trans. Amer. Math. Soc. 66 (1949), 202–252. MR 11, 311.
8. E. S. Rapaport, *On free groups and their automorphisms*, Acta Math. 99 (1958), 139–163. MR 24 #A1302.
9. J. H. C. Whitehead, *On certain sets of elements in a free group*, Proc. London Math. Soc. 41 (1936), 48–56.
10. ———, *On equivalent sets of elements in a free group*, Ann. of Math. 37 (1936), 782–800.

DEPARTMENT OF MATHEMATICS, STEVENS INSTITUTE OF TECHNOLOGY, HOBOKEN,
NEW JERSEY 07030

Current address: Department of Mathematics, Columbia University, New York,
New York 10027