





Gauge theories and Bethe eigenfunctions

based on *Quasimap counts and Bethe eigenfunctions*, Mina Aganagic & A.O., arXiv:1704.08746

"Bethe Ansatz" is the art and science of finding spectra and eigenfunctions in quantum integrable systems with a quantum group symmetry



"Bethe Ansatz" is the art and science of finding spectra and eigenfunctions in quantum integrable systems with a quantum group symmetry



Quantum groups (quantum loop algebras)

$\hat{\mathfrak{g}}$

If \mathfrak{g} is a Lie algebra, then so is $\mathfrak{g}[t^{\pm 1}] =$ Laurent poly
with values in \mathfrak{g}

Quantum groups (quantum loop algebras)

$\hat{\mathfrak{g}}$

If \mathfrak{g} is a Lie algebra, then so is $\mathfrak{g}[t^{\pm 1}] =$ Laurent poly
with values in \mathfrak{g}

Has representations of the form

$$V_1(a_1) \otimes V_2(a_2) \otimes \dots \otimes V_n(a_n)$$

a representation
of \mathfrak{g}

evaluate at $t = a_1$

Quantum groups (quantum loop algebras)

$\hat{\mathfrak{g}}$

If \mathfrak{g} is a Lie algebra, then so is $\mathfrak{g}[t^{\pm 1}] =$ Laurent poly
with values in \mathfrak{g}

Has representations of the form

$$V_1(a_1) \otimes V_2(a_2) \otimes \dots \otimes V_n(a_n)$$

a representation
of \mathfrak{g}

evaluate at $t = a_1$


A quantum group $U_{\hbar}(\hat{\mathfrak{g}})$ is a deformation such that

$$V_1(a_1) \otimes V_2(a_2) \not\cong V_2(a_2) \otimes V_1(a_1)$$

R-matrices

For generic a_1/a_2 there is a **nontrivial** intertwiner

$$V_1(a_1) \otimes V_2(a_2)$$



$$V_2(a_2) \otimes V_1(a_1)$$

$$R_{V_1, V_2}(a_1/a_2)$$


rational function
of a_1/a_2

vertex interaction in integrable
vertex models

R-matrices

For generic a_1/a_2 there is a **nontrivial** intertwiner

$$V_1(a_1) \otimes V_2(a_2)$$



$$V_2(a_2) \otimes V_1(a_1)$$

$$\text{X} = \text{X} \text{YB}$$

$$R_{V_1, V_2}(a_1/a_2)$$

rational function
of a_1/a_2

vertex interaction in integrable
vertex models

R-matrices

For generic a_1/a_2 there is a **nontrivial** intertwiner

$$V_1(a_1) \otimes V_2(a_2)$$

$$\text{X} = \text{X} \text{YB}$$

$$R_{V_1, V_2}(a_1/a_2)$$

$$V_2(a_2) \otimes V_1(a_1)$$


rational function
of a_1/a_2

vertex interaction in integrable
vertex models

One can reconstruct the whole quantum group from R-matrices
[Faddeev - Reshetikhin - Takhtajan]

Quantum integrals of motion

a.k.a. Baxter subalgebra




Quantum integrals of motion

a.k.a. Baxter subalgebra in

physical space

Here $z \in e^{\mathfrak{f}} \subset \mathcal{U}_{\hbar}(\hat{\mathfrak{g}})$

where $\mathfrak{f} \subset \mathfrak{g}$ are diagonal matrices




auxilliary space

quasiperiodic boundary conditions

Quantum integrals of motion

a.k.a. Baxter subalgebra in



physical space

Here $z \in e^{\mathfrak{f}} \subset \mathcal{U}_{\hbar}(\hat{\mathfrak{g}})$

where $\mathfrak{f} \subset \mathfrak{g}$ are diagonal matrices

Commute for all W and u

for fixed z


quasiperiodic boundary conditions

The textbook Bethe ansatz diagonalizes these for

$$\mathfrak{g} = \mathfrak{sl}_2$$

$$V = \mathbb{C}^2(a_1) \otimes \dots \otimes \mathbb{C}^2(a_n)$$

spin $\frac{1}{2}$ chain
of length n



The textbook Bethe ansatz diagonalizes these for

$$\mathfrak{g} = \mathfrak{sl}_2$$

$$V = \mathbb{C}^2(a_1) \otimes \dots \otimes \mathbb{C}^2(a_n)$$

spin $\frac{1}{2}$ chain
of length n

the challenge is to do it for a very general \mathfrak{g} (including ∞ -dimensional)

The textbook Bethe ansatz diagonalizes these for

$$\mathfrak{g} = \mathfrak{sl}_2$$

$$V = \mathbb{C}^2(a_1) \otimes \dots \otimes \mathbb{C}^2(a_n)$$

spin $\frac{1}{2}$ chain
of length n

the challenge is to do it for a very general \mathfrak{g} (including ∞ -dimensional)

A more general problem is to solve certain q -difference eq. for


$$\Psi(a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)$$

unrelated to
any other
variables
used so far

concretely, the Quantum Knizhnik-Zamolodchikov eq. for

reads
$$\Psi(a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)$$

$$\Psi(q^{a_1}, \dots, a_n) = (z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$




[I. Frenkel - N. Reshetikhin]

[F. Smirnov]

concretely, the Quantum Knizhnik-Zamolodchikov eq. for

reads
$$\Psi(a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)$$

$$\Psi(q^{a_1}, \dots, a_n) = (z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$




there are commuting "dynamical" equations in z

[Etingof, Felder, Tarasov, Varchenko, ...]

concretely, the Quantum Knizhnik-Zamolodchikov eq. for

reads
$$\Psi(a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)$$

$$\Psi(q^{a_1}, \dots, a_n) = (z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$



there are commuting "dynamical" equations in z

as $q \rightarrow 1$ become an eigenvalue problem

a generalization of Bethe Ansatz is the search for
 integral solutions of the q -difference equations

depends on the representation

$$\Psi_\alpha = \int f_\alpha(x_1, x_2, \dots, a) K(x, z, a, \dots, q)$$

(Annotations: α is the index in the physical space; x_1, x_2, \dots, a are integration variables; K is a fixed kernel function.)

index in the physical space $V = \otimes V_i(a_i)$

$$e^{\frac{1}{\ln q}} S(x, z, a) + \dots$$

as $q \rightarrow 1$

studied by [Tarasov-Varchenko, ...]
 for $\mathfrak{g} = \mathfrak{sl}(n)$

in the $q \rightarrow 1$ limit, we get

$$\frac{\partial S}{\partial x_i} = 0$$

← *Bethe* equations for
= "Bethe roots" x_1, x_2, \dots

↙ Spectrum of
the problem

in the $q \rightarrow 1$ limit, we get

Spectrum of
the problem

$$\frac{\partial S}{\partial x_i} = 0$$

← Bethe equations for
= "Bethe roots" x_1, x_2, \dots

$a_i \frac{\partial S}{\partial a_i} \rightsquigarrow$ eigenvalues of the qKZ operators, etc.

1

in the $q \rightarrow 1$ limit, we get

$$\frac{\partial S}{\partial x_i} = 0$$

↔ **Bethe** equations for
= "Bethe roots" x_1, x_2, \dots

Spectrum of
the problem

$$a_i \frac{\partial S}{\partial a_i} = \text{eigenvalues of the } q\text{KZ operators, etc.}$$

and the map

$$\text{Hilbert space} \ni \alpha \mapsto f_\alpha \Big|_{\text{Bethe}} \in \text{functions on the spectrum}$$

is the diagonalization!

So, the main problem is to find

$$\text{Hilbert space } \mathcal{V} \ni \alpha \mapsto f_{\alpha}(x_1, x_2, \dots)$$

functions



"off-shell Bethe function"



name introduced by Babujian

So, the main problem is to find

$$\text{Hilbert space } \mathcal{V} \ni \alpha \mapsto f_{\alpha}(x_1, x_2, \dots)$$

functions

"off-shell Bethe function"

and this is the problem we solve in the setup

discovered by Nekrasov and Shatashvili

\mathcal{H} embeds the problem in 3d susy gauge theories on


$$M^3 = \text{Riemann surface } \mathcal{C} \times S^1$$

↑ Riemann surface \mathcal{C}

NS correspondence

$$\text{gauge group} = \prod_{i=1}^{\text{rank of}} U(v_i)$$

records the **weight** of α ,
e.g. the number of \uparrow



NS correspondence

gauge group = $\prod_{i=1}^{\text{rank of}} U(\nu_i)$

records the **weight** of α ,
e.g. the number of \uparrow

matter = $\bigoplus \mathbb{C}^{\nu_i} \otimes \text{flavor space } W_i$

here act $\begin{pmatrix} a_1 & & 0 \\ & a_2 & \\ 0 & & \ddots \end{pmatrix}$

NS correspondence

gauge group = $\prod_{i=1}^{\text{rank of } \mathfrak{g}} U(\nu_i)$

records the **weight** of α ,
e.g. the number of \uparrow

matter = $\bigoplus \mathbb{C}^{\nu_i} \otimes \text{flavor space } W_i$

here act $\begin{pmatrix} a_1 & & 0 \\ & a_2 & \\ 0 & & \dots \end{pmatrix}$


\bigoplus bifundamental $(\mathbb{C}^{\nu_i})^* \otimes \mathbb{C}^{\nu_j}$

$i \rightarrow j$

sum over arrows in a

quiver = Dynkin* diagram of \mathfrak{g}

arbitrary graph



NS correspondence

gauge group = $\prod_{i=1}^{\text{rank of}} U(\nu_i)$

records the **weight** of α ,
e.g. the number of \uparrow

matter = $\bigoplus \mathbb{C}^{\nu_i} \otimes \text{flavor space } W_i$

here act $\begin{pmatrix} a_1 & & 0 \\ & a_2 & \\ 0 & & \dots \end{pmatrix}$

\bigoplus bifundamental $(\mathbb{C}^{\nu_i})^* \otimes \mathbb{C}^{\nu_j}$


$i \rightarrow j$

sum over arrows in a

quiver = Dynkin* diagram of \mathfrak{g}

\bigoplus duals, for more susy


NS correspondence

$$M^3 = \mathbb{C}^* \times S^1$$


Hilbert space of the
quantum integrable
system

= line operators

= equivariant K-theory of




already is a module over a certain (smaller) quantum
group by the original construction of Nakajima

mathematically, the susy indices for

$$M^3 = \text{S}^2 \times S^1$$


LGSM

are integrals in K-theory of the space of (quasi-)maps




weighted by $\sum^{\text{deg } g}$ ← multiindex, a subject both formally and conceptually related to other kinds of curve counting such as Gromov-Witten theories (← top strings).

in particular, the subject has the **quantum K-theory** ring,
with structure constants given by



as well as the quantum **q**-difference equations, which
record the response to twisting the geometry over \mathbb{P}^1



one of the key insights of NS :

quantum K-theory ring = $\frac{\text{sym. polynomials in } x_{ij}}{\text{Bethe equations}}$

standard generators, Chern roots of universal bundles


See [Pushkar-Smirnov-Zeitlin] for a discussion aimed at mathematicians

· \curvearrowright Baxter's Q-operators


the problem may be studied from a geometric representation theory angle [Maulik-0.] and one of the end results of this analysis is

Theorem [A.O., A. Smirnov - A.O.] $qKZ + \text{dynamical eq} + \dots$


are the quantum difference equations for $\mathcal{M}_{\text{Higgs}}$ where the step q is the equivariant variable that rotates the domain of the quasimap




Now back to the main problem of Bethe ansatz:



Now back to the main problem of Bethe ansatz:



Now back to the main problem of Bethe ansatz:




↑
observables

in our paper, we give several equivalent descriptions of this map, some of which make no explicit reference to gauge theories


mathematically, the dictionary

$$K_{eq}(\mathcal{M}_{\text{Higgs}}) \begin{array}{c} \xrightarrow{\hspace{2cm}} \\ \xleftarrow{\hspace{2cm}} \end{array} \begin{array}{l} \text{representations} \\ \text{of the gauge group} \end{array}$$

is given by 2-point functions with



relative insertions:



Description #1

$f_{\alpha}(x)$ is the descendent observable that corresponds to the relative observable α

Description #2 $f_\alpha(x)$ is the **stable envelope**


$$K_{eq}(\mathcal{M}_{\text{Higgs}}) \longrightarrow K_{eq}\left(\left[\begin{array}{c} \text{matter} \\ \hline \text{gauge} \end{array} \right]\right)$$

ambient stack
↓


↑
no stability

in the style of Halpern-Leistner-Maulik - A.O.

This is the $q \rightarrow 0$ limit of the elliptic stable envelopes of [Aganagic-0.], which are boundary conditions for gauge theories in question



Description #3 $f_\alpha(x)$ is a matrix element of R-matrix



specific eigenvector of $U_{\mathfrak{h}}(\hat{\mathfrak{f}})$

vacuum $\leftarrow \emptyset$

this specializes to the
 $B(x_1)B(x_2)\dots |\emptyset\rangle$
 formula for $\mathfrak{g} = \mathfrak{sl}_2$

Description #4

Explicit **abelianization** formula in the style of Shenfeld-Smirnov-....

arXiv.org > math-ph > arXiv:1704.08746

Mathematical Physics

Quasimap counts and Bethe eigenfunctions

Mina Aganagic, Andrei Okounkov

(Submitted on 27 Apr 2017)

We associate an explicit equivalent descendent insertion to any relative insertion in quantum K-theory. We give an explicit formula for off-shell Bethe eigenfunctions for general quantum loop algebras associated to the corresponding quantum Knizhnik-Zamolodchikov and dynamical q-difference equations.

Subjects: **Mathematical Physics (math-ph)**; High Energy Physics - Theory (hep-th); Algebraic Geometry (math-al)
Cite as: **arXiv:1704.08746 [math-ph]**
(or **arXiv:1704.08746v1 [math-ph]** for this version)

Section 3.2

as a T-module. In particular, the A-weights in V are given by minus contents of the boxes. As a polarization, we may take

$$T^{1/2} = V + (t_1 - 1) \text{Hom}(V, V) \\ = \sum x_i + (t_1 - 1) \sum_{i,j} x_i/x_j$$

where $\{x_i\}$ are the Chern roots of V. A fixed point is specified by the assignment of x_i to the boxes of λ , up to permutation.

If we take t_1 to be a repelling weight for A then

$$T_{\geq}^{1/2} = \sum_{c(i) \geq 0} x_i + t_1 \sum_{c(i) \geq c(j)+1} x_i/x_j - \sum_{c(i) \geq c(j)} x_i/x_j$$

where

$$T_{>}^{1/2} = T_{\text{attracting}}^{1/2}, \quad T_{<}^{1/2} = T_{\text{repelling}}^{1/2}$$

and $c(i)$ is the content of the box in λ assigned to x_i . Therefore, up to an \hbar multiple, we have

$$f_{\lambda} = \text{symmetrization of } \frac{\Pi_1 \Pi_2}{\Pi_3}$$

where

$$\Pi_1 = \prod_{c(i) < 0} (1 - x_i) \prod_{c(i) > 0} (t_1 t_2 - x_i)$$

and

$$\Pi_2 = \prod_{c(i) < c(j)+1} (x_j - t_1 x_i) \prod_{c(i) > c(j)+1} (t_2 x_j - x_i)$$

$$\Pi_3 = \prod_{c(i) < c(j)} (x_j - x_i) \prod_{c(i) > c(j)} (t_1 t_2 x_j - x_i)$$

These are formulas for K-theoretic stable envelopes for $\text{Hilb}(\mathbb{C}^2, n)$ with the polarization and slope as in Proposition 7. They are a direct K-theoretic generalization of the formulas from [48, 50].

Note that in all cases treated by the formula (79) the slope is near an integral line bundle. Much more interesting functions appear at fractional slopes, but they seem to be not required in the context of Bethe Ansatz.

3.2.7

The proof of Proposition 7 takes several steps. As a first step, we clarify the geometric meaning of the formula (79).

We separate the numerator and denominator in (79) by writing

$$(T^{1/2})_{\text{repelling}} \oplus \hbar (T^{1/2})_{\text{attracting}} = \rho_+ - \rho_-$$

why is geometry effective
in proving explicit formulas like ?

for a number of both theoretical and
very practical reasons such as:

- it lets one be inexplicit about
many features of R , or $\mathcal{U}_h(\hat{o}_j)$, or
- it automatically selects the right contour of integration
- it is awfully good at showing that poles cancel
-

