

## Homework 2

Additive number theory seminar

Due March 27, 2023 by 11:40 AM

Choose **one** of the following problems and write a clear and readable solution.

**Problem 1.** Let  $k \geq 2$  be an integer, and say that a positive integer  $n$  is  $k$ -free if it is not divisible by any  $k$ th power (so for example 2-free numbers are the squarefree numbers). Show that the number of  $k$ -free numbers less than or equal to  $x$  is  $\frac{1}{\zeta(k)}x + O(x^{1/k})$ , where  $\zeta(s)$  is the Riemann zeta function. You can use without proof the fact that

$$\frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} = \prod_p \left(1 - \frac{1}{p^s}\right),$$

and the fact that

$$\sum_{d^k | n} \mu(d) = \begin{cases} 1 & n \text{ is } k\text{-free} \\ 0 & \text{otherwise} \end{cases}$$

**Problem 2.** Treating the probability of an integer  $n$  being divisible by another integer  $q$  as a random event with probability  $\frac{1}{q}$  and assuming that these events are independent, derive the following heuristics, neglecting any error terms:

- (a) The probability of a positive integer  $n$  being squarefree is  $\prod_p \left(1 - \frac{1}{p^2}\right)$ . (This product is  $\frac{6}{\pi^2} = \frac{1}{\zeta(2)}$ , so this guess is confirmed by our calculation from class.)
- (b) Using Mertens's theorem, the probability of a positive integer  $n$  being prime is  $\frac{2e^{-\gamma}}{\log n}$ , where  $\gamma$  is the Euler–Mascheroni constant. (This is contradicted by the prime number theorem, which can be interpreted as saying the probability of a random positive integer  $n$  being prime is  $\frac{1}{\log n}$ ; since  $1 \neq 2e^{-\gamma} \approx 1.1229$ , this means this sort of heuristic is not always correct! This is because in fact being divisible by different primes is not quite independent, but related in complicated ways.)

**Problem 3.** Apply Brun's sieve to show that the number of *triplet* primes up to  $x$ , i.e. numbers  $n \leq x$  such that  $n$ ,  $n+2$ , and  $n+6$  are all prime, is  $O\left(\frac{x(\log \log x)^3}{(\log x)^3}\right)$ . (If we'd instead asked for  $n$ ,  $n+2$ , and  $n+4$  to all be prime, the only example is  $n=3$ .) The error bounding is the same as for twin primes after taking  $z = e^{\frac{1}{\gamma \log \log x}}$ , so omit it and focus on the main term.