

DEGREES OF UNITARY DELIGNE–LUSZTIG VARIETIES

CHAO LI

ABSTRACT. We prove an explicit degree formula for certain unitary Deligne–Lusztig varieties. Combining with an alternative degree formula in terms of Schubert calculus, we deduce several algebraic combinatorial identities which may be of independent interest.

1. INTRODUCTION

1.1. Degrees of unitary Deligne–Lusztig varieties. Let $k_0 = \mathbb{F}_q$ be a finite field of size q . Let $k = \mathbb{F}_{q^2}$ be the quadratic extension of k_0 . Let V be a (nondegenerate) k/k_0 -hermitian space of dimension n . Let $\text{Gr}_m(V)$ be the Grassmannian of m -dimensional subspaces of V , which is a smooth projective variety over k of dimension $m(n - m)$.

We assume that V has dimension $n = 2d + 1$ ($d \geq 0$) over k and take $m = d + 1$. Define $\text{DL}(V) \subseteq \text{Gr}_{d+1}(V)$ to be the closed k -subscheme parametrizing $(d + 1)$ -dimensional subspaces U such that $U^\perp \subseteq U$, where U^\perp is the orthogonal complement of U in V (see the more precise Definition 2.1). It is smooth, projective and geometrically irreducible of dimension d , and is known as a *unitary Deligne–Lusztig variety*, as it can be identified as a generalized Deligne–Lusztig variety associated to a parabolic subgroup of the odd unitary group $\text{U}(V)$ ([VW11, §4.5]). This class of varieties shows up in the study of the supersingular locus of unitary Shimura varieties ([VW11]) and plays an important role in the arithmetic of unitary Shimura varieties, such as the Kudla–Rapoport conjecture ([KR11, KR14]).

Our first main result is a simple degree formula for $\text{DL}(V)$. Recall that the *degree* of a projective variety $X \subseteq \mathbb{P}^N$ of pure dimension d is given by the geometric intersection number of X with d general hyperplanes in \mathbb{P}^N . Denote by $\deg \text{DL}(V)$ the degree of $\text{DL}(V)$ under the Plücker embedding $\text{Gr}_{d+1}(V) \hookrightarrow \mathbb{P}(\wedge^{d+1} V)$.

Theorem 1.1. *Let V be a k/k_0 -hermitian space of dimension $n = 2d + 1$. Then*

$$(1.1.1) \quad \deg \text{DL}(V) = \prod_{i=1}^d \frac{1 - q^{2i}}{1 - q}.$$

Remark 1.2. Here (and below) we regard the empty product (when $d = 0$) as 1. The right hand side of (1.1.1) can also be interpreted as the q -analogue of the double factorial

$$[2d]_q!! := [2d]_q [2d - 2]_q \cdots [2]_q,$$

where $[n]_q = \frac{1 - q^n}{1 - q}$ is the q -analogue of n .

Theorem 1.1 will be proved in §2.6. Its proof is inspired by the *higher local modularity* in the recent proof of the Kudla–Rapoport conjecture (cf. [LZ, §6.4]). In fact, the key formulas (Propositions 2.11, 2.12) can be extracted from certain vertical intersection formulas ([LZ, Lemmas 6.4.5, 6.4.6]) on unitary Rapoport–Zink spaces. To make the ideas more transparent, here we work directly on $\mathrm{DL}(V)$ and introduce the notions of *special cycles* $Z(W) \subseteq \mathrm{DL}(V)$ (Definition 2.3) and a *special line bundle* $\mathcal{L}_{\mathrm{DL}(V)}$ (Definition 2.5) on $\mathrm{DL}(V)$. These notions may be viewed respectively as finite field analogues of special cycles and the Hodge line bundle on unitary Shimura varieties. From this perspective, the degree formula in Theorem 1.1, or more precisely the formula for $c_1(\mathcal{L}_{\mathrm{DL}(V)}^*)^d$ in Proposition 2.12, may be viewed as a finite field analogue of the constant term formula in Kudla’s geometric Siegel–Weil formula (relating the geometric volume of unitary Shimura varieties and an abelian L -value, cf. [Kud04, (4.4)] for orthogonal Shimura varieties).

The proof of Theorem 1.1 ultimately relies on identifying the special cycles $Z(W) \subseteq \mathrm{DL}(V)$ as unitary Deligne–Lusztig subvarieties and the (proved) Tate conjecture for $\mathrm{DL}(V)$, in order to perform induction on the dimension d . These ingredients are available for Deligne–Lusztig varieties beyond those of unitary types (e.g., the type 2D_n considered in [LZ21, §7.6] and two other types B_n, C_n listed in [HLZ19, §3]), and it would be interesting to extend the method to obtain degree formulas for more general Deligne–Lusztig varieties.

1.2. Schubert calculus and applications to algebraic combinatorics. We also prove a different formula for $\deg \mathrm{DL}(V)$ in terms of Schubert calculus on Grassmannians (see also the related general works [Kim20, HP20]).

Theorem 1.3. *The following identity holds in $\mathrm{Ch}^{d(d+1)}(\mathrm{Gr}_{d+1}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$:*

$$(1.3.1) \quad \deg \mathrm{DL}(V) = \sum_c \sigma_c \sigma_{\hat{c}} \sigma_1^d q^{|c|},$$

where the sum runs over all integer tuples $c = (c_1, \dots, c_d)$ with $d \geq c_1 \geq \dots \geq c_d \geq 0$. Here (as recalled in §3.1):

- $|c| = c_1 + \dots + c_d$,
- $\sigma_c \in \mathrm{Ch}^{|c|}(\mathrm{Gr}_{d+1}(V)_{\bar{k}})_{\mathbb{Q}}$ is the Schubert class,
- \hat{c} is the complement of c defined by $\hat{c} := (d - c_d, \dots, d - c_1)$,
- \hat{c}' is the conjugate of \hat{c} .

Combining Theorems 1.1 and 1.3 we deduce several combinatorial identities.

Corollary 1.4. (i) *The right hand side of (1.3.1) is equal to $\prod_{i=1}^d \frac{1-q^{2i}}{1-q}$.*

(ii) *The $x_1^{2d} x_2^{2d-1} \dots x_d^{d+1}$ -coefficient of*

$$\left(\prod_{i,j=1}^d (qx_i + x_j) \right) (x_1 + \dots + x_d)^d \left(\prod_{i<j} (x_i - x_j) \right)$$

is equal to $\prod_{i=1}^d \frac{1-q^{2i}}{1-q}$.

(iii) *Let $l \geq 0$. The following two sets have the same size:*

- The set of standard Young tableaux of skew shape $(\tilde{c}')^*/c$, where c runs over (c_1, \dots, c_d) such that $d \geq c_1 \geq \dots \geq c_d \geq 0$ and $|c| = l$. Here $(\tilde{c}')^*$ is the dual of \tilde{c}' (recalled in §3.1).
- The set of ordered partitions (l_1, \dots, l_d) of l satisfying $l = l_1 + \dots + l_d$ and $0 \leq l_i \leq 2i - 1$ for all $1 \leq i \leq d$.

Theorem 1.3 and Corollary 1.4 will be proved §3.3 and §3.4 respectively. We notice that the combinatorial identities in Corollary 1.4 seem to be quite nontrivial even for small values of d (see Example 3.4). It would be very interesting to find more direct combinatorial proofs of these identities.

1.3. Acknowledgments. It is the author's pleasure to dedicate this paper to Steve Kudla on the occasion of his 70th birthday. The influence of his original insights on the geometric and arithmetic Siegel–Weil formula on this paper should be evident to the readers. The author is also grateful to Z. Yun and W. Zhang for helpful conversations. The author's work is partially supported by the NSF grant DMS-2101157.

1.4. Notation. Let X be a smooth projective variety over a finite field k . Let $\text{Ch}^r(X_{\bar{k}})$ be the Chow group of codimension r algebraic cycles of $X_{\bar{k}}$ defined modulo rational equivalence. For any prime $\ell \neq \text{char}(k)$, denote by

$$\text{cl}_r : \text{Ch}^r(X_{\bar{k}})_{\mathbb{Q}} \rightarrow \text{H}^{2r}(X_{\bar{k}}, \mathbb{Q}_{\ell})(r)$$

the ℓ -adic cycle class map. Let $\text{Ch}_{\text{hom}}^r(X_{\bar{k}})$ be the quotient of $\text{Ch}^r(X_{\bar{k}})$ modulo homological equivalence. Then cl_r factors through $\text{Ch}_{\text{hom}}^r(X_{\bar{k}})$. Denote by $\text{Tate}_{\ell}^{2r}(X_{\bar{k}}) \subseteq \text{H}^{2r}(X_{\bar{k}}, \mathbb{Q}_{\ell})(r)$ the subspace of Tate classes, i.e., the elements fixed by an open subgroup of $\text{Gal}(\bar{k}/k)$. Then cl_r intertwines the intersection product \cdot on the Chow ring and the cup product \cup on the cohomology ring, namely the following diagram commutes,

$$(1.4.1) \quad \begin{array}{ccccc} \text{Ch}^r(X_{\bar{k}})_{\mathbb{Q}} & \times & \text{Ch}^s(X_{\bar{k}})_{\mathbb{Q}} & \longrightarrow & \text{Ch}^{r+s}(X_{\bar{k}})_{\mathbb{Q}} \\ \downarrow \text{cl}_r & & \downarrow \text{cl}_s & & \downarrow \text{cl}_{r+s} \\ \text{H}^{2r}(X_{\bar{k}}, \mathbb{Q}_{\ell})(r) & \times & \text{H}^{2s}(X_{\bar{k}}, \mathbb{Q}_{\ell})(s) & \xrightarrow{\cup} & \text{H}^{2(r+s)}(X_{\bar{k}}, \mathbb{Q}_{\ell})(r+s). \end{array}$$

When $r = 1$, we have $\text{Ch}^1(X_{\bar{k}})_{\mathbb{Q}} = \text{Ch}_{\text{hom}}^1(X_{\bar{k}})_{\mathbb{Q}}$. When $r = \dim X$, we often identify $\text{Ch}^{\dim X}(X_{\bar{k}})_{\mathbb{Q}} = \text{Ch}_{\text{hom}}^{\dim X}(X_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$ via the degree isomorphism (cf. [Ful98, Example 1.6.6]).

Recall that the Tate conjecture ([Tat65, Conjecture 1], or [Tat94, Conjecture T^i]) asserts that for any $r \geq 0$ and for any prime $\ell \neq \text{char}(k)$, the image of ℓ -adic cycle class map cl_r is equal to $\text{Tate}_{\ell}^{2r}(X_{\bar{k}})$ (in particular, induces an isomorphism $\text{Ch}_{\text{hom}}^r(X_{\bar{k}}) \otimes \mathbb{Q}_{\ell} \simeq \text{Tate}_{\ell}^{2r}(X_{\bar{k}})$).

For a subvariety $Z \subseteq X$ of pure codimension r , we denote by $[Z]_X$ its class in $\text{Ch}^r(X_{\bar{k}})$. When the ambient variety X is clear we suppress the subscript and simply write $[Z]$. For any vector bundle \mathcal{V} on X , denote by $c_r(\mathcal{V}) \in \text{Ch}^r(X_{\bar{k}})$ its r -th Chern class. By abusing notation we also denote by the same symbols $[Z]_X$, $[Z]$ and $c_r(\mathcal{V})$ their images in $\text{Ch}^r(X_{\bar{k}})_{\mathbb{Q}}$ or $\text{Ch}_{\text{hom}}^r(X_{\bar{k}})_{\mathbb{Q}}$ (which makes no difference when computing intersection products by the commutativity of the diagram (1.4.1)). For any vector bundle \mathcal{V} on X , denote by $\mathcal{V}^* := \text{Hom}_{\mathcal{O}_X}(\mathcal{V}, \mathcal{O}_X)$ its dual bundle.

2. DEGREES OF UNITARY DELIGNE–LUSZTIG VARIETIES

Let p be a prime and let q be a power of p . Let $k_0 = \mathbb{F}_q$ be a finite field of size q . Let $k = \mathbb{F}_{q^2}$ be the quadratic extension of k_0 . Denote by σ the absolute q -Frobenius endomorphism on any scheme of characteristic p .

2.1. Hermitian spaces. Let V be a (nondegenerate) k/k_0 -hermitian space of dimension n , i.e., a k -vector space of dimension n equipped with a (nondegenerate) pairing $(\ , \) : V \times V \rightarrow k$ that is linear in the first variable, σ -linear in the second variable and satisfies

$$(2.0.2) \quad (x, y) = (y, x)^\sigma$$

for any $x, y \in V$. For any k -subspace $U \subseteq V$, denote by

$$U^\perp = \{x \in V : (x, U) = 0\}$$

its orthogonal complement.

More generally, for any k -scheme S , put $V_S := V \otimes_k \mathcal{O}_S$. Then there is a unique pairing $(\ , \)_S : V_S \times V_S \rightarrow \mathcal{O}_S$ extending $(\ , \)$ that is \mathcal{O}_S -linear in the first variable and (\mathcal{O}_S, σ) -linear in the second variable, given by

$$(x \otimes \lambda, y \otimes \mu) = \lambda \mu^\sigma (x, y)$$

for any $x, y \in V$ and $\lambda, \mu \in \mathcal{O}_S$. For any subbundle $U \subseteq V_S$, we denote by

$$U^\perp = \{x \in V_S : (x, U) = 0\}$$

its left orthogonal complement. (Notice that unlike the case $S = \text{Spec } k$, in general the symmetry (2.0.2) does not necessarily hold for $x, y \in V_S$, and the left orthogonal complement U^\perp does not necessarily agree with the right orthogonal complement.)

2.2. Unitary Deligne–Lusztig varieties and special cycles. From now on fix a k/k_0 -hermitian space V of dimension $n = 2d + 1$ ($d \geq 0$).

Definition 2.1. Define the *unitary Deligne–Lusztig variety* $\text{DL}(V) \subseteq \text{Gr}_{d+1}(V)$ be the closed k -subscheme parametrizing $(d+1)$ -dimensional subspaces U such that $U^\perp \subseteq U$, i.e., for any k -scheme S ,

$$\text{DL}(V)(S) = \{\text{subbundles } U \subseteq V_S : \text{rank } U = d + 1, U^\perp \subseteq U\}.$$

Definition 2.2. Define a *special subspace* $W \subseteq V$ to be a k -subspace such that $W^\perp \subseteq W$. If W is a special subspace, then W/W^\perp is a (nondegenerate) k/k_0 -hermitian space under the pairing induced from V .

Definition 2.3. Let $W \subseteq V$ be a k -subspace. Define the *special cycle* $Z(W) \subseteq \text{DL}(V)$ be the closed subscheme parametrizing subspaces U satisfying $U \subseteq W$, i.e., for any k -scheme S ,

$$Z(W)(S) = \{U \in \text{DL}(V)(S) : U \subseteq W_S\}.$$

By definition it is clear that

- $Z(W)$ is nonempty only if $W \subseteq V$ is a *special subspace*.
- $Z(V) = \text{DL}(V)$,

- $Z(W) \cap Z(W') = Z(W \cap W')$ for two subspaces $W, W' \subseteq V$,
- $Z(W) \subseteq Z(W')$ if $W \subseteq W'$.

We summarize several known results on $\mathrm{DL}(V)$ and $Z(W)$ which we will need later.

Proposition 2.4. (i) $\mathrm{DL}(V)$ is smooth, projective and geometrically irreducible of dimension d .

(ii) Let $W \subseteq V$ be a special subspace of codimension r . Then there is a canonical isomorphism of k -schemes

$$(2.4.1) \quad Z(W) \simeq \mathrm{DL}(W/W^\perp).$$

In particular, $Z(W)$ is smooth, projective and geometrically irreducible, and has codimension r in $\mathrm{DL}(V)$, and we call $Z(W)$ a codimension r special cycle in $\mathrm{DL}(V)$.

(iii) The Tate conjecture (see §1.4) holds for $\mathrm{DL}(V)$.

(iv) For any $r \geq 0$, the space $\mathrm{Ch}_{\mathrm{hom}}^r(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}}$ is generated by the codimension r special cycles $[Z(W)]$, where $W \subseteq V$ runs over all special subspace of codimension r .

(v) For any $r \geq 0$, the intersection pairing induces a perfect pairing

$$\mathrm{Ch}_{\mathrm{hom}}^r(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}} \times \mathrm{Ch}_{\mathrm{hom}}^{d-r}(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}} \xrightarrow{\cdot} \mathbb{Q}.$$

Proof. (i) This is [VW11, Lemma 4.5], or the more general [LTX⁺19, Proposition A.1.3 (2)].

(ii) Consider the morphism $Z(W) \rightarrow \mathrm{DL}(W/W^\perp)$ defined by

$$Z(W)(S) \rightarrow \mathrm{DL}(W/W^\perp)(S), \quad U \mapsto \bar{U} = U/W_S^\perp$$

for any k -scheme S . It is an isomorphism, with inverse given by the morphism defined by $\bar{U} \mapsto W_S^\perp + \bar{U}$. The rest follows from Item (i) since W/W^\perp has k -dimension $n - 2r$.

(iii) This is [LZ, Theorem 5.3.2 (i)].

(iv) This is a combination of [LZ, Theorem 5.3.2 (i)] and [Vol10, Corollary 2.17].

(v) By [LZ, (6.4.0.4)], the cup product induces a perfect pairing

$$\mathrm{Tate}_\ell^{2r}(\mathrm{DL}(V)_{\bar{k}}) \times \mathrm{Tate}_\ell^{2d-2r}(\mathrm{DL}(V)_{\bar{k}}) \rightarrow \mathbb{Q}_\ell.$$

The result then follows from the Tate conjecture and the commutativity of (1.4.1). □

2.3. Natural vector bundles on $\mathrm{DL}(V)$ and $Z(W)$.

Definition 2.5. Define

- $\mathcal{V}_{\mathrm{DL}(V)} := V \otimes_k \mathcal{O}_{\mathrm{DL}(V)}$ the universal bundle on $\mathrm{DL}(V)$,
- $\mathcal{U}_{\mathrm{DL}(V)} \subseteq \mathcal{V}_{\mathrm{DL}(V)}$ and $\mathcal{U}_{\mathrm{DL}(V)}^\perp \subseteq \mathcal{V}_{\mathrm{DL}(V)}$ the two universal subbundles on $\mathrm{DL}(V)$,
- $\mathcal{Q}_{\mathrm{DL}(V)} := \mathcal{V}_{\mathrm{DL}(V)}/\mathcal{U}_{\mathrm{DL}(V)}$ the universal quotient bundle on $\mathrm{DL}(V)$,
- $\mathcal{L}_{\mathrm{DL}(V)} := \mathcal{U}_{\mathrm{DL}(V)}/\mathcal{U}_{\mathrm{DL}(V)}^\perp$, which we call the *special line bundle* on $\mathrm{DL}(V)$.

Similarly for a special cycle $Z(W) \subseteq \mathrm{DL}(V)$ associated to a special subspace $W \subseteq V$, define

- $\mathcal{W}_{Z(W)} := (W/W^\perp) \otimes_k \mathcal{O}_{Z(W)}$ the universal bundle on $Z(W) \simeq \mathrm{DL}(W/W^\perp)$,
- $\mathcal{U}_{Z(W)} \subseteq \mathcal{W}_{Z(W)}$ and $\mathcal{U}_{Z(W)}^\perp \subseteq \mathcal{W}_{Z(W)}$ the two universal subbundles on $Z(W)$,

- $\mathcal{Q}_{Z(W)} := \mathcal{W}_{Z(W)}/\mathcal{U}_{Z(W)}$ the universal quotient bundle on $Z(W)$,
- $\mathcal{L}_{Z(W)} := \mathcal{U}_{Z(W)}/\mathcal{U}_{Z(W)}^\perp$, which we call the *special line bundle* on $Z(W)$.

Denote by

$$j_W : Z(W) \hookrightarrow \text{DL}(V)$$

the natural closed embedding. By definition, we have

$$(2.5.1) \quad \mathcal{L}_{Z(W)} = j_W^* \mathcal{L}_{\text{DL}(V)}.$$

Also by definition, the canonical isomorphism (2.4.1) induces a canonical isomorphism $\mathcal{L}_{Z(W)} \simeq \mathcal{L}_{\text{DL}(W/W^\perp)}$ and thus

$$(2.5.2) \quad j_W^* \mathcal{L}_{\text{DL}(V)} \simeq \mathcal{L}_{\text{DL}(W/W^\perp)}.$$

Proposition 2.6. *Let $W \subseteq V$ be a special subspace of codimension r .*

(i) *We have a canonical isomorphism*

$$(2.6.1) \quad j_W^* \mathcal{Q}_{\text{DL}(V)}/\mathcal{Q}_{Z(W)} \simeq (V/W) \otimes_k \mathcal{O}_{Z(W)}.$$

(ii) *Let $\mathcal{N}_{Z(W)/\text{DL}(V)}$ be the normal bundle of $Z(W)$ in $\text{DL}(V)$. Then we have an isomorphism*

$$\mathcal{N}_{Z(W)/\text{DL}(V)} \simeq \left(\mathcal{L}_{Z(W)}^{\oplus r} \right)^*.$$

In particular, when $W \subseteq V$ has codimension 1, we have an isomorphism of line bundles

$$(2.6.2) \quad \mathcal{N}_{Z(W)/\text{DL}(V)} \simeq \mathcal{L}_{Z(W)}^*.$$

Proof. (i) The result follows immediately from the definition of $\mathcal{Q}_{\text{DL}(V)}$ and $\mathcal{Q}_{Z(W)}$.

(ii) By [LTX⁺19, Proposition A.1.3 (2)], we have canonical isomorphisms for the tangent bundles

$$\mathcal{T}_{\text{DL}(V)/k} \simeq \mathcal{H}om(\mathcal{L}_{\text{DL}(V)}, \mathcal{Q}_{\text{DL}(V)}), \quad \mathcal{T}_{Z(W)/k} \simeq \mathcal{H}om(\mathcal{L}_{Z(W)}, \mathcal{Q}_{Z(W)}).$$

Hence by the exact sequence

$$0 \rightarrow \mathcal{T}_{Z(W)/k} \rightarrow j_W^* \mathcal{T}_{\text{DL}(V)/k} \rightarrow \mathcal{N}_{Z(W)/\text{DL}(V)} \rightarrow 0$$

and (2.5.1), we obtain a canonical isomorphism

$$\mathcal{N}_{Z(W)/\text{DL}(V)} \simeq \mathcal{H}om(\mathcal{L}_{Z(W)}, j_W^* \mathcal{Q}_{\text{DL}(V)}/\mathcal{Q}_{Z(W)}).$$

The result then follows from (2.6.1) as $(V/W) \otimes_k \mathcal{O}_{Z(W)} \simeq \mathcal{O}_{Z(W)}^r$. □

2.4. Relations with Grassmannians. For $0 \leq m \leq n$, let $\text{Gr}_m(V)$ be the Grassmannian of m -dimensional subspaces of V , which is a smooth projective variety over k of dimension $m(n-m)$. We have the *Plücker embedding*

$$\text{Pl} : \text{Gr}_m(V) \hookrightarrow \mathbb{P}(\wedge^m(V)) \simeq \mathbb{P}^N, \quad N = \binom{n}{m} - 1,$$

defined by sending an m -dimensional subspace with basis $\{e_1, \dots, e_m\}$ to the line generated by $e_1 \wedge \dots \wedge e_m \in \wedge^m(V)$ (independent of the choice of the basis).

Definition 2.7. Define $\mathcal{S}_{\text{Gr}_{d+1}(V)}$ (resp. $\mathcal{Q}_{\text{Gr}_{d+1}(V)}$) to be the universal subbundle (resp. universal quotient subbundle) on $\text{Gr}_{d+1}(V)$. Denote by

$$i : \text{DL}(V) \hookrightarrow \text{Gr}_{d+1}(V)$$

the natural closed embedding.

Lemma 2.8. *The following identities holds:*

- (i) $\mathcal{U}_{\text{DL}(V)} = i^* \mathcal{S}_{\text{Gr}_{d+1}(V)}$,
- (ii) $\mathcal{U}_{\text{DL}(V)}^\perp \simeq i^*(\sigma^* \mathcal{Q}_{\text{Gr}_{d+1}(V)})^*$,
- (iii) $c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}) = -c_1(\mathcal{Q}_{\text{Gr}_{d+1}(V)})$,
- (iv) $c_1(\mathcal{L}_{\text{DL}(V)}) = (1 - q)i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)})$.

Proof. (i) It follows from the definition of $\mathcal{S}_{\text{Gr}_{d+1}(V)}$ and $\mathcal{U}_{\text{DL}(V)}$.

(ii) It follows from Item (i) and the nondegenerate pairing $(\ , \)$ (σ -linear in the second variable).

(iii) It follows from the defining exact sequence

$$0 \rightarrow \mathcal{S}_{\text{Gr}_{d+1}(V)} \rightarrow (V \otimes_k \mathcal{O}_{\text{Gr}_{d+1}(V)}) \rightarrow \mathcal{Q}_{\text{Gr}_{d+1}(V)} \rightarrow 0.$$

(iv) By the definition of $\mathcal{L}_{\text{DL}(V)}$ together with Item (i) and Item (ii), we have

$$c_1(\mathcal{L}_{\text{DL}(V)}) = c_1(\mathcal{U}_{\text{DL}(V)}) - c_1(\mathcal{U}_{\text{DL}(V)}^\perp) = i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}) - i^* c_1((\sigma^* \mathcal{Q}_{\text{Gr}_{d+1}(V)})^*).$$

By Item (iii), this evaluates to

$$i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}) - q(i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)})) = (1 - q)i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}),$$

as desired. □

Example 2.9. When $d = 1$ (i.e., when $\dim V = 3$), we have $\text{Gr}_{d+1}(V) \simeq \mathbb{P}^2$ with $\mathcal{Q}_{\text{Gr}_{d+1}(V)} \simeq \mathcal{O}_{\mathbb{P}^2}(1)$. Thus $c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}) = c_1(\mathcal{O}_{\mathbb{P}^2}(-1))$ by Lemma 2.8 (iii). By definition $\text{DL}(V) \subseteq \text{Gr}_{d+1}(V)$ is isomorphic to a Fermat curve of degree $1 + q$ ([Vol10, Remark 4.7]),

$$\{[x, y, z] \in \mathbb{P}^2 : x^{q+1} + y^{q+1} + z^{q+1} = 0\} \subseteq \mathbb{P}^2.$$

Hence by Lemma 2.8 (iv), the following identity holds in $\text{Ch}^1(\text{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$,

$$c_1(\mathcal{L}_{\text{DL}(V)}) = (1 - q)i^* c_1(\mathcal{O}_{\mathbb{P}^2}(-1)) = -(1 - q)(1 + q) = -(1 - q^2),$$

and so

$$(2.9.1) \quad c_1(\mathcal{L}_{\text{DL}(V)}^*) = -c_1(\mathcal{L}_{\text{DL}(V)}) = 1 - q^2.$$

2.5. Chern classes of the special line bundle.

Lemma 2.10. *Let $Z(W) \subseteq \mathrm{DL}(V)$ be a 1-dimensional special cycle associated to a special subspace $W \subseteq V$ of codimension $d - 1$. Then the following identity holds in $\mathrm{Ch}^d(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$:*

$$c_1(\mathcal{L}_{\mathrm{DL}(V)}^*) \cdot [Z(W)] = 1 - q^2.$$

Proof. By the projection formula we know that

$$c_1(\mathcal{L}_{\mathrm{DL}(V)}^*) \cdot [Z(W)] = j_{W,*}(c_1(j_W^* \mathcal{L}_{\mathrm{DL}(V)}^*)).$$

By (2.5.2), we have (under the isomorphism (2.4.1))

$$c_1(j_W^* \mathcal{L}_{\mathrm{DL}(V)}^*) = c_1(\mathcal{L}_{\mathrm{DL}(W/W^\perp)}^*).$$

As W/W^\perp is of dimension 3, we know the latter evaluates to $1 - q^2$ in $\mathrm{Ch}^1(\mathrm{DL}(W/W^\perp)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$ by (2.9.1). The result then follows. \square

Proposition 2.11. *The following identity holds in $\mathrm{Ch}^1(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}}$:*

$$(2.11.1) \quad c_1(\mathcal{L}_{\mathrm{DL}(V)}^*) = \frac{1 - q^2}{1 + q^{2d+1}} \sum_{\mathrm{codim} W=1} [Z(W)],$$

here the sum runs over all special subspaces $W \subseteq V$ of codimension 1.

Proof. When $d = 1$, the number of special subspaces $W \subseteq V$ is equal to the number of isotropic lines in the 3-dimensional k/k_0 -hermitian space V , which is $1 + q^3$ (e.g., by [LZ, Lemma 1.9.1]). Hence the right hand side of (2.11.1) evaluates to

$$\frac{1 - q^2}{1 + q^3} \cdot (1 + q^3) = 1 - q^2$$

in $\mathrm{Ch}^1(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$, which agrees with the left hand side by (2.9.1).

When $d > 1$, by Proposition 2.4 (iv) (v), to prove (2.11.1) it suffices to show that for any 1-dimensional special cycle $Z(W') \subseteq \mathrm{DL}(V)$ (associated to any codimension $d - 1$ special subspace $W' \subseteq V$), the following identity holds in $\mathrm{Ch}^d(\mathrm{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$,

$$(2.11.2) \quad c_1(\mathcal{L}_{\mathrm{DL}(V)}^*) \cdot [Z(W')] = \frac{1 - q^2}{1 + q^{2d+1}} \sum_{\mathrm{codim} W=1} [Z(W)] \cdot [Z(W')].$$

For the terms of the right hand side of (2.11.2), we have three cases.

(i) When $W' \subseteq W$, we have $Z(W') \subseteq Z(W)$. By the excess intersection formula [Ful98, Corollary 6.3], we know that

$$[Z(W)] \cdot [Z(W')] = j_{W,*}(c_1(\mathcal{N}_{Z(W')/Z(W)}) \cdot [Z(W')]_{Z(W)}).$$

By (2.6.2) and (2.5.2) we have (under the isomorphism (2.4.1))

$$c_1(\mathcal{N}_{Z(W')/Z(W)}) \cdot [Z(W')]_{Z(W)} = c_1(\mathcal{L}_{\mathrm{DL}(W/W^\perp)}^*) \cdot [Z(W')]_{\mathrm{DL}(W/W^\perp)},$$

which evaluates to $1 - q^2$ in $\mathrm{Ch}^{d-1}(\mathrm{DL}(W/W^\perp)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$ by Lemma 2.10 applied to the 1-dimensional special cycle $Z(W') \subseteq \mathrm{DL}(W/W^\perp)$. Thus in this case

$$[Z(W)] \cdot [Z(W')] = 1 - q^2.$$

(ii) When $W' \not\subseteq W$ and $W' \cap W \subseteq V$ is a special subspace, we know that $W' \cap W$ has codimension d in V and thus $Z(W) \cap Z(W') = Z(W' \cap W)$ is a single k -point by Proposition 2.4 (i), and thus

$$[Z(W)] \cdot [Z(W')] = 1.$$

(iii) When $W' \not\subseteq W$ and $W' \cap W \subseteq V$ is not a special subspace, we know that $Z(W) \cap Z(W') = Z(W' \cap W)$ is empty, and thus

$$[Z(W)] \cdot [Z(W')] = 0.$$

Now we count the number of terms on the right hand side of (2.11.2) in first two cases.

- The association $W \mapsto W^\perp$ gives a bijection between the set of codimension 1 special subspaces $W \subseteq V$ in Case (i) and the set of isotropic k -lines in W'^\perp . Hence the number of terms in Case (i) is equal to $\frac{1-q^{2(d-1)}}{1-q^2}$, as W'^\perp is a totally isotropic k/k_0 -hermitian space of dimension $d-1$.
- The association $W \mapsto W^\perp$ gives a bijection between the set of codimension 1 special subspaces $W \subseteq V$ in Case (ii) and the set of isotropic k -lines in $W' \setminus W'^\perp$. Hence the number of terms in Case (ii) is equal to the number of vectors in W'^\perp times the number of isotropic lines in the k/k_0 -hermitian space W'/W'^\perp . This is $q^{2(d-1)}(1+q^3)$, as W'^\perp is of dimension $d-1$ and W'/W'^\perp is of dimension 3.

Thus the right hand side of (2.11.2) evaluates to

$$\frac{1-q^2}{1+q^{2d+1}} \cdot \left((1-q^2) \cdot \frac{1-q^{2(d-1)}}{1-q^2} + 1 \cdot q^{2(d-1)}(1+q^3) \right) = 1-q^2,$$

which is equal to the left hand side by Lemma 2.10 applied to the 1-dimensional special cycle $Z(W') \subseteq \text{DL}(V)$. \square

Proposition 2.12. *The following identity holds in $\text{Ch}^d(\text{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$:*

$$c_1(\mathcal{L}_{\text{DL}(V)}^*)^d = \prod_{i=1}^d (1-q^{2i}).$$

Proof. When $d=1$, this is (2.9.1). In general, we induct on d . By Proposition 2.11, we obtain

$$(2.12.1) \quad c_1(\mathcal{L}_{\text{DL}(V)}^*)^d = \frac{1-q^2}{1+q^{2d+1}} \cdot c_1(\mathcal{L}_{\text{DL}(V)}^*)^{d-1} \cdot \sum_{\text{codim } W=1} [Z(W)].$$

By the projection formula and (2.5.2), we have (under the isomorphism (2.4.1)),

$$c_1(\mathcal{L}_{\text{DL}(V)}^*)^{d-1} \cdot [Z(W)] = j_{W,*}(c_1(\mathcal{L}_{\text{DL}(W/W^\perp)}^*)^{d-1}),$$

which evaluates to $\prod_{i=1}^{d-1} (1-q^{2i})$ by the induction hypothesis.

The association $W \mapsto W^\perp$ gives a bijection the set of codimension 1 special subspaces $W \subseteq V$ and the set of isotropic k -lines in V . Thus the total number of terms in (2.12.1) is the number of isotropic k -lines in the $(2d+1)$ -dimensional k/k_0 -hermitian space V , which is $\frac{(1+q^{2d+1})(1-q^{2d})}{1-q^2}$ (e.g., by [LZ, Lemma 1.9.1]). Thus (2.12.1) evaluates to

$$\frac{1-q^2}{1+q^{2d+1}} \cdot \prod_{i=1}^{d-1} (1-q^{2i}) \cdot \frac{(1+q^{2d+1})(1-q^{2d})}{1-q^2} = \prod_{i=1}^d (1-q^{2i}).$$

This completes the proof. \square

2.6. Proof of Theorem 1.1. Recall $\text{Pl} : \text{Gr}_{d+1}(V) \rightarrow \mathbb{P}(\wedge^{d+1}V)$ is the Plücker embedding. By definition

$$(2.12.2) \quad \deg \text{DL}(V) = [\text{DL}(V)]_{\mathbb{P}(\wedge^{d+1}V)} \cdot c_1(\mathcal{O}_{\mathbb{P}(\wedge^{d+1}V)}(1))^d$$

is the intersection number of $\text{DL}(V)$ with d general hyperplanes in $\mathbb{P}(\wedge^{d+1}V)$. By the projection formula, we obtain that

$$\deg \text{DL}(V) = c_1((\text{Pl} \circ i)^* \mathcal{O}_{\mathbb{P}(\wedge^{d+1}V)}(1))^d$$

in $\text{Ch}^d(\text{DL}(V)_{\bar{k}})_{\mathbb{Q}} \simeq \mathbb{Q}$. By the definition of the Plücker embedding, we have

$$\text{Pl}^* \mathcal{O}_{\mathbb{P}(\wedge^{d+1}V)}(1) \simeq \det \mathcal{S}_{\text{Gr}_{d+1}(V)}^*.$$

Thus

$$c_1((\text{Pl} \circ i)^* \mathcal{O}_{\mathbb{P}(\wedge^{d+1}V)}(1)) = i^* c_1(\det \mathcal{S}_{\text{Gr}_{d+1}(V)}^*) = i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}^*).$$

By Lemma 2.8 (iv), we have

$$i^* c_1(\mathcal{S}_{\text{Gr}_{d+1}(V)}^*) = \frac{c_1(\mathcal{L}_{\text{DL}(V)}^*)}{1 - q},$$

and hence

$$\deg \text{DL}(V) = \frac{c_1(\mathcal{L}_{\text{DL}(V)}^*)^d}{(1 - q)^d}.$$

The result then follows from Proposition 2.12.

3. SCHUBERT CALCULUS

3.1. Reminder on Schubert calculus (cf. [EH16, Chapter 4], [Ful98, §14.7]). Let $0 \leq m \leq n$. The Schubert classes of $\text{Gr}_m(V)$ are indexed by m -tuples $a = (a_1, \dots, a_m)$ of integers satisfying

$$n - m \geq a_1 \geq a_2 \geq \dots \geq a_m \geq 0,$$

in other words, indexed by Young diagrams inside the $m \times (n - m)$ rectangles. For such an m -tuple a , define a *Schubert cycle*

$$\Sigma_a(V_{\bullet}) := \{U \in \text{Gr}_m(V) : \dim U \cap V_{n-m+i-a_i} \geq i, i = 1, \dots, m\} \subseteq \text{Gr}_m(V),$$

where

$$V_{\bullet} : 0 \subset V_1 \subset V_2 \subset \dots \subset V_{n-1} \subset V_n := V$$

is a complete flag in V . The Schubert cycle $\Sigma_a(V_{\bullet}) \subseteq \text{Gr}_m(V)$ is a closed subvariety of codimension $|a| := \sum_{i=1}^m a_i$. Define the *Schubert class*

$$\sigma_a := [\Sigma_a(V_{\bullet})] \in \text{Ch}^{|a|}(\text{Gr}_m(V)_{\bar{k}}),$$

which is independent of the choice of the complete flag V_{\bullet} . We use the standard notation suppressing trailing zeros in the indices. In particular, by definition $\sigma_1 = \sigma_{1,0,\dots,0} \in \text{Ch}^1(\text{Gr}_m(V)_{\bar{k}})$ is the hyperplane class under the Plücker embedding.

Define the *dual* $a^* := (n - m - a_m, \dots, n - m - a_1)$. The Schubert classes form a \mathbb{Q} -basis of $\text{Ch}^*(\text{Gr}_m(V)_{\bar{k}})_{\mathbb{Q}}$ and the intersection pairing

$$\text{Ch}^r(\text{Gr}_m(V)_{\bar{k}})_{\mathbb{Q}} \times \text{Ch}^{m(n-m)-r}(\text{Gr}_m(V)_{\bar{k}})_{\mathbb{Q}} \rightarrow \mathbb{Q}$$

is perfect and has Schubert classes as dual basis, with σ_a and σ_b dual to each other if and only if $b = a^*$.

Define the *conjugate* $a' := (a'_1, \dots, a'_{n-m})$ such that a'_j is the number of i 's such that $a_i \geq j$. The Young diagrams of a and a' are mutual reflections along the main diagonal. The canonical isomorphism

$$(3.0.3) \quad \mathrm{Gr}_m(V) \simeq \mathrm{Gr}_{n-m}(V^*), \quad U \mapsto (V/U)^*$$

maps $\sigma_a \in \mathrm{Ch}^{|a|}(\mathrm{Gr}_m(V))$ to $\sigma_{a'} \in \mathrm{Ch}^{|a|}(\mathrm{Gr}_{n-m}(V^*))$.

3.2. The class of DL(V). Let $\mathrm{Gr}_{d,d+1}(V)$ be the variety parametrizing partial flags

$$U_\bullet : 0 \subseteq U_d \subseteq U_{d+1} \subseteq V,$$

where $\dim U_d = d$ and $\dim U_{d+1} = d + 1$. Define a closed embedding

$$\psi : \mathrm{Gr}_{d,d+1}(V) \rightarrow \mathrm{Gr}_d(V) \times \mathrm{Gr}_{d+1}(V), \quad U_\bullet \mapsto (U_d, U_{d+1}).$$

Also consider the closed embedding

$$(\phi, \mathrm{id}) : \mathrm{Gr}_{d+1}(V) \rightarrow \mathrm{Gr}_d(V) \times \mathrm{Gr}_{d+1}(V), \quad U \mapsto (U^\perp, U).$$

Then by definition (ϕ, id) induces an isomorphism between $\mathrm{DL}(V)$ and $\mathrm{im}(\phi, \mathrm{id}) \cap \mathrm{im}(\psi)$.

Proposition 3.1. *The following identity holds in $\mathrm{Ch}^*(\mathrm{Gr}_d(V)_{\bar{k}} \times \mathrm{Gr}_{d+1}(V)_{\bar{k}})_{\mathbb{Q}}$:*

$$[\mathrm{im} \psi] = \sum_{a,b} \sigma_{a^*} \times \sigma_{b^*},$$

where the sum runs over

- $a = (a_1, \dots, a_d)$ satisfying $d + 1 \geq a_1 \geq \dots \geq a_d \geq 0$,
- $b = (b_1, \dots, b_{d+1})$ satisfying $b_1 = d$, $b_i = d + 1 - a_{d+2-i} \geq 0$ for $i = 2, \dots, d + 1$.

Proof. Let V_\bullet and W_\bullet be two transverse complete flags in V ([EH16, Definition 4.4]). Let

$$V^{(i)} := V_{n-d+i-a_i}, \quad W^{(i)} := W_{n-(d+1)+i-b_i}.$$

Then by definition

$$\Sigma_a(V_\bullet) = \{U_d \in \mathrm{Gr}_d(V) : \dim U_d \cap V^{(i)} \geq i, \quad i = 1, \dots, d\}$$

and

$$\Sigma_b(W_\bullet) = \{U_{d+1} \in \mathrm{Gr}_{d+1}(V) : \dim U_{d+1} \cap W^{(i)} \geq i, \quad i = 1, \dots, d + 1\}.$$

By the transversality it is easy to see that the intersection of $\mathrm{im}(\psi)$ with $\Sigma_a(V_\bullet) \times \Sigma_b(W_\bullet) \subseteq \mathrm{Gr}_d(V) \times \mathrm{Gr}_{d+1}(V)$ is nonempty if and only if

$$\dim W^{(1)} = 1, \quad \dim W^{(i)} \cap V^{(d+2-i)} = 1, \quad i = 2, \dots, d + 1,$$

in which case

$$U_d = \bigoplus_{i=2}^{d+1} W^{(i)} \cap V^{(d+2-i)}, \quad U_{d+1} = U_d \oplus W^{(1)}.$$

Therefore

$$[\mathrm{im} \psi] \cdot (\sigma_a \times \sigma_b) = \begin{cases} 1, & b_1 = d, \quad b_i = d + 1 - a_{d+2-i}, i = 2, \dots, d + 1, \\ 0, & \text{otherwise.} \end{cases}$$

The desired result then follows from the fact that Schubert classes form dual basis under the intersection pairing. \square

Corollary 3.2. *The following identity holds in $\mathrm{Ch}^*(\mathrm{Gr}_{d+1}(V)_{\bar{k}})_{\mathbb{Q}}$:*

$$[\mathrm{DL}(V)] = \sum_c \sigma_c \sigma_{\hat{c}} q^{|\hat{c}|},$$

where the sum runs over $c = (c_1, \dots, c_d)$ such that $d \geq c_1 \geq \dots \geq c_d \geq 0$, and \hat{c} is the complement of c defined by $\hat{c} := (d - c_d, \dots, d - c_1)$.

Proof. By definition we have $[\mathrm{DL}(V)] = (\phi, \mathrm{id})^*[\mathrm{im}(\psi)]$. Since $U^\perp \simeq \sigma^*(V/U)^*$, by (3.0.3) it is easy to see that

$$(\phi, \mathrm{id})^*(\sigma_{a^*} \times \sigma_{b^*}) = q^{|a^*|} \sigma_{(a^*)'} \sigma_{b^*} \in \mathrm{Ch}^*(\mathrm{Gr}_{d+1}(V)_{\bar{k}})_{\mathbb{Q}}.$$

Since $b = (d, d + 1 - a_d, d + 1 - a_2, \dots, d + 1 - a_1)$, we know that $b^* = (a_1 - 1, \dots, a_d - 1, 0)$. Let $c = (a_1 - 1, \dots, a_d - 1)$. Then $\hat{c} = (d + 1 - a_d, \dots, d + 1 - a_1) = a^*$. It follows from Proposition 3.1 that

$$[\mathrm{DL}(V)] = \sum_c \sigma_{\hat{c}} \sigma_c q^{|\hat{c}|} = \sum_c \sigma_c \sigma_{\hat{c}} q^{|\hat{c}|}.$$

This completes the proof. \square

Example 3.3. • When $d = 1$, by Corollary 3.2 we obtain

$$[\mathrm{DL}(V)] = \sigma_1 + \sigma_1 q = (1 + q)\sigma_1.$$

This agrees with the fact that $\mathrm{DL}(V) \subseteq \mathrm{Gr}_{d+1}(V) \simeq \mathbb{P}^2$ is the Fermat curve of degree $1 + q$ (Example 2.9).

• When $d = 2$, by Corollary 3.2 we obtain

$$[\mathrm{DL}(V)] = \sigma_{2,2} + \sigma_1 \sigma_{2,1} q + \sigma_2 \sigma_{1,1} q^2 + \sigma_{1,1} \sigma_2 q^2 + \sigma_{2,1} \sigma_1 q^3 + \sigma_{2,2} q^4.$$

3.3. Proof of Theorem 1.3. By (2.12.2) and the projection formula, we have

$$\deg \mathrm{DL}(V) = [\mathrm{DL}(V)] \cdot c_1(\mathrm{Pl}^* \mathcal{O}_{\mathbb{P}(\wedge^{d+1} V)}(1))^d.$$

The result then follows from Corollary 3.2 and $\sigma_1 = c_1(\mathrm{Pl}^* \mathcal{O}_{\mathbb{P}(\wedge^{d+1} V)}(1))$ is the hyperplane place class under the Plücker embedding.

3.4. Proof of Corollary 1.4.

(i) This follows immediately from Theorems 1.1 and 1.3.

(ii) Let

$$S_c(x_1, \dots, x_d) = \det(x_i^{c_j+d-j}) / \det(x_i^{d-j})$$

($1 \leq i, j \leq d$) be the *Schur polynomial* associated to $c = (c_1, \dots, c_d)$ ([Mac92]). It is a symmetric polynomial of degree $|c|$. Write

$$S := \sum_c S_c(qx_1, \dots, qx_d) S_{\mathcal{C}'}(x_1, \dots, x_d) S_1^d(x_1, \dots, x_d) =: \sum_\lambda \kappa_\lambda S_\lambda,$$

where λ runs over $\lambda = (\lambda_1, \dots, \lambda_d)$ with $|\lambda| = d(d+1)$. By definition, the coefficient κ_λ is given by the coefficient of $x_1^{\lambda_1+d-1} x_2^{\lambda_2+d-2} \dots x_d^{\lambda_d}$ in

$$(3.3.1) \quad S \cdot \det(x_i^{d-j}) = S \cdot \prod_{i < j} (x_i - x_j).$$

Since the class of a point is $\sigma_{(d+1, \dots, d+1)}$, we know that

$$\sum_c \sigma_c \sigma_{\mathcal{C}'} \sigma_1^d q^{|c|} = \kappa_{(d+1, d+1, \dots, d+1)},$$

i.e., the coefficient of $x_1^{2d} x_2^{2d-1} \dots x_d^{d+1}$ in (3.3.1). It remains to compute (3.3.1). By the dual Cauchy identity for Schur polynomials ([Mac92, 0.11']), we have

$$\sum_c S_c(x_1, \dots, x_d) S_{\mathcal{C}'}(y_1, \dots, y_d) = \prod_{i,j=1}^d (x_i + y_j).$$

By definition, we have

$$S_1(x_1, \dots, x_d) = x_1 + \dots + x_d.$$

Therefore

$$S = \sum_c S_c(qx_1, \dots, qx_d) S_{\mathcal{C}'}(x_1, \dots, x_d) S_1^d(x_1, \dots, x_d) = \left(\prod_{i,j=1}^d (qx_i + x_j) \right) (x_1 + \dots + x_d)^d,$$

and thus

$$(3.3.2) \quad S \cdot \prod_{i < j} (x_i - x_j) = \left(\prod_{i,j=1}^d (qx_i + x_j) \right) (x_1 + \dots + x_d)^d \left(\prod_{i < j} (x_i - x_j) \right).$$

The result then follows from Item (i).

(iii) By applying Pieri's formula ([EH16, Proposition 4.9]) d times, we know that the term $\sigma_c \sigma_{\mathcal{C}'} \sigma_1^d$ is equal to the number of sequences of Young diagrams $c^{(0)}, c^{(1)}, \dots, c^{(d)}$ starting with $c^{(0)} = c$ and ending with $c^{(d)} = (\mathcal{C}')^*$ such that each $c^{(i+1)}$ has exactly one more box than $c^{(i)}$. Equivalently, it is the number of standard Young tableaux of skew shape $(\mathcal{C}')^*/c$. Now Item (i) shows that the number of such standard Young tableaux with $|c| = l$ is equal the coefficient of q^l in

$$\prod_{i=1}^d \frac{1 - q^{2i}}{1 - q} = \prod_{i=1}^d \sum_{j=0}^{2i-1} q^j,$$

which is equal to the number of ordered partitions (l_1, \dots, l_d) of l satisfying the extra conditions

$$0 \leq l_i \leq 2i - 1, \quad i = 1, \dots, d.$$

Example 3.4. We end with an example illustrating Corollary 1.4 (ii) (iii).

- When $d = 1$, the polynomial (3.3.2) is equal to $(1 + q)x_1^2$. The coefficient of x_1^2 is given by $1 + q = \frac{1-q^2}{1-q}$ as in Corollary 1.4 (ii).
- When $d = 2$, the polynomial (3.3.2) is equal to

$$(q + 2q^2 + q^3) x_1^6 x_2 + (1 + 3q + 4q^2 + 3q^3 + q^4) x_1^5 x_2^2 + (1 + 2q + 2q^2 + 2q^3 + q^4) x_1^4 x_2^3 + (-1 - 2q - 2q^2 - 2q^3 - q^4) x_1^3 x_2^4 + (-1 - 3q - 4q^2 - 3q^3 - q^4) x_1^2 x_2^5 + (-q - 2q^2 - q^3) x_1 x_2^6$$

The coefficient of $x_1^4 x_2^3$ is given by

$$1 + 2q + 2q^2 + 2q^3 + q^4 = (1 + q)(1 + q + q^2 + q^3),$$

which equals $\frac{1-q^2}{1-q} \cdot \frac{1-q^4}{1-q}$ as in Corollary 1.4 (ii).

- When $d = 3$, the coefficient of $x_1^6 x_2^5 x_3^4$ is equal to

$$1 + 3q + 5q^2 + 7q^3 + 8q^4 + 8q^5 + 7q^6 + 5q^7 + 3q^8 + q^9 = (1 + q)(1 + q^2 + q^3)(1 + q + q^2 + q^3 + q^4 + q^5),$$

as in Corollary 1.4 (ii). In Table 1 we list all standard Young tableaux of skew shape $(\hat{\mathcal{C}})^*/c$ with $|c| = 4$ as in Corollary 1.4 (iii). Notice the total number of such Young tableaux is 8, which indeed agrees with the coefficient of q^4 .

c	$(\hat{\mathcal{C}})^*$	$(\hat{\mathcal{C}})^*/c$

TABLE 1. standard Young tableaux of skew shape $(\hat{\mathcal{C}})^*/c$

REFERENCES

- [EH16] David Eisenbud and Joe Harris. *3264 and all that—a second course in algebraic geometry*. Cambridge University Press, Cambridge, 2016. 10, 11, 13

- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998. 3, 8, 10
- [HLZ19] Xuhua He, Chao Li, and Yihang Zhu. Fine Deligne-Lusztig varieties and arithmetic fundamental lemmas. *Forum Math. Sigma*, 7:e47, 55, 2019. 2
- [HP20] Thomas Hudson and Dennis Peters. On the K -theoretic fundamental class of Deligne-Lusztig varieties. *J. Pure Appl. Algebra*, 224(8):106335, 8, 2020. 2
- [Kim20] Dongkwan Kim. Homology class of a Deligne-Lusztig variety and its analogs. *Int. Math. Res. Not. IMRN*, (4):1246–1280, 2020. 2
- [KR11] Stephen S. Kudla and Michael Rapoport. Special cycles on unitary Shimura varieties I. Unramified local theory. *Invent. Math.*, 184(3):629–682, 2011. 1
- [KR14] Stephen S. Kudla and Michael Rapoport. Special cycles on unitary Shimura varieties II: Global theory. *J. Reine Angew. Math.*, 697:91–157, 2014. 1
- [Kud04] Stephen S. Kudla. Special cycles and derivatives of Eisenstein series. In *Heegner points and Rankin L -series*, volume 49 of *Math. Sci. Res. Inst. Publ.*, pages 243–270. Cambridge Univ. Press, Cambridge, 2004. 2
- [LTX⁺19] Yifeng Liu, Yichao Tian, Liang Xiao, Wei Zhang, and Xinwen Zhu. On the Beilinson-Bloch-Kato conjecture for Rankin-Selberg motives. *arXiv e-prints*, page arXiv:1912.11942, December 2019. 5, 6
- [LZ] Chao Li and Wei Zhang. Kudla-Rapoport cycles and derivatives of local densities. *J. Amer. Math. Soc.* to appear. 2, 5, 8, 9
- [LZ21] Chao Li and Wei Zhang. On the arithmetic Siegel-Weil formula for GSpin Shimura varieties. *arXiv e-prints*, page arXiv:2106.15038, June 2021. 2
- [Mac92] I. G. Macdonald. Schur functions: theme and variations. In *Séminaire Lotharingien de Combinatoire (Saint-Nabor, 1992)*, volume 498 of *Publ. Inst. Rech. Math. Av.*, pages 5–39. Univ. Louis Pasteur, Strasbourg, 1992. 13
- [Tat65] John T. Tate. Algebraic cycles and poles of zeta functions. In *Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963)*, pages 93–110. Harper & Row, New York, 1965. 3
- [Tat94] John Tate. Conjectures on algebraic cycles in l -adic cohomology. In *Motives (Seattle, WA, 1991)*, volume 55 of *Proc. Sympos. Pure Math.*, pages 71–83. Amer. Math. Soc., Providence, RI, 1994. 3
- [Vol10] Inken Vollaard. The supersingular locus of the Shimura variety for $\mathrm{GU}(1, s)$. *Canad. J. Math.*, 62(3):668–720, 2010. 5, 7
- [VW11] Inken Vollaard and Torsten Wedhorn. The supersingular locus of the Shimura variety of $\mathrm{GU}(1, n - 1)$ II. *Invent. Math.*, 184(3):591–627, 2011. 1, 5

COLUMBIA UNIVERSITY, DEPARTMENT OF MATHEMATICS, 2990 BROADWAY, NEW YORK, NY 10027, USA
E-mail address: chaoli@math.columbia.edu