

Calculus I
Section 005
Evan Fink
3/30/08

Practice Midterm II

1. Find the derivatives of the following functions. You do not have to simplify your answers.

a. $f(x) = 3x^2$

b. $g(x) = \tan x + \cos x + \ln(x^2 - 1)$

c. $q(t) = \frac{1-e^t}{\sqrt[3]{1+t}}$

d. $h(x) = \frac{(x+2)^{20} \sqrt[5]{x-1}}{(x-49)(x-26)}$ (Hint: this can be done directly, but there's an easier way involving logarithms.)

2. Consider the equation $x^2y + xy^2 = 3x$.

a. Using implicit differentiation, compute $\frac{dy}{dx}$ in terms of x and y .

b. Find the equation of the tangent line to the curve given by this equation at the point $(2,1)$.

3. After an oil spill, a circular oil slick grows in such a way that its area increases at the rate of $15 \frac{\text{mi}^2}{\text{hr}}$. When the area of the slick is $225\pi \text{ mi}^2$, at what rate is the radius of the slick increasing? (Remember that the area of a circle is given by $A = \pi r^2$.)

4. A bug crawls along a yard stick, starting at time $x = 0$ seconds. Its position at time x seconds is y inches, where $y = \sqrt{x+1} - \frac{1}{4}x + 8$.

a. In $\frac{\text{in}^2}{\text{sec}}$, what is the **acceleration** of the bug after 2 seconds?

b. At what time is the bug at rest?

c. Using a linear approximation (or differentials), estimate the bug's position at 0.08 seconds to 2 decimal places.

5. Let $f(x) = xe^{x-1}$, $-4 \leq x \leq 4$.

a. On what intervals is $f(x)$ increasing? Decreasing?

b. On what intervals is $f(x)$ concave up? Down?

c. Find the absolute maximum and minimum values of $f(x)$, and the x values where they occur. Hint: there's a slightly subtle comparison of two numbers. To deal with this, it might help either to look at the first derivative, or to remember that e is greater than 2 – I recommend the first one.

6. a. State the Mean Value Theorem.

b. Let $f(x)$ be a function that is continuous on the interval $[1,5]$, differentiable on the interval $(1,5)$, and satisfies

$$f'(x) = 5 + \sin(g(x))$$

for some function $g(x)$. If $f(1) = 10$, use the Mean Value Theorem to explain why it is impossible for $f(5)$ to equal 50. (Hint: what is the greatest possible number that $f'(x)$ could ever be? What is the least possible number of $f'(x)$? Even though you don't know what $g(x)$ is, you should be able to answer these if you know basic facts about the sine function.)