

# TYPE A LEVEL ONE CONFORMAL BLOCKS DIVISORS REVISITED

MAKSYM FEDORCHUK

ABSTRACT. We prove that all type A level one conformal blocks divisors on  $\overline{M}_{0,n}$  are effective sums of boundary divisors. This leads to an elementary proof of their nefness.

Denote by  $\langle a \rangle_m$  the representative in  $\{0, 1, \dots, m-1\}$  of the residue of  $a$  modulo  $m$ .

**Definition 1.1.** Consider  $n$  integer numbers  $(d_1, \dots, d_n)$  and let  $m$  be an integer dividing  $\sum_{i=1}^n d_i$ . We define a divisor on  $\overline{M}_{0,n}$  by the following formula

$$(1) \quad D((d_1, \dots, d_n), m) = \sum_{i=1}^n \langle d_i \rangle_m \langle m - d_i \rangle_m \psi_i - \sum_{I,J} \langle d(I) \rangle_m \langle d(J) \rangle_m \Delta_{I,J},$$

where  $d(I) = \sum_{i \in I} d_i$  for any  $I \subset \{1, \dots, n\}$ .

The motivation for this definition comes from the following result [Fed11, Proposition 4.8]:

**Proposition 1.2.** For a weight vector  $\vec{d} = (d_1, \dots, d_n)$ , let  $m$  be an integer dividing  $\sum_{i=1}^n d_i$ . Let  $\mathbb{E}$  be the pullback to  $\overline{M}_{0,n}$  of the Hodge bundle over  $\overline{M}_g$  via the weighted cyclic  $m$ -covering morphism  $f_{\vec{d},m}$  and let  $\mathbb{E}_j$  be the eigenbundle of  $\mathbb{E}$  associated to the character  $j$  of  $\mu_m$ . Then

$$\lambda_{\vec{d},m}(j) := \det \mathbb{E}_j = \frac{1}{2m^2} \left[ \sum_{i=1}^n \langle jd_i \rangle_m \langle m - jd_i \rangle_m \psi_i - \sum_{I,J} \langle jd(I) \rangle_m \langle jd(J) \rangle_m \Delta_{I,J} \right].$$

**Remark 1.3.** To obtain a closed form formula for the  $\mathfrak{sl}_m$  level 1 conformal blocks divisor  $\mathbb{D}(\mathfrak{sl}_m, 1, (d_1, \dots, d_n))$ , set  $j = 1$  and multiply the class of  $\lambda_{\vec{d},m}(1)$  by  $m$ .

The divisor  $D((d_1, \dots, d_n), m)$  has at least three incarnations:

- (1) It is a determinant of a Hodge eigenbundle [Fed11].
- (2) It is a type A level one conformal blocks divisor [Fak09, AGSS10].
- (3) It is a pullback of a natural polarization on a GIT quotient of a parameter space of  $n$ -pointed rational normal curves [Gia10, GG11].

Each interpretation leads to an independent proof of nefness of  $D = D((d_1, \dots, d_n), m)$ : the first via the semipositivity of the Hodge bundle over  $\overline{M}_g$ , which comes from Hodge theory [Kol90]; the second via the theory of conformal blocks which realizes  $D$  as a quotient of a trivial vector bundle over  $\overline{M}_{0,n}$  [Fak09]; the third via GIT.

---

Date: November 1, 2011.

We now propose a fourth proof of nefness of  $D((d_1, \dots, d_n), m)$  which is independent of all of the above and is completely elementary.

**Proposition 1.4.**  $D((d_1, \dots, d_n), m)$  is a sum of effective boundary divisors on  $\overline{M}_{0,n}$ .

*Proof of Proposition 1.4.* We use the following elementary relation in  $\text{Pic}(\overline{M}_{0,n})$ :

$$(2) \quad \psi_i + \psi_j = \sum_{i \in I, j \in J} \Delta_{I,J}.$$

The claim of the proposition will follow from combining a number of Relations (2). To specify which relations we use, we introduce some notation. By replacing each  $d_i$  by  $\langle d_i \rangle_m$  we can assume that  $d_i \leq m - 1$ . Let  $d_1 + \dots + d_n = ms$ . First, we let  $S = \{p_1, \dots, p_{ms}\}$  be a collection of indices where each index  $i \in \{1, \dots, n\}$  appears  $d_i$  times. We subdivide  $S$  into  $m$  subsets  $S_1, \dots, S_m$  of cardinality  $s$  with the property that each  $S_k$  contains precisely one occurrence of each  $i \in \{1, \dots, n\}$ . We note that this can be done in many ways. One way is to arrange elements of  $S$  at the vertices of a regular  $ms$ -gon so that  $d_i$  occurrences of each  $i$  are adjacent and to take  $S_k$ 's to be regular  $s$ -gons formed by the chords divisible by  $m$ . Since  $d_i \leq m - 1$ , this subdivision satisfies all desired properties. We also denote by  $\Gamma(S)$  the set of all unordered pairs  $(p_i, p_j) \in S$  such that  $p_i \neq p_j$  as elements of  $\{1, \dots, n\}$ . We define  $\Gamma(S_k)$  for each  $k = 1, \dots, m$  in a similar fashion. Note that by our construction,  $\Gamma(S_k)$  is simply the set of all unordered pairs of distinct elements of  $S_k$ .

Consider now the following sum

$$(3) \quad \Sigma = \sum_{(p_i, p_j) \in \Gamma(S)} (\psi_{p_i} + \psi_{p_j}) - m \sum_{k=1}^m \sum_{(p_i, p_j) \in \Gamma(S_k)} (\psi_{p_i} + \psi_{p_j})$$

**Claim 1.5.**

$$\Sigma = \sum_{i=1}^n \langle d_i \rangle_m \langle m - d_i \rangle_m \psi_i$$

*Proof of Claim.* We need to count the number of occurrences in  $\Sigma$  of each  $\psi_i$ ,  $i = 1, \dots, n$ . First, in the first sum of (3), each  $\psi_i$  occurs  $d_i(ms - d_i)$  times. In the second sum, each  $\psi_i$  occurs  $d_i(s - 1)$  times. The claim follows.  $\square$

By rewriting each occurrence of  $\psi_i + \psi_j$  in Equation (3) using Relation (2), we see that  $\Sigma$  is an effective combination of the boundary divisors. Namely, we have

$$(4) \quad \Sigma = \sum_{(p_i, p_j) \in \Gamma(S)} (\psi_{p_i} + \psi_{p_j}) - m \sum_{k=1}^m \sum_{(p_i, p_j) \in \Gamma(S_k)} (\psi_{p_i} + \psi_{p_j})$$

$$(5) \quad = \sum_{(p_i, p_j) \in \Gamma(S)} \left( \sum_{p_i \in I, p_j \in J} \Delta_{I,J} \right) - m \sum_{k=1}^m \sum_{(p_i, p_j) \in \Gamma(S_k)} \left( \sum_{p_i \in I, p_j \in J} \Delta_{I,J} \right)$$

To prove the proposition it remains to show that each boundary divisor  $\Delta_{I,J}$  occurs with coefficient at least  $\langle d(I) \rangle_m \langle d(J) \rangle_m$  in (5). Recall that  $d(I) = \sum_{i \in I} d_i$ . Write  $d(I) = mq + r$ . Let  $x_1, \dots, x_m$  be the number of indices from  $I$  occurring in each of the

sets  $S_1, \dots, S_m$ . Then  $x_1 + \dots + x_m = d(I) = mq + r$ . Tracing through the construction we see that the coefficient with which  $\Delta_{I,J}$  occurs in  $\Sigma$  is

$$(6) \quad d(I)(ms - d(I)) - \sum_{k=1}^m x_k(s - x_k).$$

Since  $x(s - x)$  is a concave function, the minimum in (6) is achieved when all  $x_i$  differ by at most 1 from each other, i.e., when there are  $r$   $x_i$ 's equal to  $q + 1$  and  $m - r$  equal to  $q$ . A straightforward computation now shows that for these  $x_i$ 's Equation (6) evaluates to

$$r(m - r) = \langle d(I) \rangle_m \langle d(J) \rangle_m.$$

This finishes the proof.  $\square$

**Corollary 1.6.**  $D((d_1, \dots, d_n), m)$  is a nef divisor on  $\overline{M}_{0,n}$ .

*Proof.* Since  $D = D((d_1, \dots, d_n), m)$  is an effective combination of boundary divisors, it intersects any irreducible curve meeting the interior  $M_{0,n} \subset \overline{M}_{0,n}$  non-negatively. Next, we use the fact that  $D((d_1, \dots, d_n), m)$  is functorial with respect to the boundary stratification (i.e., it restricts to a tensor product (of pullbacks) of divisors of the same form on each boundary) to conclude inductively that  $D((d_1, \dots, d_n), m)$  intersects all curves non-negatively.  $\square$

We would like to end with the following conjecture

**Conjecture 1.7.** Suppose  $m \geq 3$  is prime and  $m \mid \sum_{i=1}^n d_i$ . Then the divisor class

$$(7) \quad f_{\vec{d},m}^*(2m^2 \lambda_{\vec{d},m}(j) - \delta_{\text{irr}}) \\ = \sum_{i=1}^n \langle jd_i \rangle_m \langle m - jd_i \rangle_m \psi_i - \sum_{I,J} \langle jd(I) \rangle_m \langle jd(J) \rangle_m \Delta_{I,J} - m \sum_{I: m \mid d(I)} \Delta_{I,J}$$

is nef on  $\overline{M}_{0,n}$  and generates an extremal ray of  $\text{Nef}(\overline{M}_{0,n}/S_n)$ .

**Example 1.8.** By taking  $n = 9$ ,  $m = 3$ , and  $j = 1$ , we obtain the extremal ray  $\Delta_2 + \Delta_3 + 2\Delta_4$  of  $\text{Nef}(\overline{M}_{0,9}/S_9)$  not accounted for by conformal blocks divisors.

We proved the conjecture for  $m = 3, 5$  (see [Fed11]) but for  $m \geq 7$  the question seems to be open.

## REFERENCES

- [AGSS10] Maxim Arap, Angela Gibney, James Stankewicz, and David Swinarski.  $sl_n$  level 1 conformal blocks divisors on  $\overline{M}_{0,n}$ , 2010. [arXiv:1009.4664](#) [math.AG].
- [Fak09] Najmuddin Fakhruddin. Chern classes of conformal blocks, 2009. [arXiv:0904.2918v3](#) [math.AG].
- [Fed11] Maksym Fedorchuk. Cyclic covering morphisms on  $\overline{M}_{0,n}$ , 2011. [arXiv:1105.0655](#) [math.AG].
- [GG11] Noah Giansiracusa and Angela Gibney. The cone of type A, level one conformal blocks divisors, 2011. [arXiv:1105.3139](#) [math.AG].
- [Gia10] Noah Giansiracusa. Conformal blocks and rational normal curves, 2010. [arXiv:1012.4835](#) [math.AG].
- [Kol90] János Kollár. Projectivity of complete moduli. *J. Differential Geom.*, 32(1):235–268, 1990.