

TOWARDS A DYNAMICAL MANIN-MUMFORD CONJECTURE

DRAGOS GHIOCA, THOMAS J. TUCKER, AND SHOUWU ZHANG

ABSTRACT. We provide a family of counterexamples to a first formulation of the dynamical Manin-Mumford conjecture. We propose a revision of this conjecture and prove it for endomorphisms of abelian varieties and for endomorphisms of $\mathbb{P}^1 \times \mathbb{P}^1$.

INTRODUCTION

The Manin-Mumford conjecture, proved by Raynaud [Ray83a, Ray83b], states that if V is a subvariety of an abelian variety A , defined over \mathbb{C} , then V contains a dense set of torsion points if and only if V is a torsion translate of an abelian subvariety of A . Over number fields, a stronger theorem, conjectured by Bogomolov and proved by Ullmo [Ull98] and Zhang [Zha98], states that V contains a dense set of points with Néron-Tate height tending to zero if and only if V is a torsion translate of an abelian subvariety of A .

The proofs of [Ull98] and [Zha98] make important use of an equidistribution theorem for points of small canonical height on abelian varieties (see [SUZ97]). Recently, dynamical analogues of this equidistribution theorem have been proved by various authors [BR06, CL06, FRL06, Yua08]. Thus, it seems quite natural to attempt to find dynamical versions of the Bogomolov and Manin-Mumford conjectures.

Here, we will present a family of counterexamples to a first formulation of the dynamical Manin-Mumford and Bogomolov conjectures (see [Zha95, Conjecture 2.5] and [Zha06, Conjecture 1.2.1, Conjecture 4.1.7]). The family is fairly simple. It involves the diagonal subgroup of $E \times E$ under the action of $([\omega_1], [\omega_2])$ for $[\omega_1], [\omega_2] \in \text{End}(E)$ elements of equal degree, where E is an elliptic curve with complex multiplication. This is explained in Theorem 1.2. We then propose a revision of this conjecture, which we show that it holds in the cases of group endomorphisms of abelian varieties and of lines under the action of arbitrary endomorphisms of $(\mathbb{P}^1)^2$.

Acknowledgements. We would like to thank Pascal Autissier, Philipp Habegger, Fabien Pazuki, Thomas Scanlon, Lucien Szpiro, Xinyi Yuan, and Michael Zieve for many helpful conversations. We would also like to thank the The

The first author was partially supported by NSERC. The second author was partially supported by NSF Grants 0801072 and 0854839.

Centro di Ricerca Matematica Ennio De Giorgi and the University of Bordeaux 1 for their hospitality.

1. COUNTEREXAMPLES AND REVISION

Our examples are on algebraic varieties, whereas [Zha06, Conjecture 1.2.1] is a conjecture about more general Kähler varieties; on the other hand, [Zha06, Conjecture 4.1.7] is a statement about general points of small canonical height, while our counterexample is for the more specific case of preperiodic points (which all have canonical height equal to 0). Thus, for simplicity, we will state an “algebraic dynamical Manin-Mumford” conjecture, which is implied by both (but is weaker than either of) [Zha06, Conjecture 1.2.1] and [Zha06, Conjecture 4.1.7]. We will then provide a family of counterexamples in Theorem 1.2.

Before stating the conjecture, we define a few terms:

- An endomorphism $\varphi : X \rightarrow X$ of a projective variety is said to have a *polarization* if there exists an ample line bundle \mathcal{L} on X such that $\varphi^*\mathcal{L} = \mathcal{L}^{\otimes d}$ for some $d > 1$.
- A subvariety Y of a projective variety X with endomorphism φ is said to be *preperiodic* (under the action of φ) if there exist positive integers m and k such that $\varphi^{m+k}(Y) = \varphi^m(Y)$.
- $\text{Prep}_\varphi = \text{Prep}_\varphi(X)$ is defined to be the set of all preperiodic points of X (under the action of φ).

Conjecture 1.1. (*Algebraic Dynamical Manin-Mumford*) *Let $\varphi : X \rightarrow X$ be an endomorphism of a projective variety defined over the complex numbers with a polarization, and let Y be a subvariety of X . Then Y is preperiodic if and only if $Y \cap \text{Prep}_\varphi(X)$ is Zariski dense in Y .*

The following theorem shows that Conjecture 1.1 is often false for $E \times E$ when E has complex multiplication.

Theorem 1.2. *Let E be an elliptic curve with complex multiplication defined over the complex numbers. Let R be an order in an imaginary quadratic extension of \mathbb{Q} such that there is an isomorphism $\iota : R \rightarrow \text{End}(E)$ written as $\iota(\omega) = [\omega]$. Suppose that $\omega_1, \omega_2 \in R$ such that $|\omega_1| = |\omega_2| > 1$. Then the diagonal of $E \times E$ is preperiodic under the action of $([\omega_1], [\omega_2])$ if and only if ω_1/ω_2 is a root of unity.*

Remark 1.3. In Theorem 1.2, note that all torsion points of E are preperiodic under both $[\omega_1]$ and $[\omega_2]$ (see also Claim 2.2); therefore the diagonal subvariety of $E \times E$ has a Zariski dense set of preperiodic points under the map $([\omega_1], [\omega_2])$. Hence Theorem 1.2 gives rise to many counterexamples. For example, take any ω_1 such that $|\omega_1| > 1$ is the square-root of an odd integer and $\omega_1/\bar{\omega}_1$ is not a root of unity, and let $\omega_2 = \bar{\omega}_1$. There are also examples in which ω_1 is an integer; take for example $\omega_1 = 5$ and $\omega_2 = 3 + 4i$ in $\mathbb{Z}[i]$. Note that for each $\omega \in R$, we have $|\omega|^2 = \#\ker([\omega]) = \deg([\omega])$.

The lowest degree counterexamples come from multiplication by $\frac{1+\sqrt{-7}}{2}$ on the curve $y^2 = x^3 - 35x + 98$ (see [Sil94, II.2], where this is worked out in detail). Pazuki [Paz] has worked out a number of higher-dimensional counterexamples.

We propose the following reformulation of Conjecture 1.1.

Conjecture 1.4. *Let X be a projective variety, let $\varphi : X \rightarrow X$ be an endomorphism defined over \mathbb{C} with a polarization, and let Y be a subvariety of X which has no component included into the singular part of X . Then Y is preperiodic under φ if and only if there exists a Zariski dense subset of smooth points $x \in Y \cap \text{Prep}_\varphi(X)$ such that the tangent subspace of Y at x is preperiodic under the induced action of φ on the Grassmanian $\text{Gr}_{\dim(Y)}(T_{X,x})$. (Here we denote by $T_{X,x}$ the tangent space of X at the point x .)*

One direction is clear: if Y is preperiodic then it will lift to a preperiodic subvariety of the Grassmanian, which will contain a dense set of preperiodic points, by work of Fakhruddin [Fak03]. The other direction seems more difficult. In the following we show that Conjecture 1.4 is true in the cases of group endomorphisms of abelian varieties and of arbitrary endomorphisms of $X = \mathbb{P}^1 \times \mathbb{P}^1$ when Y is a line.

2. ABELIAN VARIETIES

Theorem 2.1. *Conjecture 1.4 holds for group endomorphisms φ of abelian varieties X .*

Proof. Clearly, it suffices to prove our result assuming Y is irreducible. We start with an easy claim.

Claim 2.2. *For a polarizable algebraic group endomorphism φ of an abelian variety X , we have $\text{Prep}_\varphi(X) = X_{\text{tor}}$.*

Proof of Claim 2.2. Clearly, each torsion point of X is preperiodic under φ since φ is an algebraic group endomorphism, and so, for each $m \in \mathbb{N}$, we have $\varphi(X[m]) \subset X[m]$. Since $X[m]$ is a finite set, we conclude that each torsion point in $X[m]$ is preperiodic under φ .

Conversely, if $\alpha \in \text{Prep}_\varphi(X)$, then there are $k > 0$ and $m \geq 0$ such that $\varphi^m(\alpha) = \varphi^{m+k}(\alpha)$ or $(\varphi^{k+m} - \varphi^m)(\alpha) = 0$, or α is in the kernel of $\varphi^{k+m} - \varphi^m$. Since φ is polarized, then φ acts on $\text{Lie}(X)$ with eigenvalues whose absolute values are $|\deg \varphi|^{1/2 \dim X}$. Thus $(\varphi^{k+m} - \varphi^m)$ has all eigenvalues nonvanishing on $\text{Lie}(X)$. It follows that $\varphi^{k+m} - \varphi^m$ is finite over X and then has finite kernel. Thus α is torsion. \square

Now, assume $Y \subset X$ contains a Zariski dense set of points from $\text{Prep}_\varphi(X)$. By Claim 2.2, it follows that Y contains a Zariski dense set of torsion points of X . Hence, the classical Manin-Mumford conjecture (proved by Raynaud [Ray83b]) implies that Y is a torsion translate of an algebraic subgroup of

X , i.e. $Y = \gamma + H$, where $\gamma \in X_{\text{tor}}$ and H is an abelian subvariety of X . Since $\gamma \in \text{Prep}_\varphi(X)$ (also according to Claim 2.2), we are left to show that H is preperiodic under φ .

Now, using the exponential uniformization map on X , we see that the preperiodicity of H under the endomorphism φ is equivalent with the preperiodicity of the tangent subspace of H at *any* point $x \in H$ under the induced action of φ on $\text{Gr}_{\dim(Y)}(T_{X,x})$. This concludes our proof. \square

Proof of Theorem 1.2. The action of $[\omega_1, \omega_2]$ on

$$\text{Lie}(E \times E) = \text{Lie}(E) \oplus \text{Lie}(E)$$

is given as multiplication by (ω_1, ω_2) . The diagonal in $\text{Lie}(E) \oplus \text{Lie}(E)$ is preperiodic under (ω_1, ω_2) if and only if ω_1/ω_2 is a root of unity. \square

3. LINES IN $\mathbb{P}^1 \times \mathbb{P}^1$

Theorem 3.1. *Let $f, g : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be nonconstant rational maps over \mathbb{C} such that $\deg f = \deg g > 1$, let*

$$(f, g) : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$$

be the map given by $(f, g)(a, b) = (f(a), g(b))$, and let $L \subset \mathbb{P}^1 \times \mathbb{P}^1$ be a line. If there exists an infinite subset of points $x \in \text{Prep}_{(f,g)} \cap L$ such that the tangent subspace of L at x is preperiodic under the induced action of φ on $\text{Gr}_1(T_{\mathbb{P}^1 \times \mathbb{P}^1, x})$, then L is preperiodic under (f, g) .

The proof is somewhat similar to the proof of the main result from [GT10] which dealt with the case of polynomial maps.

Step 1: reduction to the case L is the diagonal subvariety. If L does not project dominantly to both copies of \mathbb{P}^1 , then (without loss of generality) we may assume $L = \{a\} \times \mathbb{P}^1$. Since L contains infinitely many preperiodic points for the action of (f, g) on $\mathbb{P}^1 \times \mathbb{P}^1$, we conclude that a is preperiodic for f . Therefore, L is indeed preperiodic under the action of (f, g) (no extra hypothesis is necessary in this case).

So, from now on, we assume L projects dominantly on $\mathbb{P}^1 \times \mathbb{P}^1$. Thus, there exists a linear map σ such that the equation of L is $y = \sigma(x)$, where (x, y) is a generic pair of projective coordinates for $\mathbb{P}^1 \times \mathbb{P}^1$. Then, replacing g by $\sigma^{-1} \circ g \circ \sigma$ yields that the diagonal line Δ contains infinitely many preperiodic points for $(f, \sigma^{-1}g\sigma)$. Therefore, from now on, we assume $L = \Delta$ is the diagonal subvariety of $\mathbb{P}^1 \times \mathbb{P}^1$. \square

Step 2: using theorems of equidistribution in dynamics. First we note that $(x, x) \in \Delta$ is in $\text{Prep}_{(f,g)}$ if and only if $x \in \text{Prep}_f \cap \text{Prep}_g$. Thus, $\text{Prep}_{(f,g)} \cap \Delta$ is infinite if and only if $\text{Prep}_f \cap \text{Prep}_g$ is infinite. By recent work of Baker–DeMarco [BD] and Yuan–Zhang [YZ], we have $\text{Prep}_f = \text{Prep}_g$. Denote the equilibrium measure and the Julia set of a map h of degree greater than 1 as μ_h and \mathcal{J}_h . Then, recent work of Yuan–Zhang [YZ], Theorem 1.6, yields that $\mu_f = \mu_g$ and $\mathcal{J}_f = \mathcal{J}_g$. \square

Step 3: applying a theorem of Levin. Thus, f and g satisfy the conditions (i)-(ii) of Theorem 3.2 proved in [Lev90].

Theorem 3.2. (*Levin*) *Let $f, g \in \mathbb{C}(t)$ be rational maps of degree greater than 1 satisfying the following two conditions:*

- (i) $\mathcal{J}_f = \mathcal{J}_g$ and $\mu_f = \mu_g$.
- (ii) *there exists a repelling periodic point for g which is also preperiodic for f .*

Then one of the following conclusions hold:

- (1) *f and g share a common parabolic orbifold.*
- (2) *there exist $k, \ell \in \mathbb{N}$ such that $f^\ell \circ g^k = f^{2\ell}$.*

Assume first that (2) holds. Using the fact that $\deg(f) = \deg(g) \geq 2$, we conclude that $k = \ell$ in conclusion (2) of Theorem 3.2. Hence $f^\ell \circ g^\ell = f^{2\ell}$. We claim that this relation yields that the diagonal $\Delta \in \mathbb{P}^1 \times \mathbb{P}^1$ is (f, g) -preperiodic.

Indeed, because $g^\ell : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is surjective, for each $x \in \mathbb{P}^1(\mathbb{C})$, there exists $y \in \mathbb{P}^1(\mathbb{C})$ such that $x = g^\ell(y)$. Furthermore, using that $f^\ell \circ g^\ell = f^{2\ell}$, we obtain that $f^{2\ell}(y) = f^\ell(g^\ell(y))$, and so, $(f^{2\ell}(y), g^{2\ell}(y)) = (f^\ell(x), g^\ell(x))$. Therefore $(f^\ell, g^\ell)(\Delta) \subset (f^{2\ell}, g^{2\ell})(\Delta)$. Moreover, since Δ is irreducible, we conclude that $(f^{2\ell}, g^{2\ell})(\Delta) = (f^\ell, g^\ell)(\Delta)$, and thus Δ is (f, g) -preperiodic, as conjectured. Note that so far we did not need the extra assumption about the preperiodicity of the tangent subspaces. \square

Step 4: the case when f and g share a common parabolic orbifold. Now, assume that (1) holds in Theorem 3.2. Douady-Hubbard [DH93] (following Thurston) show that when f has a parabolic orbifold, it means that modulo conjugation, $f(t)$ is either

- (a) t^m for some $m \in \mathbb{Z}$; or
- (b) T_m for some $m \in \mathbb{N}$ (where T_m denotes the m -th Chebychev polynomial); or
- (c) a Lattès map.

Douady-Hubbard-Thurston's result is proved in a general topological setting, for branched coverings of the Riemann sphere. It can be proved much more simply for rational functions, using Galois theory (see [GZ]). For example, consider the parabolic orbifold that arises when f is a degree four map which maps six points with ramification index two onto three points in its image. Taking a genus 1 cover of \mathbb{P}^1 that ramifies at exactly these three points plus one more point then gives rise to the multiplication-by-2 map on an elliptic curve that descends to f under the hyperelliptic involution (this can be checked easily using Abhyankar's lemma for ramification indices under composita).

Now, in the case where one of the maps is conjugate to a powering map or a Chebychev polynomial (cases (a) and (b) above), the other cannot be Lattès map since the two maps must have the same Julia set; the Julia

set of a Lattès map is the entire Riemann sphere, which can never be the case for a polynomial or for a rational map t^m for $m \in \mathbb{Z}$. Similarly, if one map is conjugate to t^m , then the other one cannot be conjugate to T_m (and vice-versa) since the Julia set of t^m is the unit circle, while the Julia set for T_m is the closed interval $[-2, 2]$. Now, if both f and g are conjugate to t^m (which is an endomorphism of \mathbb{G}_m), then our problem reduces to the classical Manin-Mumford question for \mathbb{G}_m^2 which was first proved by Lang [Lan65] (see also the results of Raynaud [Ray83a, Ray83b] for the general case of abelian varieties). A similar argument yields the desired result if both f and g are conjugate to T_m .

Therefore we reduced the proof of Theorem 3.1 to the case when both f and g are Lattès maps. In this case, there exist two elliptic curves E_1 and E_2 , there exist two projection maps $\pi_i : E_i \rightarrow \mathbb{P}^1$ for $i = 1, 2$, and there exist two isogenies $\omega_i : E_i \rightarrow E_i$ witnessing the fact that f and g are Lattès maps, i.e.,

$$\pi_1 \circ \omega_1 = f \circ \pi_1 \text{ and } \pi_2 \circ \omega_2 = g \circ \pi_2.$$

For each unramified point $x \in E_1 \times E_2$ for the morphism (π_1, π_2) , the induced map on the tangent space is an isomorphism. Now, the ramified locus for (π_1, π_2) is a finite union of divisors of $E_1 \times E_2$ of the form $\{a\} \times E_2$ or $E_1 \times \{b\}$. On the other hand, each irreducible component of $(\pi_1, \pi_2)^{-1}(\Delta)$ has dominant projection onto each E_i for $i = 1, 2$. Therefore, for each irreducible component V of $(\pi_1, \pi_2)^{-1}(\Delta)$, there exist at most finitely many ramified points for the map (π_1, π_2) . Also, for each $(x, x) \in \text{Prep}_{(f,g)}(\mathbb{P}^1 \times \mathbb{P}^1)$ each point in $(\pi_1, \pi_2)^{-1}(x, x) \cap V$ is preperiodic for the action of $\Phi := (\omega_1, \omega_2)$ on $E_1 \times E_2$. Since there are only finitely many ramified points of (π_1, π_2) lying on V , and because $(\pi_1, \pi_2)(V) = \Delta$, we obtain that there are infinitely many preperiodic points x under Φ lying on V such that the tangent subspace of V is preperiodic under the induced action of Φ on $\text{Gr}_1(T_{E_1 \times E_2, x})$. Because Conjecture 1.4 holds for algebraic group endomorphisms of abelian varieties, we conclude that V is preperiodic under Φ ; therefore Δ is preperiodic under (f, g) as desired. \square

Remark 3.3. It is worth noting that if not both f and g are Lattès maps, then we actually get that any line $L \subset \mathbb{P}^1 \times \mathbb{P}^1$ is preperiodic once it contains infinitely many preperiodic points; we do *not* need in this case the assumption about the preperiodicity of the tangent subspaces. We only need this last assumption when *both* f and g are Lattès maps.

Remark 3.4. We can extend Theorem 3.1 in a number of ways:

- (a) Over number fields, Theorem 3.1 also holds under the weaker “Bogomolov type” condition that there is an infinite nonrepeating sequence of points $(\alpha_i) \in \mathbb{P}^1(\mathbb{Q})$ such that

$$\lim_{i \rightarrow \infty} h_f(\alpha_i) = \lim_{i \rightarrow \infty} h_g(\alpha_i) = 0.$$

This follows immediately from the fact that the equidistribution results of [BR06, CL06, FRL06, Yua08] apply to such families of points.

- (b) Theorem 3.1 can be generalized to give a complete answer to Question 29 (1) of [KS07] in the case of rational functions defined over the complex numbers. In particular, if f and g are two rational functions satisfying $\text{Prep}(f) = \text{Prep}(g)$, then either $f^k g^\ell = f^{2k}$ for some ℓ and k , or f and g are both Chebychev, powering, or Lattès maps. This is an immediate application of [YZ] and [Lev90] as in our proof of Theorem 3.1.

We can also give the same answer as above to part (2) of Question 29 from [KS07]. Indeed, let $Z \subset \text{Prep}_f$ be an infinite subset such that $g(Z) \subset \text{Prep}_f$, we are going to prove that the invariant metrics $\|\cdot\|_f$ and $\|\cdot\|_g$ on $O(1)$ defined by f and g respectively are proportional. This implies the condition of Levin's theorem as above.

Let K be the subfield of \mathbb{C} generated by the coefficients of f and g and \bar{K} the algebraic closure of K in \mathbb{C} , then f and g are the base changes of endomorphisms f_K, g_K of \mathbb{P}_K^1 , and Z is the base change of a subset Z_K in $\text{Prep}_f(\bar{K})$. With Z_K replaced by the union of its conjugates under $\text{Gal}(\bar{K}/K)$, we may assume that Z_K is a union of closed points in \mathbb{P}_K^1 . Now we further extend f_K, g_K to endomorphisms

$$f_U, g_U : \mathbb{P}_U^1 \longrightarrow \mathbb{P}_U^1$$

over a variety U over \mathbb{Q} of finite type and with function field K . For each valuation v of K , the bundle $O(1)$ has invariant measures $\|\cdot\|_{v, f_U}$ and $\|\cdot\|_{v, g_U}$ defined for f_U and g_U on the Berkovich space X_v^{an} . Let Z_U be the union of Zariski closure of closed points in Z_K . For each geometric point $t \in U(\bar{\mathbb{Q}})$, the objects (f_U, g_U, Z_U) have reductions (f_t, g_t, Z_t) such that $Z_t \in \text{Prep}_{f_t}$ with $g_t(Z_t) \subset \text{Prep}_{f_t}$. By [YZ2], Theorem 4.7, Z_t is still infinite.

Applying the main result of [Mim97] we get that for *all* $x \in \text{Prep}_{f_t}$, we have $g_t(x) \in \text{Prep}_{f_t}$. Therefore $g_t(\text{Prep}_{f_t}) \subset \text{Prep}_{f_t}$. By [YZ], this implies that the invariant adelic metrics $\|\cdot\|_{f_t}$ and $\|\cdot\|_{g_t}$ on $O(1)$ defined by f_t and g_t respectively are proportional. In particular, this is true for all $t \in U(\bar{\mathbb{Q}})$ and the archimedean place corresponding to

$$\bar{\mathbb{Q}} \subset \bar{K} \subset \mathbb{C}.$$

By continuity of $\|\cdot\|_{f_U}$ and $\|\cdot\|_{g_U}$, this is true for *all* points in $U(\mathbb{C})$ including the point corresponding to original embedding $K \subset \mathbb{C}$. Thus we have shown that the invariant metrics $\|\cdot\|_f$ and $\|\cdot\|_g$ are proportional.

- (c) Theorem 3.1 extends easily to the case of any line in $(\mathbb{P}^1)^N$ under the action of (f_1, \dots, f_N) for arbitrary rational maps f_i ; the proof

in this case is reduced to the case $N = 2$ by projecting on any set of two coordinates of $(\mathbb{P}^1)^N$.

4. OTHER QUESTIONS

In this section we explore other questions related to the Dynamical Manin-Mumford problem. Each time we show that our question is valid for both polarizable group endomorphisms of abelian varieties, and for lines under the action of polarizable endomorphisms of $(\mathbb{P}^1)^m$ for any $m \in \mathbb{N}$.

Our first question refers to a classical principle in arithmetic geometry: for a given ambient variety X , one defines the notion of being *special* both for points and for irreducible subvarieties of X . Generically, one expects that if an irreducible subvariety Y of X contains a Zariski dense set of special points, then Y is a special subvariety. This principle lies at the heart of the classical Manin-Mumford conjecture - in that case, the ambient variety is an abelian variety G , the special points are the torsion points of G , while the special irreducible subvarieties are torsion translates of abelian subvarieties of G . For the Algebraic Dynamical Manin-Mumford Conjecture (see Conjecture 1.1), the ambient variety is any projective variety endowed with a polarizable endomorphism Φ , the special points are the preperiodic points for Φ , and the special irreducible subvarieties of X are the preperiodic irreducible subvarieties of X . Our counterexample from Section 1 shows that an irreducible subvariety may contain a Zariski dense set of special points (in the algebraic dynamics context) without being itself a special subvariety. However, it is natural to raise the following question regarding subvarieties containing a Zariski dense set of preperiodic points.

Question 4.1. *Let X be a projective variety and let $\Phi : X \rightarrow X$ be a polarizable endomorphism defined over the complex numbers. Assume X_1 and X_2 are any two subvarieties of X containing a Zariski dense set of preperiodic points under Φ . Then $X_1 \cap X_2$ contains a Zariski dense set of preperiodic points under Φ .*

In particular, our Question 4.1 asks that two (distinct) irreducible curves containing infinitely many preperiodic points intersect in finitely many preperiodic points.

Question 4.1 has a positive answer when X is an abelian variety and Φ is a polarizable algebraic group endomorphism. Indeed, in that case the preperiodic points for Φ are the torsion points of X (see Claim 2.2), and we know (by the classical Manin-Mumford conjecture proved by Raynaud [Ray83a, Ray83b]) that any irreducible subvariety of X which contains a Zariski dense set of torsion points must be a torsion translate of an abelian subvariety of X (note also that the torsion points are dense in each abelian subvariety).

We also have a positive answer for Question 4.1 when $X = (\mathbb{P}^1)^m$ (for some $m \in \mathbb{N}$) and X_1, X_2 are lines. Indeed, first we may reduce to the case $m = 2$ as observed in Remark 3.4 (c). Secondly, our proof of Theorem 3.1

shows that if $Y \subset \mathbb{P}^1 \times \mathbb{P}^1$ is a line which contains infinitely many preperiodic points for Φ , then either Y is preperiodic under Φ , or $\Phi = (f, g)$ for two Lattès maps f and g (see Remark 3.3). So, if Φ is not given by the action of two Lattès maps, we conclude that both lines X_1 and X_2 satisfying the hypothesis of Question 4.1 are preperiodic under Φ , and thus their point of intersection is also preperiodic under Φ . Now, if $\Phi = (f, g)$ and f, g are Lattès maps, then exactly as in the proof of Theorem 3.1 we may reduce the question from $\mathbb{P}^1 \times \mathbb{P}^1$ to a question on a split abelian surface. The reasoning from the above paragraph regarding abelian varieties finishes the proof for Question 4.1 for lines in $\mathbb{P}^1 \times \mathbb{P}^1$.

A positive answer to Question 4.1 has the following interesting consequence. Using the automatic uniformity theorem proved by Scanlon in [Sca04] we deduce that the set of preperiodic points of Φ satisfies automatic uniformity (since the hypothesis of Theorem 2.4 from [Sca04] is contained in a positive answer to Question 4.1). Therefore, for each positive integer d , there exists an integer $B(d)$ (depending on d , X and Φ only) such that for any subvariety Y contained in X of degree at most d (computed with respect to a fixed embedding of X into the projective space as given by the polarization), if $Y \cap \text{Prep}_\Phi(X)$ is not Zariski dense in Y , the Zariski closure of $Y \cap \text{Prep}_\Phi(X)$ has at most $B(d)$ irreducible components. In particular, if $Y \cap \text{Prep}_\Phi(X)$ is a finite set, then $|Y \cap \text{Prep}_\Phi(X)| \leq B(d)$.

Remark 4.2. It is worth noting that the hypothesis in Theorem 2.4 of [Sca04] is that if X_1 and X_2 contain a Zariski dense set of special points, then each irreducible component of $X_1 \cap X_2$ which contains at least one special point must contain a Zariski dense set of special points. Hence, one may consider a weakening of our Question 4.1 by asking that if X_1 and X_2 contain a Zariski dense set of preperiodic points for the endomorphism Φ , then each irreducible component of $X_1 \cap X_2$ which contains *at least one* preperiodic point for Φ must contain a Zariski dense set of preperiodic points for Φ .

Remark 4.3. Phrased in terms of canonical heights of varieties, Question 4.1 asks whether $X_1 \cap X_2$ must have canonical height zero if X_1 and X_2 each have canonical height zero, via results of [Zha95]. By the Bost-Gillet-Soulé arithmetical Bézout theorem ([BGS94]), one has the bound

$$(4.3.1) \quad h(X_1 \cdot X_2) \leq (\deg X_1)h(X_2) + (\deg X_2)h(X_1) + c(\deg X_1)(\deg X_2)$$

for h the height on \mathbb{P}^n coming from the Fubini-Study metric and X_1, X_2 two subvarieties that intersect properly. One might hope to show that the extra “error term” $c(\deg X_1)(\deg X_2)$ is zero; indeed, Bost-Gillet-Soulé state that they believe this “error term” is zero for the Fubini-Study metric and that this can be proved for linear subvarieties (see [BGS94, page 905]). However, Pascal Autissier has shown us that there is indeed such an error term for the canonical height associated to a polarized dynamical system even for lines with respect to the canonical height coming from the squaring map on \mathbb{P}^2 , so a different approach is needed. More precisely, let n be any integer larger

than 1, let L_1 be the line given by the equation $Ne - (n - 1)y = 0$, and let L_2 be the line given by the equation $nx + (n - 1)y - z = 0$. Since for points in \mathbb{P}^2 , the canonical height associated to the squaring map is the usual (naive) Weil height, and since $L_1 \cdot L_2 = [n - 1 : n : 2n(n - 1)]$, we conclude that $h(L_1 \cdot L_2) = \log(2n(n - 1))$. Using the Mahler measure one obtains $h(L_1) = h(L_2) = \log(n)$, and so, $h(L_1 \cdot L_2) = h(L_1) + h(L_2) + \log(2 \cdot (1 - 1/n))$. Hence there exists an error term in (4.3.1) for the canonical height associated to the squaring map. The best error term for the naive height on \mathbb{P}^2 should be $\log(2)$, i.e. the above examples are most likely asymptotically sharp.

We may also ask the following natural question which connects the dynamical Manin-Mumford and the Dynamical Bogomolov Conjectures in the context of algebraic dynamics (note that this connection is already known in the case of the classical Bogomolov and Manin-Mumford Conjecture for abelian varieties).

Question 4.4. *Let X be a projective variety over a number field, and let Φ be a polarizable endomorphism of X defined over \mathbb{Q} . We denote by \hat{h} the canonical height associated to Φ , and for each $\epsilon > 0$ we define*

$$X_\epsilon := \{x \in X(\bar{\mathbb{Q}}) : \hat{h}(x) < \epsilon\}.$$

Then each subvariety Y of X contains a Zariski dense set of preperiodic points for Φ if and only if for each $\epsilon > 0$ the subset $X_\epsilon \cap Y$ is Zariski dense in Y .

Since the preperiodic points for Φ are precisely the points of canonical height 0, we immediately note that once a subvariety Y contains a Zariski dense set of preperiodic points, then $Y \cap X_\epsilon$ is Zariski dense in Y .

Our proof of Theorem 3.1 yields a positive answer to Question 4.4 for lines embedded in $(\mathbb{P}^1)^m$ for any $m \in \mathbb{N}$. Indeed, as explained in Remark 3.4 (c), one reduces the question to lines Y inside $\mathbb{P}^1 \times \mathbb{P}^1$ under the action of $\Phi = (f, g)$, and then we have two cases. If not both f and g are Lattès maps, then Y is preperiodic as long as it contains infinitely many points of small height (see Remark 3.4 (a)). If both f and g are Lattès maps, then arguing as in the proof of Theorem 3.1 our problem reduces to Question 4.4 for abelian varieties, which we already know that it holds.

REFERENCES

- [BD] M. Baker and L. DeMarco, *Preperiodic points and unlikely intersections*, available online at arxiv.org/abs/0911.0918, 24 pages.
- [BGS94] J.-B. Bost, H. Gillet, and C. Soulé, *Heights of projective varieties and positive green forms*, J. Amer. Math. Soc. **7** (1994), 903–1027.
- [BR06] M. Baker and R. Rumely, *Equidistribution of small points, rational dynamics, and potential theory*, Ann. Inst. Fourier (Grenoble) **56** (2006), 625–688.
- [CL06] A. Chambert-Loir, *Mesures et équidistribution sur les espaces de Berkovich*, J. Reine Angew. Math. **595** (2006), 215–235.
- [DH93] A. Douady and J. H. Hubbard, *A proof of Thurston’s topological characterization of rational functions*, Acta Math. **171** (1993), 263–297.

- [Fak03] N. Fakhruddin, *Questions on self maps of algebraic varieties*, J. Ramanujan Math. Soc. **18** (2003), 109–122.
- [FRL06] C. Favre and J. Rivera-Letelier, *Équidistribution quantitative des points de petite hauteur sur la droite projective*, Math. Ann. **335** (2006), 311–361.
- [GT10] D. Ghioca and T. J. Tucker, *Proof of a dynamical Bogomolov conjecture for lines under polynomial actions*, Proc. Amer. Math. Soc. **138** (2010), 937–942.
- [GZ] D. Ghioca and M. E. Zieve, *Lattès maps in every characteristic*, in preparation.
- [KS07] S. Kawaguchi and J. H. Silverman, *Dynamics of projective morphisms having identical canonical heights*, Proc. Lond. Math. Soc. (3) **95** (2007), 519–544.
- [Lan65] S. Lang, *Division points on curves*, Ann. Mat. Pura Appl. (4) **70** (1965), 229–234.
- [Lev90] G. M. Levin, *Symmetries on Julia sets*, Mat. Zametki **48** (1990), 72–79, 159.
- [Mim97] A. Mimar, *On the preperiodic points of an endomorphism of $\mathbb{P}^1 \times \mathbb{P}^1$ which lie on a curve*, Ph.D. thesis, Columbia University, 1997.
- [Paz] F. Pazuki, *Zhang’s conjecture and squares of abelian surfaces*, to appear in C. R. Math. Acad. Sci. Paris.
- [Ray83a] M. Raynaud, *Courbes sur une variété abélienne et points de torsion*, Invent. Math. **71** (1983), 207–233.
- [Ray83b] ———, *Sous-variétés d’une variété abélienne et points de torsion*, Arithmetic and geometry, vol. I, Progr. Math., vol. 35, Birkhäuser, Boston, MA, 1983, pp. 327–352.
- [Sca04] Thomas Scanlon, *Automatic uniformity*, Int. Math. Res. Not. (2004), 3317–3326.
- [Sil94] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994.
- [SUZ97] L. Szpiro, E. Ullmo, and S. Zhang, *Équirépartition des petits points* (French) [Uniform distribution of small points], Invent. Math. **127** (1997), 337–347.
- [Ull98] E. Ullmo, *Positivité et discrétion des points algébriques des courbes*, Ann. of Math. (2) **147** (1998), 167–179.
- [Yua08] X. Yuan, *Big line bundles over arithmetic varieties*, Invent. Math. **173** (2008), 603–649.
- [YZ] X. Yuan and S.-W. Zhang, *Calabi theorem and algebraic dynamics*, preprint, 23 pages.
- [YZ2] X. Yuan and S.-W. Zhang, *Small points and Berkovich metrics*, preprint, 28 pages.
- [Zha95] S. Zhang, *Small points and adelic metrics*, J. Algebraic Geometry **4** (1995), 281–300.
- [Zha98] S. Zhang, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), 159–165.
- [Zha06] S. Zhang, *Distributions in Algebraic Dynamics*, Survey in Differential Geometry, vol. 10, International Press, 2006, pp. 381–430.

DRAGOS GHIOCA, DEPARTMENT OF MATHEMATICS, 1984 MATHEMATICS ROAD, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2
E-mail address: `dghioca@math.ubc.ca`

THOMAS TUCKER, DEPARTMENT OF MATHEMATICS, HYLAN BUILDING, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627
E-mail address: `ttucker@math.rochester.edu`

SHOUWU ZHANG, DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027
E-mail address: `szhang@math.columbia.edu`