

# On the cohomology of some twisted Shimura varieties of PEL-type

Cristian Virdol  
Department of Mathematics  
Columbia University

July 9, 2008

## 1 Introduction

In this article we consider the moduli problem of a "twist" of some simple Shimura varieties of PEL-type that appear in Kottwitz's papers [K1] and [K2] (these varieties have a hyperspecial level subgroup at some prime  $p$ ). The twist of the Shimura varieties is done by a mod  $q$  (for some prime  $q \neq p$ ) representation of the absolute Galois group of the reflex field of the Shimura varieties. Then we consider the Newton polygon stratification of the special fiber at  $p$  of twisted Shimura varieties and show that each Newton polygon stratum can be described in terms of the products of the reduced fibers of the corresponding PEL-type Rapoport-Zink spaces with certain smooth twisted Igusa varieties and of the action on them of a  $p$ -adic group that depends on the stratum. Then we extend our results to characteristic 0, and if  $l \neq p$  is a prime number, we describe the  $l$ -adic cohomology of proper twisted Shimura varieties of not necessarily hyperspecial level at  $p$ , in terms of the  $l$ -adic cohomology with compact supports of the twisted Igusa varieties and of the Rapoport-Zink spaces. These results generalize Mantovan's paper [M].

More exactly, we consider a simple  $\mathbb{Q}$ -algebra  $B$  and  $*$  a positive involution on  $B$ . Let  $V$  be a non-degenerate skew-Hermitian  $B$ -module and let  $G$  be the algebraic group over  $\mathbb{Q}$  of similitudes of the skew-Hermitian  $B$ -module  $V$  (other conditions are imposed on  $B$  and  $V$  to insure that  $G$  is unramified at some given rational prime  $p$ ; see §2 for details). Let  $h : \mathbb{C} \rightarrow \text{End}_B(V_{\mathbb{R}})$  be an  $\mathbb{R}$ -algebra homomorphism such that  $h(\bar{z}) = h(z)^*$  and the symmetric real-valued bilinear form  $(v, h(i)w)$  on  $V_{\mathbb{R}}$  is positive definite, where we have written  $*$  for the involution on  $\text{End}_B(V)$  obtained from the alternating form on  $V$ . By restricting  $h$  to  $\mathbb{C}^{\times}$  we obtain a homomorphism  $h : \mathbb{C}^{\times} \rightarrow G_{\mathbb{R}}$  of algebraic group over  $\mathbb{R}$ . For  $q$  a rational prime distinct from  $p$ , such that  $G$  is unramified at  $q$ , let  $\mathbf{S}_{\mathbf{K}}$  be the Shimura variety associated to  $(G, h^{-1})$  and to an open compact subgroup  $\mathbf{K} := K_q \times H$  of  $G(\mathbb{A}_f)$ , where  $K_q$  is a normal subgroup of a hyperspecial maximal compact subgroup  $H_q$  of  $G(\mathbb{Q}_q)$ ,  $H$  is a sufficiently

small open compact subgroup of  $G(\mathbb{A}_f^q)$ , and  $\mathbb{A}_f$  and  $\mathbb{A}_f^q$  are the finite part of the ring of adèles  $\mathbb{A}_{\mathbb{Q}}$  of  $\mathbb{Q}$  and the finite part of the ring of adèles of  $\mathbb{Q}$  away from  $q$ , respectively. Then  $\mathbf{S}_{\mathbf{K}}$  is a smooth quasi-projective variety defined over a number field  $E$  called the reflex field.

Consider the following moduli problem over  $O_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ . If  $S$  is a locally noetherian scheme over  $O_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ , then  $S$ -valued points on the moduli problem are the isomorphism classes of quadruples  $(A, \lambda, i, \bar{\eta}^p)$  where:  $A \rightarrow S$  is a projective abelian scheme over  $S$  up to prime-to- $p$  isogeny,  $\lambda : A \rightarrow \hat{A}$  is a polarization of  $A$ ,  $i : \mathcal{O}_B \rightarrow \text{End}(A)$  is a  $*$ -homomorphism for  $*$  on  $\mathcal{O}_B$  and the Rosati involution on  $\text{End}(A)$  obtained from  $\lambda$ , and  $\bar{\eta}^p$  is a level structure of type  $K^p$  on  $A$  and we require that  $(A, \lambda, i, \bar{\eta}^p)$  satisfy some "determinant condition" (see §3 below), which depends on  $h$ .

For  $K^p$  sufficiently small our moduli problem is representable by a quasi-projective scheme  $S_{K^p}$  over  $O_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ . The variety over  $E$  obtained from  $S_{K^p}$  is a disjoint union of  $|\ker^1(\mathbb{Q}, G)|$  copies of  $\mathbf{S}_{\mathbf{K}}$ , where our  $\mathbf{K}$  defined above has a decomposition of the form  $\mathbf{K} = K^p K_p$ , where  $K_p \subset G(\mathbb{Q}_p)$  is some hyperspecial maximal compact (see §2 below for details).

The varieties  $S_{K^p}$  and  $\mathbf{S}_{\mathbf{K}}$  have natural actions of  $H_q/K_q$ . For  $H$  sufficiently small these actions are free. We fix such a small group  $H$  and consider a continuous Galois representation  $\rho : \text{Gal}(\bar{\mathbb{Q}}/E) \rightarrow H_q/K_q$  unramified at  $p$  and let  $S'_{K^p}$  be the variety defined over  $E$  obtained from  $S_{K^p}$  via twisting by  $\rho$  composed with the natural action of  $H_q/K_q$  on  $S_{K^p}$  (see §2 for details). We show that each Newton polygon stratum at  $p$  of  $S'_{K^p}$  or  $\mathbf{S}'_{\mathbf{K}}$  can be described in terms of the products of the reduced fibers of the corresponding PEL-type Rapoport-Zink spaces with certain smooth twisted Igusa varieties and the action on them of a  $p$ -adic group that depends on the stratum (see theorems 9.4 and 9.5 and corollary 9.6 for details). Then we show (see theorem 9.7) that if  $l \neq p$  is a prime number, the  $l$ -adic cohomology of proper twisted Shimura varieties  $S'_K$  or  $\mathbf{S}'_{\mathbf{K}}$ , of level not necessarily hyperspecial at  $p$ , can be described in terms of the  $l$ -adic cohomology with compact supports of the twisted Igusa varieties and of the Rapoport-Zink spaces.

## 2 Shimura varieties of PEL type

Let  $p$  be a prime number. We are interested in certain Shimura varieties of PEL type that appear in [K2] having good reduction at all places of the reflex field lying above the prime  $p$  (in the next few sections we are following very closely [K2]).

We consider the PEL data  $(B, *, V, ( , ))$ , where:

- 1)  $B$  is a finite-dimensional simple algebra over  $\mathbb{Q}$ ,
- 2)  $*$  is a positive involution on  $B$  over  $\mathbb{Q}$ ,
- 3)  $V$  is a nonzero finitely generated left  $B$ -module,
- 4)  $( , )$  is a nondegenerate  $\mathbb{Q}$ -valued  $*$ -Hermitian alternating pairing on  $V$ ,

that satisfy the following conditions:

a) there exists a  $\mathbb{Z}_{(p)}$ -order  $\mathcal{O}_B$  on  $B$  whose  $p$ -adic completion is a maximal order in  $B_{\mathbb{Q}_p}$  and which is preserved by  $*$ ;

b) there exists a lattice  $\Lambda$  in  $V_{\mathbb{Q}_p}$  which is self-dual for  $(\ , \ )$  and is preserved by  $\mathcal{O}_B$ ;

c)  $B_{\mathbb{Q}_p}$  is a product of matrix algebras over unramified extensions of  $\mathbb{Q}_p$ .

Let  $F$  be the center of  $B$ . Then c) implies that  $F$  is a number field unramified at  $p$ . Let  $D$  be the  $\mathbb{Q}$ -algebra  $\text{End}_B(V)$ ; it is a simple  $\mathbb{Q}$ -algebra with center  $F$  and has an involution  $*$  (the adjoint map for  $(\cdot, \cdot)$ ). Let  $G$  be the algebraic group over  $\mathbb{Q}$  of the  $B$ -linear automorphisms of  $V$ , which preserve the pairing  $(\ , \ )$  up to a scalar multiple; then for any  $\mathbb{Q}$ -algebra  $R$  we have  $G(R) = \{x \in D \otimes_{\mathbb{Q}} R \mid xx^* \in R^\times\}$ . Let  $h : \mathbb{C} \rightarrow D_{\mathbb{R}}$  be an  $\mathbb{R}$ -algebra homomorphism such that  $h(\bar{z}) = h(z)^*$  and the symmetric real-valued bilinear form  $(v, h(i)w)$  on  $V_{\mathbb{R}}$  is positive definite, i.e  $(v, h(i)v) > 0$  for all nonzero  $v \in V_{\mathbb{R}}$ .

The homomorphism  $h$  determines a decomposition of the  $B_{\mathbb{C}}$ -module  $V \otimes_{\mathbb{Q}} \mathbb{C} = V_1 \oplus V_2$ , where  $V_1$  (respectively,  $V_2$ ) is the subspace of  $V \otimes_{\mathbb{Q}} \mathbb{C}$  on which  $h(z)$  acts by  $z$  (respectively, by  $\bar{z}$ ). The field of definition of the isomorphism class of the complex representation  $V_1$  of  $B$  is a number field  $E$  called the *reflex field*, whose ring of integers we denote by  $\mathcal{O}_E$ .

We denote also by  $h$  the restriction of  $h$  to  $\mathbb{C}^\times$ , regarded as a homomorphism of real algebraic groups  $\mathbb{C}^\times \rightarrow G$ . Then from Lemma 4.1 of [K2], we know that the pair  $(G, h^{-1})$  satisfies the conditions (1.5.1), (1.5.2), (1.5.3) of [D]. Therefore the set  $X_\infty$  of conjugates of  $h^{-1}$  under  $G(\mathbb{R})$  is a finite union of copies of the symmetric space for the identity component of  $G(\mathbb{R})$  and this symmetric space is of Hermitian type. Let  $K^p$  be a sufficiently small open compact subgroup of  $G(\mathbb{A}_f^p)$ , where  $\mathbb{A}_f^p$  denotes the ring of finite adeles of  $\mathbb{Q}$  with trivial  $p$ -adic component. The stabilizer  $K_p$  of  $\Lambda$  in  $G(\mathbb{Q}_p)$  is a hyperspecial maximal compact subgroup of  $G(\mathbb{Q}_p)$ . We define  $\mathbf{K} := K^p K_p$ . We denote by  $\mathbf{S}_{\mathbf{K}}$  the Shimura variety associated to  $(G, h^{-1})$  and  $\mathbf{K}$ . Then  $\mathbf{S}_{\mathbf{K}}$  is a smooth quasi-projective variety and has a canonical model defined over  $\mathcal{O}_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ .

For any compact open subgroup  $K_1^p$  of  $K^p$  there is an etale covering  $\mathbf{S}_{\mathbf{K}_1} \rightarrow \mathbf{S}_{\mathbf{K}}$ , where  $\mathbf{K}_1 := K_1^p K_p$  and this covering map is Galois with Galois group  $\mathbf{K}/\mathbf{K}_1 = K^p/K_1^p$  if  $K_1^p$  is normal in  $K^p$ . For  $g \in G(\mathbb{A}_f^p)$  there is a natural isomorphism  $\mathbf{S}_{\mathbf{K}} \rightarrow \mathbf{S}_{g^{-1}\mathbf{K}g}$ .

### 3 Moduli problem

In this section we formulate the PEL moduli problem having good reduction at all places of the reflex field lying over a given rational prime  $p$  (in the case D of [K2], page 375, we assume that  $p > 2$ ). For an open compact subgroup  $K^p \subset G(\mathbb{A}_f^p)$ , we consider the set-valued contravariant functor from the category of locally noetherian schemes  $S$  over  $\mathcal{O}_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$  that associates to a connected scheme  $S$  and to a geometric point  $s$  on  $S$  the set of isomorphism classes of quadruples  $(A, \lambda, i, \bar{\eta}^p)$ , where

- 1)  $A$  is a projective abelian scheme over  $S$ ;

- 2)  $\lambda : A \rightarrow \hat{A}$  is a prime-to- $p$  polarization;
- 3)  $i : \mathcal{O}_B \hookrightarrow \text{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$  is a morphism of  $\mathbb{Z}_{(p)}$ -algebras such that  $\lambda \circ i(b^*) = i(\hat{b}) \circ \lambda$  and  $\det(b, \text{Lie}(A)) = \det(b, V_1)$  for all  $b \in \mathcal{O}_B$ ;
- 4)  $\bar{\eta}^p$  is a  $\pi_1(S, s)$ -invariant  $K^p$ -orbit of isomorphisms of skew-Hermitian  $B$ -modules  $\eta^p : V \otimes_{\mathbb{Q}} \mathbb{A}_f^p \rightarrow H_1(A_s, \mathbb{A}_f^p)$ , where  $H_1(A_s, \mathbb{A}_f^p)$  is the Tate  $\mathbb{A}_f^p$ -module of the abelian variety  $A_s$ .

We remark that the level structure  $\bar{\eta}^p$  of type  $K^p$  is independent of the choice of the geometric point  $s$  of  $S$  and thus our functor is independent of  $s$ . For  $K^p$  sufficiently small this functor is representable by a quasi-projective scheme  $S_{K^p}$  over  $O_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ .

Two quadruples  $(A, \lambda, i, \bar{\eta}^p)$  and  $(A', \lambda', i', \bar{\eta}'^p)$  are said to be isomorphic if there is a prime-to- $p$  isogeny  $\varphi : A \rightarrow A'$ , commuting with the action of  $\mathcal{O}_B$ , carrying  $\bar{\eta}^p$  into  $\bar{\eta}'^p$ , and carrying  $\lambda$  into a scalar multiple of  $\lambda'$  (a scalar in  $\mathbb{Z}_{(p)}^\times$ ) i.e.  $\varphi^* \lambda' = c\lambda$  for some  $c \in \mathbb{Z}_{(p)}^\times$ , where  $\varphi^* \lambda' = \hat{\varphi} \circ \lambda' \circ \varphi$  and  $\hat{\varphi} : \hat{A}' \rightarrow \hat{A}$  is the dual of  $\varphi$ .

For any compact open subgroup  $K_1^p$  of  $K^p$  there is an etale covering  $S_{K_1^p} \rightarrow S_{K^p}$ , sending  $(A, \lambda, i, (\bar{\eta}_1^p)^p)$  to  $(A, \lambda, i, \bar{\eta}^p)$ , where  $(\bar{\eta}_1^p)^p$  denotes the  $K_1^p$ -orbit of  $\eta^p$ , and this covering map is Galois with Galois group  $K^p/K_1^p$  if  $K_1^p$  is normal in  $K^p$ . For  $g \in G(\mathbb{A}_f^p)$  there is an isomorphism  $S_{K^p} \rightarrow S_{g^{-1}K^p g}$  sending  $(A, \lambda, i, \bar{\eta}^p)$  to  $(A, \lambda, i, \overline{\eta^p g})$ .

## 4 Complex points

We want to describe the set of complex points of  $S_{K^p}(\mathbb{C})$  relative to the map  $O_E \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)} \rightarrow \mathbb{C}$  induced by the inclusion  $E \subset \mathbb{C}$ . Hence we are interested in quadruples  $(A, \lambda, i, \bar{\eta}^p)$ , where this time  $A$  is an abelian variety over  $\mathbb{C}$  up to prime-to- $p$  isogeny. Then the homology group  $H := H_1(A, \mathbb{Q})$  is a skew-Hermitian  $B$ -module, with the  $B$ -action coming from  $i$  and the alternating form coming from  $\lambda$ .

From §8 of [K2], we know that for every place  $v$  of  $\mathbb{Q}$  the skew-Hermitian  $B_{\mathbb{Q}_v}$ -modules  $H_{\mathbb{Q}_v}$  and  $V_{\mathbb{Q}_v}$  are isomorphic.

The isomorphism classes of skew-Hermitian  $B$ -modules of the same dimension as  $V$  are classified by  $H^1(\mathbb{Q}, G)$ , and thus we get there are finitely many isomorphic classes of skew-Hermitian  $B$ -modules  $(V', (\cdot, \cdot)')$  such that  $V'_{\mathbb{Q}_p}$  and  $V_{\mathbb{Q}_p}$  are isomorphic for all places  $v$  of  $\mathbb{Q}$ . Let  $V^{(1)}, \dots, V^{(m)}$  be some representatives in these isomorphism classes. We fix local isomorphisms  $V_{\mathbb{Q}_p}^{(i)} \cong V_{\mathbb{Q}_p}$  for all places  $v$  of  $\mathbb{Q}$  and let  $G^{(i)}$  denote the group of automorphisms of  $V^{(i)}$  and thus we have  $G_{\mathbb{Q}_p}^{(i)} \cong G_{\mathbb{Q}_p}$  for all  $p$ . We obtain that the set  $S_{K^p}(\mathbb{C})$  is a disjoint union  $\coprod_{i=1}^m \mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$ , where  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$  consists of quadruples  $(A, \lambda, i, \bar{\eta}^p)$  such that  $H$  is isomorphic to  $V^{(i)}$ , and hence it is enough to study each set  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$ .

Assume that  $V^{(1)}$  is  $V$  and we start studying  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(1)}$ . Let  $(A, \lambda, i, \bar{\eta}^p)$  be a point of  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(1)}$  and choose an isomorphism between  $H$  and  $V$ . Choosing a level structure  $\bar{\eta}^p$  is equivalent to choosing an element of  $G(\mathbb{A}_f^p)/K^p$ . The

complex structure on  $H_{\mathbb{R}} = \text{Lie}(A)$  determines a complex structure on  $V_{\mathbb{R}}$  and yields a homomorphism  $h' : \mathbb{C} \rightarrow D_{\mathbb{R}}$ . From Lemma 4.2 of [K2], we get that  $h'$  belongs to the set  $X_{\infty}$  of  $G(\mathbb{R})$ -conjugates of  $h$ . The self-dual  $\mathcal{O}_B$ -lattice  $\lambda' := H_1(A, \mathbb{Z}_p)$  in  $H_{\mathbb{Q}_p}$  becomes an  $\mathcal{O}_B$ -lattice in  $V_{\mathbb{Q}_p}$  that it is self-dual up to a scalar. The group  $K_p$  is the stabilizer of the self-dual  $\mathcal{O}_B$ -lattice  $\Lambda$  in  $G(\mathbb{Q}_p)$ . From Corollary 7.3 of [K2], we know that there exists  $g \in G(\mathbb{Q}_p)$  such that  $\lambda' = g\Lambda$ , and  $g$  determines a well-defined element of  $G(\mathbb{Q}_p)/K_p$ . The choice of isomorphism between  $H$  and  $V$  gives us a well-defined element of  $(G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty}$ . Two isomorphisms between  $H$  and  $V$  differ by an automorphism of  $V$ , thus by an element of  $G(\mathbb{Q})$  and the corresponding elements of  $(G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty}$  are in the same  $G(\mathbb{Q})$ -orbit. Hence  $(A, \lambda, i, \bar{\eta}^p)$  gives rise to a well-defined element of the quotient  $G(\mathbb{Q}) \backslash (G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty}$ . We get a bijection from  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(1)}$  to  $G(\mathbb{Q}) \backslash ((G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty})$ . Replacing  $V$  by  $V^{(i)}$ , we obtain a bijection from  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$  to  $G(\mathbb{Q})^{(i)} \backslash ((G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty})$ . Then  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$  is the set of complex points of a variety  $\mathbf{S}_{\mathbf{K}}^{(i)}$  over  $E$ , which is isomorphic to  $\mathbf{S}_{\mathbf{K}}$ .

Hence, we get (see §8 of [K2]) that variety  $S_{K^p}$  over  $E$  is a disjoint union of  $|\ker^1(\mathbb{Q}, G)|$ -copies of  $\mathbf{S}_{\mathbf{K}}$ .

## 5 Twisted Shimura varieties

Let  $q \neq p$  be a rational prime such that  $G$  is unramified at  $q$ , i.e.  $G$  is quasi-split over  $\mathbb{Q}_q$  and split over an unramified extension of  $\mathbb{Q}_q$ . Consider now our  $\mathbf{K}$  of the form  $\mathbf{K} := K_q \times H$ , where  $K_q$  is a normal subgroup of a hyperspecial maximal compact subgroup  $H_q$  of  $G(\mathbb{Q}_q)$ . Then it is well known (see for example Corollary 1.4.1.3 of [CA]) that for  $H$  sufficiently small, the group  $H_q/K_q$  acts freely on  $S_{K^p}$ . The action of  $H_q/K_q$  on  $S_{K^p}(\mathbb{C})$  can be described in the following way: the set  $S_{K^p}(\mathbb{C})$  is a disjoint union  $\coprod_{i=1}^m \mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)}$  (a disjoint union of  $|\ker^1(\mathbb{Q}, G)|$  copies of  $\mathbf{S}_{\mathbf{K}}(\mathbb{C})$ ) and  $H_q/K_q$  acts on each

$$\mathbf{S}_{\mathbf{K}}(\mathbb{C})^{(i)} = G(\mathbb{Q})^{(i)} \backslash ((G(\mathbb{A}_f)/\mathbf{K}) \times X_{\infty})$$

where  $H_q \hookrightarrow G(\mathbb{A}_{\mathbb{Q}})$  by  $\alpha \mapsto (1, \dots, \alpha, 1, \dots, 1)$ ,  $\alpha$  at the  $q$ -component, and the action of an element  $\alpha \in H_q$  is given by the right multiplication at the  $q$ -component.

We fix a continuous representation

$$\rho : \text{Gal}(\bar{\mathbb{Q}}/E) \rightarrow H_q/K_q,$$

unramified at  $p$ . Let  $L$  be the finite Galois extension of  $E$  defined by  $L := (\bar{\mathbb{Q}})^{\text{Ker}(\rho)}$ .

Let

$$X = S_{K^p} \times_{\text{Spec}(E)} \text{Spec}(L).$$

The group  $H_q/K_q$  acts on  $S_{K^p}$ . Since  $\rho : \text{Gal}(L/E) \hookrightarrow H_q/K_q$ , the group  $\text{Gal}(L/E)$  acts on  $S_{K^p}$ . We denote this action of  $\text{Gal}(L/E)$  on  $S_{K^p}$  by  $\rho'$ . The

Galois group  $\text{Gal}(L/E)$  has a natural action on  $\text{Spec}(L)$  and we can descend via the quotient process  $X$  to  $S'_{K^p}/\text{Spec}(E)$  using the diagonal action

$$\text{Gal}(L/E) \ni \sigma \rightarrow \rho'(\sigma) \otimes \sigma$$

on  $X$ . Thus, we obtain a quasi-projective variety  $S'_{K^p}/\text{Spec}(E)$  having the same dimension as  $S_{K^p}$ . This is the twisted Shimura variety that we mentioned in the title.

Now we describe the PEL moduli problem for  $S'_{K^p}$ . We consider again quadruples as in §3 above, only  $\eta^p$  will be replaced by  $\eta'^p$ . Thus, with the same notations as above, for the twisted Shimura variety  $S'_{K^p}$  a level structure of type  $K^p$  on  $A$  is a  $K^p$ -orbit  $\bar{\eta}'^p$  of isomorphisms  $\eta'^p : V \otimes_{\mathbb{Q}} \mathbb{A}_f^p \rightarrow H_1(A_s, \mathbb{A}_f^p)$  of skew-Hermitian  $B$ -modules such that  $\bar{\eta}'^p$  is fixed by the action  $\rho(\tau) \times \tau$  for all  $\tau \in \pi_1(S, s)$ , where  $\rho(\tau)$  acts on the left using the natural action of  $H_q/K_q$  and  $\tau$  acts on the right using the natural action of  $\tau \in \pi_1(S, s)$  on  $A_s$ .

## 6 The Newton polygon stratification

For a place  $v$  of  $E$  above  $p$ , we denote by  $E_v$  the completion of  $E$  at  $v$ . From the condition c) of §2, we get that  $E_v/\mathbb{Q}_p$  is an unramified extension. Let  $k/\mathbb{F}_p$  be the residue field of  $\mathcal{O}_{E_v}$ ,  $|k| = q = p^f$ , and let  $\sigma : \bar{\mathbb{F}}_p \rightarrow \bar{\mathbb{F}}_p$  be the Frobenius map  $x \mapsto x^p$ .

For  $K^p \subset G(\mathbb{A}_f^p)$  a sufficiently small open compact open subgroup, we write  $S = S_{K^p}$  and  $\bar{S} = \bar{S}_{K^p}/k$  for its reduction. If  $\mathcal{A}$  is the universal abelian variety over  $\bar{S}$ , we define  $\mathcal{G} = \mathcal{A}[p^\infty]$ . Then  $\mathcal{G}$  is a Barsotti-Tate group endowed with a quasi-polarization and a compatible action (i.e it satisfies the determinant condition as in [K2]) of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$  of a polarized Barsotti-completion of  $\mathcal{O}_B$  in  $B_{\mathbb{Q}_p}$ . We call a Barsotti-Tate group endowed with a quasi-polarization and of a compatible action of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$  a Barsotti-Tate group with *additional structures*.

We denote by  $B(G)$  the set of  $\sigma$ -conjugacy classes in  $G(L)$ , where  $L = \mathbb{Q}_p^{nr} \supset E_v$ . Hence,  $B(G)$  is the set of isogeny classes over  $\bar{\mathbb{F}}_p \supset k$  of polarized Barsotti-Tate groups, which are endowed with a compatible action of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$ . If  $\mathcal{G}$  is a Barsotti-Tate group with additional structures, we denote by  $b(\mathcal{G}) \in B(G)$  the corresponding isogeny class.

For any element  $b \in B(G)$  define the strata

$$\bar{S}^{(b)} = \{x \in \bar{S} \mid b(\mathcal{G}_{\bar{x}}) = b\} \subset \bar{S}.$$

Then  $\bar{S}^{(b)} \subset \bar{S}$  is a reduced subscheme, and as the level  $K^p$  varies, these strata are preserved by the natural projection  $\bar{S}_{K_1^p} \rightarrow \bar{S}_{K^p}$  (for  $K_1^p \subset K^p$ ) and also by the morphism  $g : \bar{S}_{K^p} \rightarrow \bar{S}_{gK^pg^{-1}}$  for all  $g \in G(\mathbb{A}_f^p)$ .

We know (proposition 1 of [M]):

**Proposition 6.1.** *Let  $\Sigma/\bar{\mathbb{F}}_p \supset k$  be a polarized Barsotti-Tate group endowed with a compatible action of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$ , and denote by  $b \in B(G)$  its isogeny class. We*

define

$$C_\Sigma = \{x \in \bar{S} \mid \mathcal{G}_{\bar{x}} \simeq \Sigma_{\frac{\bar{x}}{k(x)}}\} \subset \bar{S}^{(b)} \times_k \bar{\mathbb{F}}_p.$$

This is a closed subset of the stratum  $\bar{S}^{(b)}$ ; and as a subscheme of  $\bar{S}^{(b)} \times_k \bar{\mathbb{F}}_p$  endowed with the induced reduced structure, it is smooth.

Then  $C_\Sigma$  is called a leaf. As the level  $K^p$  varies, the leaves are preserved by both the natural projection  $\bar{S}_{K_1^p} \rightarrow \bar{S}_{K^p}$  (for  $K_1^p \subset K^p$ ) and also by the morphism  $g : \bar{S}_{K^p} \rightarrow \bar{S}_{gK^p g^{-1}}$  for all  $g \in G(\mathbb{A}_f^p)$ .

The universal Barsotti-Tate group  $\mathcal{G}$  admits a slope filtration when restricted to the leaves (see page 239 of [Z]).

If  $\Sigma/\bar{\mathbb{F}}_p$  is a completely slope-divisible Barsotti-Tate group with additional structures, we write  $b \in B(G)$  for its isogeny class and  $1 \geq \lambda_1 > \lambda_2 > \cdots > \lambda_k \geq 0$  for the slopes of its Newton polygon. Then  $\Sigma = \bigoplus_i \Sigma^i$ , where for each  $i$ ,  $1 \leq i \leq k$ ,  $\Sigma^i$  is the slope-divisible isoclinic Barsotti-Tate group of the slope  $\lambda_i$ .

For  $C = C_\Sigma$  a leaf contained in  $\bar{S}^{(b)}$  associated to a completely slope-divisible Barsotti-Tate group  $\Sigma$  with additional structures, we denote by  $\mathcal{G}$  the universal Barsotti-Tate group over  $C$ . Then  $\mathcal{G}$  is a completely slope-divisible Barsotti-Tate group with slope filtration

$$0 \subset \mathcal{G}_1 \subset \cdots \subset \mathcal{G}_k = \mathcal{G},$$

where the subquotients  $\mathcal{G}^i = \mathcal{G}_i/\mathcal{G}_{i-1}$  are isoclinic slope-divisible Barsotti-Tate groups of slope  $\lambda_i$ . Then  $\mathcal{G}_i$  and  $\mathcal{G}^i$  inherit an action of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$ .

## 7 Igusa varieties

For a completely slope-divisible Barsotti-Tate group  $\Sigma$  as above, we denote by  $b = b(\Sigma) \in B(G)$  the corresponding isogeny class, by  $C_b = C_\Sigma \subset \bar{S}^{(b)}$  the corresponding leaf and by  $\mathcal{G}$  the universal Barsotti-Tate group over  $C_b$ .

**Definition 7.1.** *For any  $m \geq 1$ , the Igusa variety of level  $m$ ,  $J_{b,m} \rightarrow C_b$  is the universal space for the existence of isomorphisms*

$$j_{m,i} : \Sigma^i[p^m] \rightarrow \mathcal{G}^i[p^m]$$

of finite flat group schemes over  $C_b$  such that

- (1) they extend étale locally to any level  $m' \geq m$ ;
- (2) they are  $\mathcal{O}_{B_{\mathbb{Q}_p}}$ -equivariant;
- (3) they commute up to  $(\mathbb{Z}/p^m)^\times$ -multiple with the isomorphisms induced by the quasi-polarizations on  $\mathcal{G}$  and  $\Sigma$ .

The group  $\Gamma_b = \text{Aut}(\Sigma) = \prod_i \text{Aut}(\Sigma^i)$  acts on  $J_{b,m}$ . If  $\gamma = \bigoplus_i \gamma^i \in \Gamma_b = \text{Aut}(\Sigma) = \prod_i \text{Aut}(\Sigma^i)$ , then  $\gamma$  acts on  $J_{b,m}$  by

$$(A, \lambda, i, \bar{\eta}^p; j_{m,1}, \dots, j_{m,k}) \mapsto (A, \lambda, i, \bar{\eta}^p; j_{m,1} \circ \gamma|_{[p^m]}, \dots, j_{m,k} \circ \gamma|_{[p^m]}),$$

where a point of  $J_{b,m}$  is represented by the data  $(A, \lambda, i, \bar{\eta}^p; j_{m,1}, \dots, j_{m,k})$ , as in the definition 7.1. This action factors through the quotient  $\Gamma_{b,m}$  of  $\Gamma_b$  by the subgroup of automorphisms of  $\Sigma$  which restrict to the identity on  $\Sigma[p^m]$ .

We know (see proposition 4 of [M]):

**Proposition 7.2.** *For any  $m \geq 1$ , the Igusa variety  $J_{b,m} \rightarrow C_b$  is finite étale and Galois with Galois group  $\Gamma_{b,m}$ . In particular, the Igusa varieties are smooth.*

We have natural projections

$$q_b = q_{b,m',m} : J_{b,m'} \rightarrow J_{b,m}, \quad \forall m' \geq m,$$

which correspond to restricting the isomorphisms  $j_{m',i}$  to the  $p^m$ -torsion subgroups for all  $i$ . The projections  $q_b$  are invariant under the action of  $\Gamma_b$  on the Igusa varieties.

Thus, as the level  $m$  varies, the Igusa varieties  $J_{b,m}$  form a projective system under the morphisms  $q_b$  which is endowed with an action of  $\Gamma_b$ . In §4 of [M] it is shown that this action naturally extends to a submonoid  $S_b \supset \Gamma_b$  of the group  $T_b$  of the quasi-self-isogenies of  $\Sigma$ , preserving its additional structures.

Now we define  $S_b$ . Any  $\rho \in T_b$  can be written as  $\rho = \oplus_i \rho^i$ , where  $\rho^i$  is the quasi-self-isogeny of  $\Sigma^i$  obtained by restricting  $\rho$  to  $\Sigma^i$ . If  $\rho^{-1}$  is an isogeny, then, for each  $i = 1, \dots, k$ , define two integers  $e_i = e_i(\rho) \geq f_i = f_i(\rho) \geq 0$  to be, respectively, the minimal and maximal integers such that

$$\ker[p^{f_i}] \subset \ker[\rho^{i-1}] \subset \ker[p^{e_i}].$$

Define

$$S_b = \{\rho \in T_b \mid \rho^{-1} \text{ is an isogeny, } f_{i-1}(\rho) \geq e_i(\rho), \forall i \geq 2\}.$$

Then  $S_b$  is a submonoid of  $T_b$ , and  $p^{-1}, \text{fr}^{-B} = \oplus_i p^{-\lambda_i B} \in S_b$ , where the integer  $B \geq 1$  is defined such that for all  $i = 1, \dots, k$ ,  $F^B p^{-\lambda_i B} : \Sigma^i \rightarrow \Sigma^{i(p^B)}$  are well defined isogenies, where  $F : \Sigma \rightarrow \Sigma^{(p)}$  is the Frobenius element. Also we know that  $T_b = \langle S_b, p, \text{fr}^B \rangle$ , where  $\text{fr}^{-B} := \oplus_i p^{-\lambda_i B} \in S_b$  (see lemma 2.11 of [M1]).

The action of  $\Gamma_b$  on the system of Igusa varieties extends to  $S_b \subset T_b$  (lemma 5 of [M]):

**Lemma 7.3.** *Let  $(\mathcal{A}, \lambda, i, \bar{\eta}^p; j_{m,1}, \dots, j_{m,k})$  denote the universal object over the Igusa variety  $J_{b,m}$ , let  $\rho = \oplus_i \rho_i \in S_b$ , and assume  $m \geq e = e_1(\rho)$ .*

*There exists a unique finite flat subgroup  $\mathcal{K}_\rho \subset \mathcal{G}[p^e]$  such that the corresponding subgroups inside the isoclinic subquotients  $\mathcal{G}^i$  of  $\mathcal{G}$  are  $j_{m,i}(\ker(\rho_i^{-1}))$  for all  $i$ .*

*Moreover, the quotient abelian variety  $\mathcal{A}/\mathcal{K}_\rho$  inherits the additional structures of  $\mathcal{A}/J_{b,m}$  and thus gives rise to a morphism between the Igusa varieties:*

$$\rho : J_{b,m} \rightarrow J_{b,m-e}.$$

For all  $\rho \in S_b$ , the morphisms  $\rho : J_{b,m} \rightarrow J_{b,m-e}$  are compatible with the projections among Igusa varieties of different levels  $m \geq e = e_1(\rho)$  and they give rise to an action of the monoid  $S_b$  on the system of Igusa varieties which extends the action of  $\Gamma_b$ .

One can to define a  $\sigma$ -semilinear action of Frobenius on the varieties as arising from the Frobenius morphism of the Barsotti-Tate group  $\Sigma$ ,  $F : \Sigma \rightarrow \Sigma^{(p)}$ , that is, to define

$$\text{Frob} : J_{b,m} \rightarrow J_{b,m-1}$$

to be the  $\sigma$ -semilinear morphism corresponding to the quotient abelian variety  $\mathcal{A}^{(p)} = \mathcal{A}/\mathcal{A}[p]$  endowed with the induced additional structures. Then Frob is the absolute Frobenius of  $J_{b,m}$  followed by the projection  $q_{b,m,m-1}$ .

By reintroducing in our notation the level  $K^p$ , we write

$$J_{b,K^p,m} \rightarrow C_{b,K^p} \subset \bar{S}_{K^p}^{(b)} \subset \bar{S}_{K^p}$$

for the Igusa varieties of level  $K^p$  and level  $m$  at  $p$ . For any  $b \in B(G)$  and  $m \geq 1$ , we the group  $G(\mathbb{A}_f^p)$  acts on the Igusa varieties

$$\forall g \in G(\mathbb{A}_f^p) : J_{b,K^p,m} \rightarrow J_{b,g^{-1}K^p,g,m},$$

$$(A, \lambda, i, \bar{\eta}^p; j_{m,1}, \dots, j_{m,k}) \mapsto (A, \lambda, i, \overline{\eta^p \circ g}; j_{m,1}, \dots, j_{m,k}).$$

This action is compatible under the projection map with the action of  $G(\mathbb{A}_f^p)$  on the leaves  $C_{b,K^p}$ , commutes with the natural projections among the Igusa varieties as the level  $m$  at  $p$  varies, and also commutes with the action of  $S_b$  defined above.

Thus, for  $l \neq p$  be a prime number, the direct limit of cohomologies with compact support

$$H_c^i(J_b, \mathbb{Q}_l) = \lim_{\rightarrow K^p, m} H_c^i(J_{b,K^p,m}, \mathbb{Q}_l), \quad \forall i,$$

is endowed with an action of the submonoid  $S_b \times G(\mathbb{A}_f^p) \subset T_b \times G(\mathbb{A}_f^p)$ . This action cannot be extended to an action of  $T_b \times G(\mathbb{A}_f^p)$  on the varieties, but it is possible to extend it on the cohomology groups. Because of the equality  $T_b = \langle S_b, p, \text{fr}^B \rangle$ , it suffice to prove that the action of  $p^{-1}, \text{fr}^{-B} \in S_b$  on the cohomology groups of Igusa varieties is invertible.

We know (lemma 6 of [M]):

**Lemma 7.4.** *Maintaining the above notation, we have the following:*

- (1) for any  $m \geq 1$ ,  $(p^{-1}, 1) \in S_b \times G(\mathbb{A}_f^p)$  acts on the Igusa varieties as  $(p, 1) \circ q_{b,m,m-1}$ ;
- (2) for any  $m \geq \lambda_1 B$ ,  $(\text{fr}^{-B}, 1) \in S_b \times G(\mathbb{A}_f^p)$  acts on the Igusa varieties as  $q_{b,m,m-\lambda_1 B} \circ \text{Frob}$ .

Thus we get the first part of the following proposition:

**Proposition 7.5.** *The action of the submonoid  $S_b \times G(\mathbb{A}_f^p)$  on the cohomology groups of the Igusa varieties extends uniquely to an action of the group  $T_b \times G(\mathbb{A}_f^p)$ .*

*Moreover, as representations of  $T_b \times G(\mathbb{A}_f^p)$ , the  $H_c^i(J_b, \mathbb{Q}_l)$  are admissible for all  $i \geq 0$ .*

The second part regarding the admissibility of the representations follows from

$$H_c^i(J_b, \mathbb{Q}_l)^{\Gamma_{b,m} \times K^p} = H_c^i(J_{b,K^p,m}, \mathbb{Q}_l),$$

where  $\Gamma_{b,m}$  is the subgroup of  $T_b$  consisting of the isomorphisms of  $\Sigma$  which restrict to the identity on  $\Sigma[p^m]$ .

## 8 Twisted leaves and twisted Igusa varieties

Let  $q \neq p$  be the rational prime, and let  $K_q \subset G(\mathbb{Q}_q)$  be the open compact subgroup considered in §5. Let  $K^p \subset G(\mathbb{A}_f^p)$  be a sufficiently small open compact subgroup of the form  $K^p = K_q \times K^{pq}$ , where  $K^{pq}$  is an open compact subgroup of  $G(\mathbb{A}_f^{pq})$ , and we write  $S' = S'_{K^p}$  for the twisted variety defined in §5 and  $\bar{S}' = \bar{S}'_{K^p}/k$  for its reduction.

Using the same notations as in §6, one can define for any element  $b \in B(G)$  the twisted strata

$$\bar{S}'^{(b)} = \{x \in \bar{S}' \mid b(\mathcal{G}_{\bar{x}}) = b\} \subset \bar{S}'.$$

Then  $\bar{S}'^{(b)} \subset \bar{S}'$  is a reduced subscheme, and as the level  $K^p = K_q \times K^{pq}$  varies, these twisted strata are preserved by the natural projection  $\bar{S}'_{K_1^p} \rightarrow \bar{S}'_{K^p}$  (for  $K_1^p = K_q \times K_1^{pq} \subset K^p = K_q \times K^{pq}$ ) and also by the morphism  $g : \bar{S}'_{K^p} \rightarrow \bar{S}'_{gK^p g^{-1}}$  for all  $g \in G(\mathbb{A}_f^{pq})$ .

If  $\Sigma/\bar{\mathbb{F}}_p \supset k$  is a polarized Barsotti-Tate group endowed with a compatible action of  $\mathcal{O}_{B_{\mathbb{Q}_p}}$ , and  $b \in B(G)$  is its isogeny class, we define the twisted leaf

$$C'_b = C'_\Sigma = \{x \in \bar{S}' \mid \mathcal{G}_{\bar{x}} \simeq \Sigma_{\overline{k(x)}}\} \subset \bar{S}'^{(b)} \times_k \bar{\mathbb{F}}_p.$$

This is a closed subset of the stratum  $\bar{S}'^{(b)}$ , and as a subscheme of  $\bar{S}'^{(b)} \times_k \bar{\mathbb{F}}_p$  endowed with the induced reduced structure, it is smooth. As the level  $K^p = K_q \times K^{pq}$  varies, these twisted leaves are preserved by the natural projection  $\bar{S}'_{K_1^p} \rightarrow \bar{S}'_{K^p}$  (for  $K_1^p = K_q \times K_1^{pq} \subset K^p = K_q \times K^{pq}$ ) and also by the morphism  $g : \bar{S}'_{K^p} \rightarrow \bar{S}'_{gK^p g^{-1}}$  for all  $g \in G(\mathbb{A}_f^{pq})$ .

In the same way, one can define the twisted Igusa varieties  $J'_{b,m} \rightarrow C'_b$ , where a point of  $J'_{b,m}$  is represented by the data  $(A, \lambda, i, \bar{\eta}'^p; j_{m,1}, \dots, j_{m,k})$ . As in §7, we have that the group  $\Gamma_b$  acts on  $J'_{b,m}$ , and this action factors through  $\Gamma_{b,m}$ . From proposition 7.2, we know that the map  $J'_{b,m} \rightarrow C'_b$  is finite étale and Galois with Galois group  $\Gamma_{b,m}$ , and the twisted Igusa varieties  $J'_{b,m}$  are smooth.

As in §8, we have natural projections

$$q'_b = q_{b,m',m} : J'_{b,m'} \rightarrow J'_{b,m}, \quad \forall m' \geq m,$$

which correspond to restricting the isomorphisms  $j_{m',i}$  to the  $p^m$ -torsion subgroups for all  $i$ . The projections  $q'_b$  are invariant under the action of  $\Gamma_b$  on the Igusa varieties.

Thus, as the level  $m$  varies, the twisted Igusa varieties  $J'_{b,m}$  form a projective system under the morphism  $q_b$  which is endowed with an action of  $\Gamma_b$ . As in §7, this action naturally extends to the submonoid  $S_b \supset \Gamma_b$  of the group  $T_b$  of the quasi-self-isogenies of  $\Sigma$ , preserving its additional structures.

By reintroducing in our notation the level  $K^p = K_q \times K^{pq}$ , we write

$$J'_{b,K^p,m} \rightarrow C'_{b,K^p} \subset \bar{S}'_{K^p} \subset \bar{S}'_{K^p}$$

for the twisted Igusa varieties of level  $K^p = K_q \times K^{pq}$  and level  $m$  at  $p$ . For any  $b \in B(G)$  and  $m \geq 1$ , we define an action of the group  $G(\mathbb{A}_f^{pq})$  on the twisted Igusa varieties of level  $m$  at  $p$  (as the level  $K^p = K_q \times K^{pq}$  away from  $p$  varies) as

$$\begin{aligned} \forall g \in G(\mathbb{A}_f^{pq}) : J'_{b,K^p,m} &\rightarrow J'_{b,g^{-1}K^p g,m}, \\ (A, \lambda, i, \eta^{\bar{p}}; j_{m,1}, \dots, j_{m,k}) &\mapsto (A, \lambda, i, \overline{\eta^{\bar{p}} \circ g}; j_{m,1}, \dots, j_{m,k}). \end{aligned}$$

This action is clearly compatible under the projection map with the action of  $G(\mathbb{A}_f^{pq})$  on the twisted leaves  $C'_{b,K^p}$ , commutes with the natural projections among the Igusa varieties as the level at  $p$  varies, and also commutes with the action of  $S_b$ .

Thus, for  $l \nmid pq$  be a prime number, the direct limit of cohomologies with compact support

$$H_c^i(J'_{b,K_q}, \mathbb{Q}_l) = \lim_{\rightarrow K^p = K_q \times K^{pq,m}} H_c^i(J'_{b,K^p,m}, \mathbb{Q}_l), \quad \forall i,$$

is endowed with an action of the submonoid  $S_b \times G(\mathbb{A}_f^{pq}) \subset T_b \times G(\mathbb{A}_f^{pq})$ . As in §7, this action cannot be extended to an action of  $T_b \times G(\mathbb{A}_f^{pq})$  on the varieties, but it is possible to extend it on the cohomology groups.

We have the identities

$$H_c^i(J'_{b,K_q}, \mathbb{Q}_l)^{\Gamma_{b,m} \times K^{pq}} = H_c^i(J'_{b,K^p,m}, \mathbb{Q}_l),$$

where  $K^p = K_q \times K^{pq}$ .

## 9 The geometry of the twisted Newton polygon strata

In this section, following closely §5 of [M], we construct a system of finite morphisms

$$\pi'_N : J'_{m,b} \times \bar{\mathcal{M}}_b^{n,d} \rightarrow \bar{S}'^{(b)} \times_k \bar{\mathbb{F}}_p,$$

indexed by positive integers  $m, n, d, N$  ( $m \geq d$  and  $N \geq d/\delta f$ , where  $\delta \in \mathbb{Q}$  depends only on  $b \in B(G)$ ), which are compatible under the projections among the

twisted Igusa varieties and the inclusion among the twisted truncated Rapoport-Zink spaces  $\bar{\mathcal{M}}'^{n,d}$ , and which are also  $T_b$ -invariant and  $W_{E_v}$ -invariant, where  $W_{E_v}$  is the Weil group. Also for any geometric closed point  $x \in \bar{S}^{(b)}$ , the fibers of  $\pi'_N$  at  $x$  form a  $T_b$ -principal homogeneous space.

Now we recall the definition of the Rapoport-Zink space  $\mathcal{M}_{b/W(\bar{\mathbb{F}}_p)}$ . An isogeny class  $b \in B(G)$  (corresponding to  $S_{K^p}$ ) is called *descent* if it contains a Barsotti-Tate group defined over a finite field. In [OZ], it was proved that this is equivalent to containing a completely slope-divisible Barsotti-Tate group. We choose a completely slope-divisible Barsotti-Tate group  $\Sigma = \Sigma_b$  in the isogeny class  $b$ .

To any descent class  $b \in B(G)$ , Rapoport-Zink associated a formal scheme  $\mathcal{M}_b$  over  $W(\bar{\mathbb{F}}_p) \supset O_{E_v}$  as follows: for any scheme  $S/W(\bar{\mathbb{F}}_p)$ , where  $p$  is locally nilpotent,  $\mathcal{M}_b(S)$  is the set of isomorphism classes of Barsotti-Tate groups with additional structures  $H/S$  together with a quasi-isogeny  $\beta : \Sigma \times_{\bar{\mathbb{F}}_p} \bar{S} \rightarrow H \times_S \bar{S}$  defined over  $\bar{S} = Z(p) \subset S$ .

The group  $T_b$  acts on the Rapoport-Zink space  $\mathcal{M}_b$  by

$$\forall \rho \in T_b : (H, \beta) \mapsto (H, \beta \circ \rho).$$

One can define also a  $\sigma$ -semilinear isomorphism of  $\mathcal{M}_b$  arising from the Frobenius of  $\Sigma$ ,

$$\begin{aligned} \text{Frob} : \mathcal{M}_b &\rightarrow \mathcal{M}_b, \\ (H, \beta) &\mapsto (H, F^{-1} \circ \beta), \end{aligned}$$

which obviously commutes with the action of  $T_b$ .

Because the formal scheme  $\mathcal{M}_b$  is only locally of finite type, it is convenient to define certain subspaces of  $\mathcal{M}_b$ , which are called *truncated Rapoport-Zink spaces*, which are of finite type and which form a cover of  $\mathcal{M}_b$ . If  $n, d$  are positive integers, define  $\mathcal{M}_b^{n,d} \subset \mathcal{M}_b$  as the locus classifying pairs  $(H, \beta)$  such that the quasi-isogenies  $p^n \beta$  and  $p^{d-n} \beta^{-1}$  are isogenies (or, equivalently, such that  $p^n \beta$  is an isogeny whose kernel is killed by  $p^d$ ). Then  $\mathcal{M}_b^{n,d}$  are closed subspaces of  $\mathcal{M}_b$  and, as  $n, d$  vary, they form a direct system under the natural inclusions  $i_b = i_{n',d'}$  with  $\mathcal{M}_b$  as a limit. The truncated Rapoport-Zink spaces  $\mathcal{M}_b^{n,d}$  are not preserved by the actions of  $T_b$  and Frob, but one can think of the action of  $T_b$  and Frob on  $\mathcal{M}$  as arising from the action on the direct system. More exactly, if  $\rho \in T_b$ , let  $n(\rho)$  and  $d(\rho)$  be the smallest integer such that  $p^{n(\rho)} \rho$  and  $p^{d(\rho)-n(\rho)} \rho^{-1}$  are isogenies. Then the morphism  $\rho : \mathcal{M}_b \rightarrow \mathcal{M}_b$  induces by restriction some morphisms

$$\rho : \mathcal{M}_b^{n,d} \rightarrow \mathcal{M}_b^{n+n(\rho), d+d(\rho)}, \quad \forall n, d,$$

which obviously commute with the natural inclusions. Also by restricting Frob to  $\mathcal{M}_b^{n,d} \subset \mathcal{M}_b$ , we obtain a  $\sigma$ -linear morphism

$$\text{Frob} : \mathcal{M}_b^{n,d} \rightarrow \mathcal{M}_b^{n+1, d+1}, \quad \forall n, d.$$

We denote by  $\bar{\mathcal{M}}_b$  (resp.  $\bar{\mathcal{M}}_b^{n,d}$ ) over  $\bar{\mathbb{F}}_b$ , the reduced fiber of  $\mathcal{M}_b$  (resp.,  $\mathcal{M}_b^{n,d}$  for any  $n, d$ ). Then  $\bar{\mathcal{M}}_b$  uniformizes the isogeny classes of the abelian varieties classified by  $\bar{S}^{(b)}$ .

In the same way, starting with an isogeny class  $b \in B(G)$  corresponding to  $S'_{K^p}$ , one can define the twisted Rapoport-Zink space  $\mathcal{M}'_b$  and the truncated Rapoport-Zink spaces  $\mathcal{M}'_b^{n,d}$ , verifying the same proprieties as above.

We write  $\text{Fr} = \text{Fr}_{\bar{S}'_{K^p}}$  for the absolute Frobenius. Then  $\text{Fr}^f$  is the morphism  $\bar{S}'_{K^p} \rightarrow \bar{S}'_{K^p}$  which sends the quadruple  $(A, \lambda, i, \bar{\eta}^p)$  to  $(A^{(q)}, \lambda^{(q)}, i^{(q)}, \bar{\eta}^{p(q)})$ .

From proposition 9 of [M], and the description of  $S'$  in §5, and of  $J'_{m,b}$  in §7, we obtain the following result:

**Proposition 9.1.** *For any positive integers  $m, n, d, N, m \geq d$  and  $N \geq d/\delta f$ , there exists some morphisms*

$$\pi'_N : J'_{m,b} \times \bar{\mathcal{M}}_b^{n,d} \rightarrow \bar{S}'^{(b)} \times_k \bar{\mathbb{F}}_p,$$

such that

- (1)  $\pi'_{N+1} = (\text{Fr}^f \times 1) \circ \pi'_N$ ;
- (2)  $\pi'_N \circ q_b = \pi'_N$ ;
- (3)  $\pi'_N \circ i_b = \pi'_N$ ;
- (4)  $\pi'_N \circ \rho \times \rho = \pi'_N$  for any  $\rho \in S_b$ , provided that  $m \geq d + d(\rho) + e(\rho)$  and  $N \geq (d + d(\rho))/\delta f$ ;
- (5)  $\pi'_N \circ (\text{Frob}^f \times \text{Frob}^f) = (1 \times \sigma^f) \circ \pi'_N$  for all  $m \geq d+1$  and  $N \geq (d+1)/\delta f$ .

We now describe the fibers on the morphisms  $\pi'_N$ . If  $x$  is a point of  $\bar{S}'^{(b)}(\bar{\mathbb{F}}_p)$ , then for any  $N \geq N_0 \geq d/\delta f$ , we consider the fibers over  $x$  of the maps

$$\pi'_N : J'_{m,b} \times \bar{\mathcal{M}}_b^{n,d} \rightarrow \bar{S}'^{(b)} \times_k \bar{\mathbb{F}}_p.$$

Using proposition 9.1 (1), we can identify

$$\pi'^{-1}_N(\text{Fr}^{fN} x) = \pi'^{-1}_{N_0}(\text{Fr}^{fN_0} x).$$

As  $m$  varies, the sets  $\pi'^{-1}_N(\text{Fr}^{fN} x)$  form an inverse system under the projections  $q_b$ , and the corresponding limits are a direct system under the inclusions  $i_b$  as  $n, d$  vary. We define the *fiber above  $x$*  to be the set

$$\Pi'^{-1}(x) = \lim_{\rightarrow n, d} \lim_{\leftarrow m} \pi'^{-1}_N(\text{Fr}^{fN} x),$$

which is endowed with the topology of the direct limit of inverse limits of discrete sets. Then from proposition 9.1 (4), we get that  $\Pi'^{-1}(x)$  is also endowed with continuous action of  $S_b$ , and because of proposition 9.1 (5) and lemma 7.4, this action extends uniquely to a continuous action of the group  $T_b = \langle S_b, p, \text{fr}^B \rangle$ .

We obtain (see propositions 10 and 11 of [M]):

**Proposition 9.2.** *For any  $x \in \bar{S}'^{(b)}(\bar{\mathbb{F}}_p)$ , its fiber  $\Pi'^{-1}(x)$  is a principle homogeneous space for the continuous action of  $T_b$ .*

**Proposition 9.3.** *For any positive integers  $m, n, d, N$ ,  $m \geq d$  and  $N \geq d/\delta f$ , the morphism  $\pi'_N : J'_{m,b} \times \bar{\mathcal{M}}'^{n,d}_b \rightarrow \bar{S}'^{(b)} \times \bar{\mathbb{F}}_p$  is finite.*

Let  $U'^{n,d}_b$  be the maximal open subspaces of  $\mathcal{M}'_b$  contained in  $\bar{\mathcal{M}}'^{n,d}_b$ , and let  $\bar{U}'^{n,d}_b$  be their reduced fibers. Then the sets  $U'^{n,d}_b$  are an open cover of  $\bar{\mathcal{M}}'$ , and thus for any sheaf  $\mathcal{F}/\bar{\mathcal{M}}'_b$ , we have

$$H_c^t(\bar{\mathcal{M}}'_b, \mathcal{F}) = \lim_{\rightarrow n,d} H_c^t(\bar{U}'^{n,d}_b, \mathcal{F}), \quad t \geq 0.$$

Let  $\pi'_{N,m}$  be the restrictions of the morphisms  $\pi'_N$  to the subschemes  $J'_{b,m} \times \bar{U}'^{n,d}_b \subset J'_{b,m} \times \bar{\mathcal{M}}'^{n,d}_b$  and let  $\bar{\pi}r : J'_{b,m} \times \bar{U}'^{n,d}_b \rightarrow \bar{\mathcal{M}}'_b$  be the projections to the second factor.

We denote by  $\tau \mapsto \bar{\tau} = (\sigma^f)^{r(\tau)}$  the projection  $W_{E_v} \rightarrow \text{Gal}(\bar{\mathbb{F}}_p/k)$ . If  $\mathcal{L}$  is a torsion étale over  $\bar{S}'^{(b)} \times \bar{\mathbb{F}}_p$ , we define an action of the Weil group  $W_{E_v}$  on  $\mathcal{L}$  to be the data of isomorphisms

$$\tau : (1 \times \bar{\tau})^* \mathcal{L} \simeq \mathcal{L}, \quad \forall \tau \in W_{E_v},$$

such that  $\tau \circ \tau' = \tau' \tau$ . Also, we call an *action of the Weil group  $W_{E_v}$  on a sheaf  $\mathcal{G}$  over  $\bar{\mathcal{M}}'$*  the data of isomorphisms  $\tau : (\text{Frob}^f)^{r(\tau)} \mathcal{G} \simeq \mathcal{G}$  for all  $\tau \in W_{E_v}$ ,  $r(\tau) \geq 0$ , such that  $\tau \circ \tau' = \tau' \tau$ .

We obtain (see theorems 12 and 13 and corollary 14 of [M]):

**Theorem 9.4.** *For any  $b \in B(G)$  and  $K^p = K_q \times K^{p,q} \subset G(\mathbb{A}_f^p)$ , let  $\mathcal{L}$  be an abelian torsion étale sheaf over  $\bar{S}'^{(b)}$  with torsion orders prime to  $p$  and endowed with an action of the Weil group  $W_{E_v}$ . Then there exists a  $W_{E_v}$ -equivariant spectral sequence*

$$E_2^{p,q} = H^q(T_b, \lim_{\rightarrow m,n,d} H_c(J'_{b,m} \times \bar{U}'^{n,d}_b, \pi'^*_N \mathcal{L})) \Rightarrow H_c^{p+q}(\bar{S}'^{(b)}_{K^p} \times \bar{\mathbb{F}}_p, \mathcal{L}).$$

For  $l \neq p$  a prime and an integer  $r \geq 1$ , we denote by  $\mathcal{H}_r(T_b)$  the Hecke algebra of  $T_b$  with coefficients in  $\mathbb{Z}/l^r \mathbb{Z}$ .

**Theorem 9.5.** *For any  $b \in B(G)$  and  $K^p = K_q \times K^{p,q} \subset G(\mathbb{A}_f^p)$ , let  $\mathcal{L}$  (resp.  $\mathcal{G}$ ) be an étale sheaf of  $\mathbb{Z}/p\mathbb{Z}$ -modules over  $\bar{S}'^{(b)} \times \bar{\mathbb{F}}_p$  (resp.  $\bar{\mathcal{M}}'$ ) which is endowed with an action of the Weil group  $W_{E_v}$ .*

*Suppose there exists a system of  $W_{E_v}$ -equivariant isomorphisms of étale sheaves*

$$\{\pi'^*_N \mathcal{L} \simeq \bar{\pi}r^* \mathcal{G}\}_{m,n,d,N,m \geq d, N \geq d/\delta f}$$

*which are compatible under the natural transaction maps induced by  $q_b, i_b$  and  $\rho \in S_b$ .*

*Then there exists a  $W_{E_v}$ -equivariant spectral sequence of  $\mathbb{Z}/l^r \mathbb{Z}$ -modules:*

$$\oplus_{t+s=q} \text{Tor}_{\mathcal{H}(T_b)}^p(H_c^s(\bar{\mathcal{M}}', \mathcal{G}), \lim_{\rightarrow m} H_c^t(J'_{b,K^p,m}, \mathbb{Z}/l^r \mathbb{Z})) \Rightarrow H_c^{p+q}(\bar{S}'^{(b)}_{K^p} \times \bar{\mathbb{F}}_p, \mathcal{L}).$$

If  $\mathcal{L}$  (resp.  $\mathcal{G}$ ) is the constant étale sheaf over  $\bar{S}'^{(b)}$  (resp., over  $\bar{\mathcal{M}}'$ ) associated to  $\mathbb{Z}/l^r\mathbb{Z}$ , then the hypothesis of theorem 9.5 is satisfied and thus we get:

**Corollary 9.6.** *For any  $b \in B(G)$  and  $K^p = K_q \times K^{pq} \subset G(\mathbb{A}_f^p)$ , there exists a  $W_{E_v}$ -equivariant spectral sequence of  $\mathbb{Z}/l^r\mathbb{Z}$ -modules:*

$$\oplus_{t+s=q} \text{Tor}_{\mathcal{H}(T_b)}(H_c^s(\bar{\mathcal{M}}', \mathbb{Z}/l^r\mathbb{Z}), H_c^t(J'_{b,K^p}, \mathbb{Z}/l^r\mathbb{Z})) \Rightarrow H_c^{p+q}(\bar{S}'_{K^p}^{(b)} \times \bar{\mathbb{F}}_p, \mathbb{Z}/l^r\mathbb{Z}),$$

where we write  $H_c^t(J'_{b,K^p}, \mathbb{Z}/l^r\mathbb{Z}) = \lim_{\rightarrow m} H_c^t(J'_{b,K^p,m}, \mathbb{Z}/l^r\mathbb{Z})$  for all  $t \geq 0$ .

We remark that all the above results are proved for the varieties  $S_{K^p} = S_{K^p(0)}/O_{E_v}$ , corresponding to  $K^p(0) = K^p K_p$ , where  $K_p$  is a hyperspecial maximal compact of  $G(\mathbb{Q}_p)$  (more exactly  $K_p$  is the stabilizer of the lattice  $\Lambda$ ). For any  $m \geq 0$ , one can define (see §6 of [M]) also the varieties  $S_{K^p(m)} \rightarrow S_{K^p(0)}$ , where  $K^p(m) = K^p K_m$ , and  $K_m \subset K_p$  is a congruence subgroup of level  $m$  (i.e.  $K_m$  is the stabilizer of the lattice  $\Lambda$  modulo  $p^m$ , see [M] for details). Also for any  $m \geq 0$ , one can define the Rapoport-Zink spaces of level  $m$ ,  $\mathcal{M}_{b,m} \rightarrow \mathcal{M}_b = \mathcal{M}_{b,0}$ . Moreover, this construction could be extended (see §6 of [M]), and for any

$$g \in G(\mathbb{Q}_p)^+ = \{g \in G(\mathbb{Q}_p) | g^{-1}(\Lambda) \subset \Lambda\} \subset G(\mathbb{Q}_p),$$

and for any  $m \geq e(g)$ , where  $e(g)$  is some number which depends on  $g$ , one can define the varieties  $S_{K^p(m),g} \rightarrow S_{K^p(m)}$  and the Rapoport-Zink spaces  $\mathcal{M}_{b,m,g} \rightarrow \mathcal{M}_{b,m}$  (for  $g \in K_p$ , we have  $S_{K^p(m),g} = S_{K^p(m)}$  and  $\mathcal{M}_{b,m,g} = \mathcal{M}_{b,m}$ ).

As above for  $K^p$  of the form  $K^p = K_q \times K^{pq}$ , one can define the twisted varieties  $S'_{K^p(m),g}$ , and the twisted Rapoport-Zink spaces  $\mathcal{M}'_{b,m,g}$ , and following §6, 7 and 8 of [M], we obtain (see theorem 22 of [M]):

**Theorem 9.7.** *There is an equality of virtual representations of the group  $G(\mathbb{A}_f^q) \times 1 \times W_{E_v}$ ,*

$$\begin{aligned} & \sum_{t \geq 0} (-1)^t \lim_{\rightarrow K^q \subset G(\mathbb{A}_f^q)} H^t(S'_{K^q \times K^q \times E_v} \bar{E}_v, \mathbb{Q}_l) \\ &= \sum_{b \in B(G)} \sum_{i,j,k \geq 0} (-1)^{i+j+k} \lim_{\rightarrow U \subset G(\mathbb{Q}_p)} \mathcal{E}_{b,U}^{i,j,k}, \end{aligned}$$

where for all  $b \in B(G)$  and  $i, j, k \geq 0$ , we write

$$\mathcal{E}_{b,U}^{i,j,k} = \text{Ext}_{T_b\text{-smooth}}^i(H_c^j(\mathcal{M}'_{b,U}{}^{\text{rig}} \times_{E_v^{\text{nr}}} \bar{E}_v, \mathbb{Q}_l(-D)), H_c^k(J'_{b,K^q}, \mathbb{Q}_l)),$$

where  $D$  is the dimension of the Shimura varieties.

We remark that from §4, we know that variety  $S_{K^p}$  is a disjoint union of  $|\ker^1(\mathbb{Q}, G)|$ -copies of  $\mathbf{S}_{\mathbf{K}}$ . Also we have that the variety  $J_{b,K^p,m}$  is a disjoint union of  $|\ker^1(\mathbb{Q}, G)|$ -copies of the corresponding varieties  $\mathbf{J}_{b,\mathbf{K},m}$ , and thus all the above results remain true if we replace  $S'_{K^p}$  by  $\mathbf{S}'_{\mathbf{K}}$  and  $J'_{b,K^p,m}$  by  $\mathbf{J}'_{b,\mathbf{K},m}$ , etc.

## References

- [CA] H. Carayol, *Sur la mauvaise réduction des courbes de Shimura*, Compositio Mathematica, 59, nr.2 (1986), 151-230.
- [D] P. Deligne, *Travaux de Shimura*, Sémin. Bourbaki Février 1971, Exposé. No. 389, Lecture Notes in Math., vol. 244, Springer, Heidelberg, 1971.
- [K1] R. Kottwitz, *Shimura varieties and  $\lambda$ -adic representations*, Automorphic Forms, Shimura varieties and L-functions, part 1, Perspectives in Mathematics, vol. 10, Academic Press, San Diego, CA, 1990, pp. 161-209.
- [K2] R. Kottwitz, *Points on some Shimura varieties over finite fields*, Journal of American Mathematical Society, Vol. 5, No. 2. (Apr., 1992), pp. 373-444.
- [M] E.Mantovan, *On the cohomology of certain PEL-type Shimura varieties*. Duke Math. J., vol. 129, no.3, 2005.
- [M1] E.Mantovan, *On certain unitary group Shimura varieties* in Varieties de Shimura, espaces de Rapoport-Zink et correspondences de Langlands locales, Asterisque 291, Soc. Math. France, Montrouge, 2004, 200-331.
- [O] F.Oort, *Foliations in moduli spaces of abelian varieties*, J. Amer. Math. Soc. 17 (2004), 267-296.
- [OZ] F.Oort, T.Zink, *Families of  $p$ -divisible groups with constant Newton polygon*, Doc. Math. 7 (2002), 183-201.
- [RR] M.Rapoport, M.Richardz, *On the classification and specialization of  $F$ -isocrystals with additional structure*, Compositio Math. 103 (1996), 153-181.
- [RZ] M.Rapoport, T.Zink, *Period spaces for  $p$ -divisible Groups*, Ann. of Math. Stud. 141, Princeton Univ. Press, Princeton, 1996.
- [Z] T.Zink, *On the slope filtration*, Duke Math. J. 109 (2001), 79-95.