

# Non-solvable base change for Hilbert modular representations and zeta functions of twisted quaternionic Shimura varieties

Cristian Virdol  
Department of Mathematics  
Columbia University

September 16, 2010

Résumé: Dans cet article, nous montrons un changement de base non-résoluble pour certaines représentations modulaires de Hilbert et nous utilisons ce résultat pour établir le prolongement méromorphe à tout le plan complexe des fonctions zêta de certaines variétés de Shimura quaternioniques tordus. Les fonctions zêta des variétés de Shimura quaternioniques tordus sont calculées à toutes les places.

## Abstract

In this paper we prove some non-solvable base change for Hilbert modular representations, and we use this result to show the meromorphic continuation to the entire complex plane of the zeta functions of some twisted quaternionic Shimura varieties. The zeta functions of the twisted quaternionic Shimura varieties are computed at all places.

2000 Mathematics subject classification: 11F41, 11F80, 11R42, 11R80.

## 1 Introduction

In the first part of this article we prove the following non-solvable base change for Hilbert modular representations:

**Theorem 1.1.** *Let  $F$  be a totally real number field, and let  $\pi$  be a cuspidal automorphic representation of weight  $k \geq 2$  of  $GL(2)/F$ . Let  $F'$  be a finite solvable extension of a totally real number field containing  $F$ . Then there exists a number field  $F''$  containing  $F'$  which is a solvable extension of a totally real field, such that  $F''$  is Galois over  $\mathbb{Q}$ , and such that the representation  $\pi$  admits a base change to  $GL(2)/F''$ . If  $F'$  is a totally real number field, then  $F''$  can be chosen to be a totally real number field.*

To show this theorem, we use some results from Taylor's papers [HSBT] and [T2]. We recall that from Langlands [L] and Arthur-Clozel [AC], we know that if  $\pi$  is an automorphic representation of  $\mathrm{GL}(n)/F$ , where  $F$  is a number field, and  $F'$  is a solvable extension of  $F$ , then  $\pi$  admits a base change to  $\mathrm{GL}(n)/F'$ .

In the second part of this article, we compute the zeta function of some "twisted" quaternionic Shimura varieties in terms of automorphic representations, and as an application of Theorem 1.1 we prove that their zeta function could be meromorphically continued to the entire complex plane and satisfies a functional equation. In [BL], Brylinski-Labesse computed the zeta function of quaternionic Shimura varieties associated to a totally indefinite quaternion algebra  $D$  over a totally real field  $F$  i.e. all the infinite places of  $F$  are unramified in  $D$ . In his book [R], Reimann generalized the result in [BL] and computed the semisimple zeta function of quaternionic Shimura varieties associated to indefinite quaternion algebras  $D$ . Then in [B], Blasius generalized the result in [R] and obtained the expression of the zeta function of quaternionic Shimura varieties at all places.

More exactly, in the second part of this article, we consider  $F$  a totally real field,  $O := O_F$  the ring of integers of  $F$  and  $D$  an indefinite quaternion algebra over  $F$ . Let  $G$  be the algebraic group over  $F$  defined by the multiplicative group  $D^\times$  of  $D$  and let  $\bar{G} := \mathrm{Res}_{F/\mathbb{Q}}(G)$ . We fix a prime ideal  $\wp$  of  $O_F$ , such that  $G(F_\wp)$  is isomorphic to  $\mathrm{GL}_2(F_\wp)$ . Let  $S_{\bar{G}, \mathbf{K}} = S_{\mathbf{K}}$  be the canonical model of the quaternionic Shimura variety associated to an open compact subgroup  $\mathbf{K} := K_\wp \times H$  of  $\bar{G}(\mathbb{A}_f)$ , where  $K_\wp$  is the set of elements of  $\mathrm{GL}_2(O_\wp)$  that are congruent to 1 modulo  $\wp$ ,  $H$  is an open compact subgroup of the restricted product of  $(D \otimes_F F_{\mathfrak{p}})^\times$  where  $\mathfrak{p}$  runs over all the finite places of  $F$ ,  $\mathfrak{p} \neq \wp$  and  $\mathbb{A}_f$  is the finite part of the ring of adèles  $\mathbb{A}_{\mathbb{Q}}$  of  $\mathbb{Q}$ . Then  $S_{\mathbf{K}}$  is a quasi-projective variety defined over a totally real number field  $E$  called the canonical field of definition.

The variety  $S_{\mathbf{K}}$  has a natural action of  $\mathrm{GL}_2(O/\wp)$  (see §3.2). For  $H$  sufficiently small this action is free. We fix such a small group  $H$ . If  $K$  is a number field, we denote  $\Gamma_K := \mathrm{Gal}(\bar{\mathbb{Q}}/K)$ . Consider a continuous Galois representation  $\varphi : \Gamma_E \rightarrow \mathrm{GL}_2(O/\wp)$  and let  $S'_{\mathbf{K}}$  be the variety defined over  $E$  obtained from  $S_{\mathbf{K}}$  via twisting by  $\varphi$  composed with the natural action of  $\mathrm{GL}_2(O/\wp)$  on  $S_{\mathbf{K}}$  (see §3.2 for details).

From Corollary 11.8 of [R] and Theorem 3 of [B] (see also Propositions 3.3 and 3.4 below), we know that the zeta function  $L(s, S_{\mathbf{K}})$  of  $S_{\mathbf{K}}$  is given by the formula (see §3.1 for notations):

$$L(s, S_{\mathbf{K}}) = \prod_{\pi} L(s - d'/2, \pi, r)^{m(\pi_\infty)m(\pi_f^{\mathbf{K}})}.$$

Here the product is taken over automorphic cohomological representations  $\pi$  of  $\bar{G}(\mathbb{A}_{\mathbb{Q}})$  of weight 2,  $d'$  is the dimension of  $S_{\mathbf{K}}$ ,  $m(\pi_f^{\mathbf{K}})$  is the dimension of  $\pi_f^{\mathbf{K}}$  ( $\pi_f^{\mathbf{K}}$  denotes the subspace of  $\mathbf{K}$ -invariants of  $\pi_f$ ),  $r$  is a well specified representation of the  $L$ -group  ${}^L\bar{G}|_{\Gamma_E}$  associated to  $\bar{G}$  (see §3.1 for the definition of  $r$ ) and  $m(\pi_\infty)$  will be defined in §3.1.

In this article we obtain the following result (see §3.1 for notations and also Proposition 3.5):

**Theorem 1.2.** *The zeta function  $L(s, S'_{\mathbf{K}})$  of  $S'_{\mathbf{K}}$  is given by the formula:*

$$L(s, S'_{\mathbf{K}}) = \prod_{\pi} L(s - d'/2, \pi, r \otimes (\pi_f^{\mathbf{K}} \circ \varphi))^{m(\pi_{\infty})},$$

where the product is taken over automorphic cohomological representations  $\pi$  of  $\bar{G}(\mathbb{A}_{\mathbb{Q}})$  of weight 2, such that  $\pi_f^{\mathbf{K}} \neq 0$ .

If  $d' = 1$  or  $2$  and the field  $L := \bar{\mathbb{Q}}^{\text{Ker}(\varphi)}$  is a solvable extension of a totally real field, then the zeta function  $L(s, S'_{\mathbf{K}})$  can be meromorphically continued to the entire complex plane and satisfies a functional equation.

The first part of Theorem 1.2 is proved in §3.3 by taking the injective limit of the representations of  $\Gamma_E \times \mathbb{H}_K$  on the étale cohomology of Shimura varieties  $S_K$  and using some linear algebra. Here  $K$  is an open compact subgroup of  $\bar{G}(\mathbb{A}_f)$  and  $\mathbb{H}_K$  is the Hecke algebra of level  $K$ .

The second part of Theorem 1.2 regarding the meromorphic continuation of the zeta function  $L(s, S'_{\mathbf{K}})$  is proved in §4. We show using Theorem 1.1 (see Theorem 4.3) that if  $d' = 1$  or  $2$  and  $\pi$  is a representation as in the product of Theorem 1.2 and  $\omega$  is an Artin representation of  $\Gamma_E$  such that the field  $K := \bar{\mathbb{Q}}^{\text{Ker}(\omega)}$  is a solvable extension of a totally real field, then the  $L$ -function  $L(s, \pi, r \otimes \omega)$  can be meromorphically continued to the entire complex plane and satisfies a functional equation. We prove also the meromorphic continuation and functional equation of  $L(s, S'_{\mathbf{K}})$  when  $d' \geq 3$  and the field  $L := \bar{\mathbb{Q}}^{\text{Ker}(\varphi)}$  is a solvable extension of a totally real field, if we assume that some other Langlands  $L$ -functions can be meromorphically continued to the entire complex plane and satisfy a functional equation (see Lemma 4.1).

We remark that when  $D = M_2(F)$  the Shimura variety is not compact and in this case we use the  $l$ -adic intersection cohomologies of the Baily-Borel compactification of the Shimura variety.

In this article, if  $\pi$  is an automorphic representation of  $\bar{G}(\mathbb{A}_{\mathbb{Q}})$ , we denote the automorphic representation of  $\text{GL}_2(\mathbb{A}_F)$  ( $\mathbb{A}_F$  is the ring of adèles of  $F$ ), obtained from  $\pi$  by Jacquet-Langlands correspondence (usually denoted  $JL(\pi)$ ) by the same symbol  $\pi$ .

## 2 The proof of Theorem 1.1

Let  $\pi$  be a cuspidal automorphic representation of weight  $k \geq 2$  of  $\text{GL}(2)/F$ , where  $F$  is a totally real field. Then from [T1], we know that there exists a  $\lambda$ -adic representation (for  $\lambda$  a prime of the ring of coefficients  $\mathbf{O}$  of  $\pi$ , such that  $\lambda \nmid l$  for some rational prime  $l$ )

$$\rho_{\pi} := \rho_{\pi, \lambda} : \Gamma_F \rightarrow \text{GL}_2(\mathbf{O}_{\lambda}) \hookrightarrow \text{GL}_2(\bar{\mathbb{Q}}_l),$$

that is unramified outside the primes dividing  $\mathbf{n}l$ , where  $\mathbf{n}$  is the level of  $\pi$ . We denote by  $\bar{\rho}_{\pi} = \bar{\rho}_{\pi, \lambda}$  the reduction of  $\rho_{\pi} = \rho_{\pi, \lambda} : \Gamma_F \rightarrow \text{GL}_2(\mathbf{O}_{\lambda}) \pmod{\lambda}$ . An

$l$ -adic representation  $\rho : \Gamma_F \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_l)$  is called automorphic (or modular) if  $\rho \cong \rho_\pi$  for some  $\pi$  as above. Proving Theorem 1.1 is equivalent to proving the potential automorphy of the corresponding  $l$ -adic representations.

For  $F = \mathbb{Q}$  and  $k = 2$ , Theorem 1.1 is Theorem 3.7 of [V1]. The proof in [V1] uses the positivity of the density of the set of ordinary primes that is known for cuspidal automorphic representations of  $\mathrm{GL}(2)/\mathbb{Q}$ . This fact is not known for cuspidal automorphic representations of  $\mathrm{GL}(2)/F$  for general totally real field  $F$ . To prove Theorem 1.1 for general totally real field  $F$ , one uses some results from [HSBT] and [T2].

We say that the automorphic representation  $\pi$  of  $\mathrm{GL}(2)/F$ , where  $F$  is a totally real field, is of *CM-type* if there exists some Galois character  $\eta : I_F/F^\times \rightarrow \overline{\mathbb{Q}}_l^\times$ , where  $I_F$  denotes the idele group of  $F$ , with  $\eta \neq 1$  such that  $\pi \cong \pi \otimes \eta$ . It is known (see Theorem 7.11 of [G]) that if  $\pi$  is of CM-type, then  $\pi$  admits a base change to  $\mathrm{GL}(2)/L$  for any finite extension  $L/F$ .

So it is sufficient to prove Theorem 1.1 when the representation  $\pi$  are non-CM. We assume this fact from now on.

We assume first that the field  $F'$  from Theorem 1.1 is totally real and prove Theorem 1.1 in this case.

We know the following result (see Theorem 3.1 of [HSBT] (unitary case) and Theorem 3.3 of [T2] for details):

**Theorem 2.1.** *Suppose that  $F$  is a totally real number field and let  $k \geq 2$  be an integer. Suppose that  $l > \max\{3, k\}$  is a rational prime which is unramified in  $F$ . Suppose also that*

$$\rho : \Gamma_F \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_l)$$

*is a continuous odd representation (i.e.  $\det \rho(c) = -1$  for each complex conjugation  $c$ ) which is unramified at all but finitely many primes and satisfies the following proprieties:*

1. *The image of  $\bar{\rho}$  contains  $SL_2(\mathbb{F}_l)$ ;*
  2. *If  $w|l$  is a prime of  $F$ , then the representation  $\rho|_{D_w}$ , where  $D_w$  is the decomposition group at  $w$ , is crystalline with Hodge-Tate weights 0 and  $k - 1$ .*
- Then there exists a totally real finite extension  $M/F$  which is Galois over  $\mathbb{Q}$ , such that each  $\rho|_{\Gamma_M}$  is automorphic.*

We want to apply Theorem 2.1 to the  $l$ -adic representation  $\rho_\pi|_{\Gamma_{F'}}$  (see Theorem 1.1 and the beginning of §2 for notations). From the properties of  $\rho_\pi$  we have that  $\det \rho_\pi|_{\Gamma_{F'}}(c) = -1$  for each complex conjugation  $c$ .

Since the representation  $\pi$  is non-CM, we know the following result (see Proposition 3.8 of [D]):

**Proposition 2.2.** *For  $l$  sufficiently large, the image of the residual representation  $\bar{\rho}_\pi$  contains  $SL_2(\mathbb{F}_l)$ .*

Hence we can choose a sufficiently large  $l$  such that the images of all representations  $\bar{\rho}_\pi$  contain  $SL_2(\mathbb{F}_l)$ . Since  $F'$  is totally real, the image of  $\bar{\rho}_\pi|_{\Gamma_{F'}}$

contains  $\mathrm{SL}_2(\mathbb{F}_l)$  (see Proposition 3.5 of [V1]). Thus the condition 1 of Theorem 2.1 is satisfied.

We know the following result (see for example Corollary 2.10 of [D]):

**Proposition 2.3.** *Assume that  $\pi$  is a cuspidal automorphic representation of weight  $k \geq 2$  of  $\mathrm{GL}(2)/F$ , where  $F$  is a totally real field. Then if  $l$  is a sufficiently large rational prime, we have that for each prime  $v$  of  $F$  dividing  $l$ , the  $l$ -adic representation  $\rho_{\pi, \lambda}|_{D_v}$ , where  $\lambda|l$  is a prime of the field of coefficients of  $\pi$  and  $D_v$  is the decomposition group at  $v$ , is crystalline with Hodge-Tate weights 0 and  $k - 1$ .*

Hence for  $l$  sufficiently large the condition 2 of Theorem 2.1 is also satisfied and Theorem 1.1 is proved if  $F'$  is a totally real number field.

Now we prove Theorem 1.1 for  $F'$  a solvable extension of a totally real field  $F_0$  containing  $F$ . From Theorem 1.1 applied to the totally real extension  $F_0$  of  $F$ , we deduce that there exists a Galois totally real extension  $F''$  of  $F_0$  such that the representation  $\pi$  admits a base change to  $\mathrm{GL}(2)/F''$ . Then  $F'F''$  is a solvable extension of  $F''$ . Since  $F''$  is Galois over  $\mathbb{Q}$ , we deduce that the Galois closure  $F'''$  of  $F'F''$  over  $\mathbb{Q}$  is a solvable extension of  $F''$ . Hence from Langlands base change for solvable extensions we get that the representation  $\pi$  admits base changes to  $\mathrm{GL}(2)/F'''$  and thus Theorem 1.1 is proved. ■

### 3 Quaternionic Shimura varieties

Consider a totally real number field  $F$  of degree  $d$  over  $\mathbb{Q}$  and let  $D$  be a quaternion algebra over  $F$ . Let  $\mathbb{A}_f$  be the finite part of the ring of adèles  $\mathbb{A}_{\mathbb{Q}}$  of  $\mathbb{Q}$ . We denote by  $J_F$  the set of infinite places of  $F$  and we identify  $J_F$  as a  $\Gamma_{\mathbb{Q}}$ -set with  $\Gamma_F \setminus \Gamma_{\mathbb{Q}}$ . Let  $J'_F$  be the subset of places of  $J_F$  where  $D$  is ramified. Let  $d' :=$ the cardinal of  $J_F - J'_F$ . We assume that  $d' > 0$ , i.e.  $D$  is indefinite over  $F$ .

Let  $G$  be the algebraic group over  $F$  defined by the multiplicative group  $D^\times$  of  $D$ . Consider the algebraic group  $\bar{G} := \mathrm{Res}_{F/\mathbb{Q}}(G)$  over  $\mathbb{Q}$  defined by the propriety:  $\bar{G}(A) = G(A \otimes_{\mathbb{Q}} F)$  for all  $\mathbb{Q}$ -algebras  $A$ . The  $L$ -group associated to  $\bar{G}$  is defined by the semidirect product:

$${}^L\bar{G} := {}^L\bar{G}^0 \rtimes \Gamma_{\mathbb{Q}},$$

where  ${}^L\bar{G}^0$  is the product of  $d$  copies of  $\mathrm{GL}_2(\mathbb{C})$  indexed by elements  $\sigma \in \Gamma_F \setminus \Gamma_{\mathbb{Q}}$  and  $\Gamma_{\mathbb{Q}}$  acts on  ${}^L\bar{G}^0$  by permuting the factors in the natural way. It is easy to see that  $\bar{G}(\mathbb{R})$  is isomorphic to  $\mathrm{GL}_2(\mathbb{R})^{d'} \times \mathbf{H}^{\times(d-d')}$ , where  $\mathbf{H}$  is the algebra of quaternions over  $\mathbb{R}$ .

For  $v \in J_F - J'_F$ , we fix an isomorphism of  $G(F_v)$  with  $\mathrm{GL}_2(\mathbb{R})$ . We have  $\bar{G}(\mathbb{R}) = \prod_{v \in J_F} G(F_v)$ . Let  $J := (J_v) \in \bar{G}(\mathbb{R})$ , where

$$J_v := \begin{cases} 1 & \text{for } v \in J'_F; \\ 1/\sqrt{2} \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} & \text{for } v \in J_F - J'_F. \end{cases}$$

Let  $K_\infty$  be the centralizer of  $J$  in  $\bar{G}(\mathbb{R})$ . Put

$$X := \bar{G}(\mathbb{R})/K_\infty.$$

It is well known that  $X$  is complex analytically isomorphic to  $(\mathfrak{H}_\pm)^{d'}$ , where  $\mathfrak{H}_\pm = \mathbb{C} - \mathbb{R}$ . For each open compact subgroup  $K \subset \bar{G}(\mathbb{A}_f)$  put

$$S_K(\mathbb{C}) := \bar{G}(\mathbb{Q}) \backslash X \times \bar{G}(\mathbb{A}_f)/K.$$

For  $K$  sufficiently small,  $S_K(\mathbb{C})$  is a complex manifold which is the set of complex points of a quasi-projective variety. In general  $S_K(\mathbb{C})$  is not connected and is a finite disjoint union of quotients  $\Gamma \backslash \mathfrak{H}_\pm^{d'}$ , where  $\Gamma \subset \bar{G}(\mathbb{Q})$  is a congruence subgroup. The subfield  $E$  of  $\mathbb{Q}$  having the propriety that  $\Gamma_E$  is the stabilizer of the subset  $J'_F \subset \Gamma_F \backslash \Gamma_\mathbb{Q}$ , for the natural right action of  $\Gamma_\mathbb{Q}$  on  $\Gamma_F \backslash \Gamma_\mathbb{Q}$ , is called the canonical field of definition. It is known (see [DE]) that  $S_K(\mathbb{C})$  has a canonical model over  $E$  that is denoted by  $S_K$ . Then  $S_K$  is called a quaternionic Shimura variety. The dimension of  $S_K$  is equal to  $d'$ .

### 3.1 Zeta function of quaternionic Shimura varieties

In this section we introduce some notations and we shall expose the computation of the zeta function for quaternionic Shimura varieties following closely [RT].

Let  $\pi$  be an automorphic representation of  $\bar{G}(\mathbb{A}_\mathbb{Q}) = G(\mathbb{A}_F)$ . Then  $\pi = \otimes \pi_v$ , where the restricted tensor product is taken over all places  $v$  of  $F$  and  $\pi_v$  is a representation of  $G(F_v)$ , where  $F_v$  is the completion of  $F$  at  $v$ . An irreducible representation  $\pi_v$  of  $\mathrm{GL}_2(F_v)$  is called unramified if  $\pi_v$  contains a nonzero vector that is fixed under  $\mathrm{GL}_2(O_v)$ , where  $O_v$  the completion of the ring of integers  $O_F$  at  $v$ . For almost all  $v$ , the representation  $\pi_v$  is unramified. We define  $L(s, \pi) := \prod_v L(s, \pi_v)$ , where if  $\pi_v$  is unramified

$$L(s, \pi_v) := \det(1 - Nv^{-s}g(\pi_v))^{-1},$$

and

$$g(\pi_v) = \begin{pmatrix} s_1 & 0 \\ 0 & s_2 \end{pmatrix}$$

denotes the Langlands class of  $\pi_v$ .

For a continuous representation  $r : {}^L\bar{G}^0 \rtimes \Gamma_E \rightarrow \mathrm{GL}_n(\mathbb{C})$  and an automorphic representation  $\pi = \otimes \pi_v$  of  $\bar{G}(\mathbb{A}_\mathbb{Q}) = G(\mathbb{A}_F)$ , where  $\pi_v$  denotes a representation of  $G(F_v)$ , one can define the  $L$ -function  $L(s, \pi, r) := \prod_v L(s, \pi_v, r)$ , where if  $\pi_v$  is unramified

$$L(s, \pi_v, r) := \det(1 - Nv^{-s}r(g(\pi_v)))^{-1}.$$

If  $\omega : \Gamma_E \rightarrow \mathrm{GL}_m(\mathbb{C})$  is an Artin representation, then we denote by the same symbol the representation of  ${}^L\bar{G}^0 \rtimes \Gamma_E$  that extends  $\omega$  and restricts to the trivial representation on  ${}^L\bar{G}^0$ . Then one can define as above the  $L$ -function  $L(s, \pi, r \otimes \omega)$ .

Consider  ${}^L\bar{T}^0$  to be the subgroup of  ${}^L\bar{G}^0$  of elements  $(t_\sigma)$  such that for all  $\sigma$ ,  $t_\sigma$  is diagonal and let  $\nu$  be the character of  ${}^L\bar{T}^0$  defined by

$$\nu((t_\sigma)) := \prod \nu_\sigma(t_\sigma),$$

$$\nu_\sigma\left(\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}\right) := \begin{cases} a & \text{for } \sigma \in J_F - J'_F; \\ 1 & \text{for } \sigma \in J'_F. \end{cases}$$

Then  $\Gamma_E$  stabilizes the character  $\nu$ .

We denote by  $r$  the finite dimensional representation of  ${}^L\bar{G}^0$  whose highest weight with respect to the standard Borel subgroup is  $\nu$ . Since  $\Gamma_E$  stabilizes  $\nu$ , the representation  $r$  could be uniquely extended to  ${}^L\bar{G}^0 \rtimes \Gamma_E$  such that  $\Gamma_E$  acts as the identity on the  $\nu$ -weight space. Then, the dimension of  $r$  is  $2^{d'}$ . From now on in this paper we fix this representation  $r$ .

Let  $K$  be an open compact subgroup of  $\bar{G}(\mathbb{A}_f)$ . If  $l$  is a prime number, let  $\mathbb{H}_K$  be the Hecke algebra generated by the bi- $K$ -invariant  $\mathbb{Q}_l$ -valued compactly supported functions on  $\bar{G}(\mathbb{A}_f)$  under the convolution. If  $\pi = \pi_f \otimes \pi_\infty$  is an automorphic representation of  $\bar{G}(\mathbb{A}_\mathbb{Q})$ , we denote by  $\pi_f^K$  the space of  $K$ -invariants in  $\pi_f$ . The Hecke algebra  $\mathbb{H}_K$  acts on  $\pi_f^K$ .

We have an action of the Hecke algebra  $\mathbb{H}_K$  and an action of the Galois group  $\Gamma_E$  on the étale cohomology  $H_{et}^i(S_K, \mathbb{Q}_l)$  and these two actions commute. We say that the representation  $\pi$  is *cohomological* if  $H^*(\mathfrak{g}, K_\infty, \pi_\infty) \neq 0$ , where  $\mathfrak{g}$  is the Lie algebra of  $K_\infty$  (the cohomology is taken with respect to  $(\mathfrak{g}, K_\infty)$ -module associated to  $\pi_\infty$ ). Then we know (see for example Proposition 1.8 of [RT]):

**Proposition 3.1.** *The representation of  $\Gamma_E \times \mathbb{H}_K$  on the étale cohomology  $H_{et}^i(S_K, \mathbb{Q}_l)$  is isomorphic to*

$$\bigoplus_{\pi} \sigma^i(\pi) \otimes \pi_f^K,$$

where  $\sigma^i(\pi)$  is a representation of the Galois group  $\Gamma_E$ . The above sum is over weight 2 irreducible cohomological automorphic representations  $\pi$  of  $\bar{G}(\mathbb{A}_\mathbb{Q})$ , such that  $\pi_f^K \neq 0$  and the  $\mathbb{H}_K$ -representations  $\pi_f^K$  are irreducible and mutually inequivalent.

The automorphic representations  $\pi$  which appear in Proposition 3.1 are one-dimensional or cuspidal and infinite-dimensional and we know the following result (see Propositions 1.5 and 1.8 of [RT]):

**Proposition 3.2.** *i) If  $\pi$  is infinite-dimensional, then*

$$\dim \sigma^i(\pi) = \begin{cases} 2^{d'} & \text{for } i = d', \\ 0 & \text{for } i \neq d'. \end{cases}$$

*ii) If  $\pi$  is one-dimensional, then*

$$\dim \sigma^i(\pi) = \begin{cases} \binom{d'}{i'} & \text{for } i = 2i', \\ 0 & \text{for } i \text{ odd.} \end{cases}$$

Fix an isomorphism  $i_l : \bar{\mathbb{Q}}_l \rightarrow \mathbb{C}$ , and define the  $L$ -function

$$L(s, S_K) := \prod_{\pi} \prod_v \prod_i \det(1 - Nv^{-s} i_l(\sigma^i(\pi)(\phi_v)) | H_{et}^i(S_K, \bar{\mathbb{Q}}_l)^{I_v})^{-1^{i+1}},$$

where  $\phi_v$  is a geometric Frobenius element at a finite place  $v$  of  $E$  and  $I_v$  is the inertia group at  $v$  (in order to define the local factors at the places of  $E$  dividing  $l$  one has to use actually the  $l'$ -adic cohomology for some  $l' \neq l$  and Theorem 3 of [B] which gives us the expression of the local factors of the zeta functions of quaternionic Shimura varieties).

For  $\pi$  cohomological, we define

$$m(\pi_{\infty}) = \begin{cases} (-1)^{d'} & \text{if } \pi_{\infty} \text{ is infinite-dimensional;} \\ 1 & \text{if } \pi_{\infty} \text{ is one-dimensional.} \end{cases}$$

We know (see Corollary 1.10 of [RT]):

**Proposition 3.3.** *There exists a finite set  $S$  of primes of  $E$ , such that for all primes  $v$  of  $E$  not in  $S$ , and for  $\pi$  cohomological of weight 2:*

$$\prod_i \det(1 - Nv^{-s} i_l(\sigma^i(\pi)(\phi_v)))^{(-1)^{i+1}} = \det(1 - Nv^{-s+(d'/2)} r_v(g(\pi_v)))^{-m(\pi_{\infty})},$$

where  $r_v$  denotes the restriction of  $r$  to  ${}^L\bar{G}^0 \rtimes G_v$ , and  $G_v$  is a decomposition group at  $v$ .

### 3.2 Twisted quaternionic Shimura varieties

Let  $\wp$  be a prime ideal of  $O_F$  such that  $G(F_{\wp})$  is isomorphic to  $\mathrm{GL}_2(F_{\wp})$ . Consider  $\mathbf{K} := K_{\wp} \times H$ , where  $K_{\wp}$  is the set of elements of  $\mathrm{GL}_2(O_{\wp})$  that are congruent to 1 modulo  $\wp$  and  $H$  is some open compact subgroup of the restricted product of  $(D \otimes_F F_{\mathfrak{p}})^{\times}$ , where  $\mathfrak{p}$  runs over all the finite places of  $F$ , with  $\mathfrak{p} \neq \wp$ . Then it is well known (see for example [C], Corollary 1.4.1.3) that for  $H$  sufficiently small, the group  $\mathrm{GL}_2(O/\wp)$  acts freely on  $S_{\mathbf{K}}$ . We fix such a small  $H$ . Then the action of  $\mathrm{GL}_2(O/\wp)$  on

$$S_{\mathbf{K}}(\mathbb{C}) = \bar{G}(\mathbb{Q}) \backslash X \times \bar{G}(\mathbb{A}_f) / \mathbf{K}$$

can be described in the following way : we have that  $\mathrm{GL}_2(O_{\wp}) \hookrightarrow \bar{G}(\mathbb{A}_{\mathbb{Q}})$  by  $\alpha \mapsto (1, \dots, \alpha, 1, \dots, 1)$ ,  $\alpha$  at the  $\wp$  component. Using the isomorphism  $\mathrm{GL}_2(O/\wp) \cong \mathrm{GL}_2(O_{\wp})/K_{\wp}$ , the action of an element  $g \in \mathrm{GL}_2(O_{\wp})$  is given by the right multiplication at the  $\wp$ -component.

We fix a continuous representation

$$\varphi : \Gamma_E \rightarrow \mathrm{GL}_2(O/\wp).$$

Let  $L$  be the finite Galois extension of  $E$  defined by  $L := (\bar{\mathbb{Q}})^{\mathrm{Ker}(\varphi)}$ .

Let

$$S' = S_{\mathbf{K}} \times_{\mathrm{Spec}(E)} \mathrm{Spec}(L).$$

The group  $\mathrm{GL}_2(O/\wp)$  acts on  $S_{\mathbf{K}}$ . Since  $\varphi : \mathrm{Gal}(L/E) \hookrightarrow \mathrm{GL}_2(O/\wp)$ , the group  $\mathrm{Gal}(L/E)$  acts on  $S_{\mathbf{K}}$ . We denote this action of  $\mathrm{Gal}(L/E)$  on  $S_{\mathbf{K}}$  by  $\varphi'$ . The Galois group  $\mathrm{Gal}(L/E)$  has a natural action on  $\mathrm{Spec}(L)$  and we can descend via the quotient process  $S'$  to  $S'_{\mathbf{K}}/\mathrm{Spec}(E)$  using the diagonal action

$$\mathrm{Gal}(L/E) \ni \sigma \rightarrow \varphi'(\sigma) \otimes \sigma$$

on  $S'$ . Thus, we obtain a quasi-projective variety  $S'_{\mathbf{K}}/\mathrm{Spec}(E)$ . This is the twisted quaternionic Shimura variety that we mentioned in the title.

### 3.3 Computation of the zeta function of twisted quaternionic Shimura varieties

We consider the injective limit:

$$V^i := \varinjlim_{\mathbf{K}} H_{et}^i(S_{\mathbf{K}}, \bar{\mathbb{Q}}_l) \cong \varinjlim_{\mathbf{K}} \oplus_{\pi} U^i(\pi) \otimes_{\bar{\mathbb{Q}}_l} \pi_f^K,$$

where  $U^i(\pi)$  is the  $\bar{\mathbb{Q}}_l$ -space that corresponds to  $\sigma^i(\pi)$  (see Proposition 3.1 for notations).

Then the  $\pi$ -component  $V^i(\pi)$  of  $V^i$  is isomorphic to  $\sigma^i(\pi) \otimes \pi_f$  as  $\Gamma_E \times \mathbb{H}$ -module. Taking the  $\mathbf{K}$ -fixed vectors, we deduce that  $V^i(\pi)^{\mathbf{K}}$  is isomorphic to  $\sigma^i(\pi) \otimes \pi_f^{\mathbf{K}}$  as  $\Gamma_E \times \mathrm{GL}_2(O/\wp O)$ -module. Since the varieties  $S_{\mathbf{K}}$  and  $S'_{\mathbf{K}}$  become isomorphic over  $\bar{\mathbb{Q}}$ , we have the isomorphism  $H_{et}^i(S_{\mathbf{K}}, \bar{\mathbb{Q}}_l) \cong H_{et}^i(S'_{\mathbf{K}}, \bar{\mathbb{Q}}_l)$ . The actions of  $\Gamma_E$  on these cohomologies that give the expression of the zeta functions of these varieties are different. If we consider the component  $V^{i'}(\pi)$  that corresponds to  $\pi$  of  $H_{et}^i(S'_{\mathbf{K}}, \bar{\mathbb{Q}}_l)$  (see the decomposition of Proposition 3.1), we get that  $V^{i'}(\pi)$  is isomorphic to  $\sigma^i(\pi) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)$  as  $\Gamma_E$ -module.

We consider a local field  $L$  of characteristic 0 and residue characteristic  $p$ . Let  $W_L \subset \Gamma_L$  be the Weil group. The *Weil – Deligne* group  $WD_L$  of  $L$  is defined as the semidirect product of  $W_L$  with  $\mathbb{C}$  by the relation

$$\sigma z \sigma^{-1} = |\sigma| z$$

for all  $\sigma \in W_L$  and  $z \in \mathbb{C}$ , where  $|\cdot|$  is the norm map:  $|\cdot| : W_L \rightarrow q^{\mathbb{Z}} \subset \mathbb{Q}^{\times}$ , where  $q$  denotes the cardinality of the residue field of  $L$ , and  $|\cdot| = 1$  on the inertia group  $I_L \subset W_L$  and  $|\Phi| = q$ , where  $\Phi \in W_L$  is an arithmetic Frobenius.

Fix a prime number  $l$  different from  $p$ . For a vector space  $V$  of finite dimension over  $\bar{\mathbb{Q}}_l$ , let  $\rho : \Gamma_L \rightarrow \mathrm{GL}(V)$  be a continuous  $l$ -adic representation. We denote also by  $\rho$  its restriction to  $W_L$ . To the  $l$ -adic representation  $\rho$ , one can associate (see for example [B] for details) a pair  $(\rho^*, N)$  called *Frobenius semisimple parameter* of  $\rho$ , where  $N \in \mathrm{End}(V)$  is a nilpotent endomorphism and  $\rho^*$  is a representation of  $WD_L$  having the propriety that  $\rho^*|_{W_L}$  is semisimple and for all  $\sigma \in W_L$ ,  $\rho^*(\sigma)$  is semisimple.

We know the following result which is a generalization of Proposition 3.3 above (see Theorem 3 of [B]):

**Proposition 3.4.** *Let  $l$  be a prime number and  $\pi$  be a cuspidal automorphic representation as in Proposition 3.1. Then for each finite place  $v$  of  $E$  whose residue characteristic  $p$  is different from  $l$ , the isomorphism class of the Frobenius semisimple parameter  $(\rho_{K,v}^*, N_v)$  of the Weil-Deligne group  $WD_{E_v}$  of  $E_v$  defined by the restriction of  $\sigma^{d'}(\pi)$  to a decomposition group  $D_v$  at  $v$  coincides with the class*

$$((r \otimes | \cdot |^{d'/2}) \circ \sigma(\pi_v)), N_{K,v})$$

*obtained by the restriction of  $(r \otimes | \cdot |^{d'/2}) \circ \sigma$  to the decomposition group  $D_v$ , where  $\sigma : \Gamma_E \hookrightarrow^L \bar{G}^0 \rtimes \Gamma_E$  is the inclusion and  $\sigma(\pi_v) : WD_{E_v} \rightarrow^L \bar{G}^0 \rtimes \Gamma_E$  is the standard homomorphism.*

**Proof:** The idea of the proof of the Proposition 3.4 (for details see [B]) is that the representations  $\sigma^{d'}(\pi)$  and  $((r \otimes | \cdot |^{d'/2}) \circ \sigma)(\phi_v)$  satisfy (see §5.2 of [B]) the *Weight Monodromy Conjecture* (i.e. the eigenvalues of  $\sigma^{d'}(\pi)(\phi_v)$  and  $((r \otimes | \cdot |^{d'/2}) \circ \sigma)(\phi_v)$  are *Nv-Weil numbers of weight  $d'$* , which means that the eigenvalues are algebraic integers  $\alpha$  having the propriety that for each automorphism  $\sigma$  of  $\bar{\mathbb{Q}}$ , we have  $|\sigma(\alpha)| = |Nv|^{d'/2}$  and thus for each  $v$  their corresponding nilpotent data  $N_{K,v}$  and  $N_v$  are uniquely determined (for details see §1.12 of [B]) by the semisimple representations  $\rho_{K,v}^*$  and  $(r \otimes | \cdot |^{d'/2}) \circ \sigma(\pi_v)$ . Hence it is sufficient to prove that

$$\rho_{K,v}^* = (r \otimes | \cdot |^{d'/2}) \circ \sigma(\pi_v).$$

But we know that for almost all  $v$ , (i)  $N_{K,v} = 0$  and  $N_v = 0$ , (ii)  $\rho_{K,v}^*$  and  $(r \otimes | \cdot |^{d'/2}) \circ \sigma(\pi_v)$  are unramified and from the computation of the unramified zeta function (see [R] and [BL], or Proposition 3.3 above) shows that this formula is true. From Chebotarev theorem, we see that the semisimplification  $\sigma^{d'}(\pi)^{ss}$  is isomorphic to  $(r \otimes | \cdot |^{d'/2}) \circ \sigma$  and thus their restrictions to the decomposition group  $D_v$  for  $v \nmid l$  are isomorphic and thus they give rise to isomorphic parameters  $\rho_{K,v}^*$  and  $(r \otimes | \cdot |^{d'/2}) \circ \sigma(\pi_v)$  and we conclude Proposition 3.4. ■

We prove now the following result which implies the first part of Theorem 1.2:

**Proposition 3.5.** *Let  $l$  be a prime number and  $\pi$  be a cuspidal automorphic representation as in Proposition 3.1. Then for each finite place  $v$  of  $E$  whose residue characteristic is different from  $l$ , the isomorphism class of the Frobenius semisimple parameter  $(\rho_{\mathbf{K},v}^*, N'_v)$  of the Weil-Deligne group  $WD_{E_v}$  of  $E_v$  defined by the restriction to the decomposition group  $D_v$  at  $v$  of  $\sigma^{d'}(\pi) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)$  coincides with the class of*

$$(((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma(\pi_v), N'_{K,v})$$

*obtained by the restriction to the decomposition group  $D_v$  of  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma$ .*

**Proof:** From the proof of Proposition 3.4, we know that  $\sigma^{d'}(\pi)$  and  $(r \otimes | \cdot |^{d'/2}) \circ \sigma$  satisfy the *Weight Monodromy Conjecture*, and thus from Brauer's induction theorem, we get that the representations  $\sigma^{d'}(\pi) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)$  and  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma$  satisfy also the *Weight Monodromy Conjecture* and thus for each  $v$  their corresponding nilpotent data  $N'_v$  and  $N'_{\mathbf{K},v}$  are uniquely determined by the semisimple representations  $\rho'_{\mathbf{K},v}$  and  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma(\pi_v)$ . Thus it is sufficient to show that

$$\rho'_{\mathbf{K},v} = ((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma(\pi_v).$$

But again we know that for almost all  $v$ , (i)  $N'_v = 0$  and  $N'_{\mathbf{K},v} = 0$ , (ii)  $\rho'_{\mathbf{K},v}$  and  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma(\pi_v)$  are unramified and Proposition 3.3 combined with Brauer's induction theorem shows that this formula is true. From Cebotarev theorem, we see that the semisimplification  $(\sigma^{d'}(\pi) \otimes (\pi_f^{\mathbf{K}} \circ \varphi))^{ss}$  is isomorphic to  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma$  and thus their restrictions to the decomposition group  $D_v$  for  $v \nmid l$  are isomorphic and thus they give rise to isomorphic parameters  $\rho'_{\mathbf{K},v}$  and  $((r \otimes | \cdot |^{d'/2}) \otimes (\pi_f^{\mathbf{K}} \circ \varphi)) \circ \sigma(\pi_v)$  and we conclude Proposition 3.5. ■

## 4 Meromorphic continuation

As an application to Theorem 1.1, in this section we prove the second part of Theorem 1.2 regarding the meromorphic continuation of zeta functions of the twisted quaternionic Shimura varieties defined in §3.

In this section, under some conditions, we continue meromorphically the zeta function  $L(s, S'_{\mathbf{K}})$  to the entire complex plane and show that it satisfies also a functional equation.

Let  $\omega = \pi_f^{\mathbf{K}} \circ \varphi$ . We define  $L := \bar{\mathbb{Q}}^{\text{Ker}(\varphi)}$  and  $K := \bar{\mathbb{Q}}^{\text{Ker}(\omega)}$ . From now on, in this paper, we assume that  $L$  is a solvable extension of a totally real field. Since  $K \subseteq L$ , the field  $K$  is also a solvable extension of a totally real field.

### 4.1 Definition of the representation $\rho(\pi)$

One can find a representation  $\rho(\pi)$  of  $\Gamma_E$  ([BR] §7.2) such that

$$L(s, \rho(\pi)) = L(s - d'/2, \pi, r).$$

We describe now the representation  $\rho(\pi)$ . Let  $G$  be a group and  $K$  and  $H$  be two subgroups of  $G$ . We consider a representation

$$\tau : H \rightarrow \text{GL}(W)$$

and a double coset  $H\sigma K$  such that  $d(\sigma) = |H \setminus H\sigma K| < \infty$ . We define a representation  $\tau_{H\sigma K}$  of  $K$  on the  $d(\sigma)$ -fold tensor product  $W^{\otimes d(\sigma)}$ . Consider the representatives  $\{\sigma_1, \dots, \sigma_{d(\sigma)}\}$  such that  $H\sigma K = \cup H\sigma_j$ . If  $\gamma \in K$ , then there exists  $\xi_j \in H$  and an index  $\gamma(j)$  such that

$$\sigma_j \gamma = \xi_j \sigma_{\gamma(j)}.$$

We define the representation:

$$\tau_{H\sigma K}(\gamma)(\omega_1 \otimes \cdots \otimes \omega_{d(\sigma)}) = \tau(\xi_1)\omega_{\gamma^{-1}(1)} \otimes \cdots \otimes \tau(\xi_{d(\sigma)})\omega_{\gamma^{-1}(d(\sigma))}.$$

One can prove easily that the equivalence class of  $\tau_{H\sigma K}$  is independent of the choice of the representatives  $\sigma_1, \dots, \sigma_{d(\sigma)}$ .

Let  $J_F - J'_F = \{\delta_1, \dots, \delta_{d'}\}$ , and  $S := \bigcup \Gamma_F \delta_i$ . We write  $S$  as a disjoint union of double cosets

$$S = \bigcup_{j=1}^k \Gamma_F \sigma_j \Gamma_E$$

and we denote by  $\rho_j$  the representation of  $\Gamma_E$  defined by  $\rho_{\pi, \lambda}$ , and the double coset  $\Gamma_F \sigma_j \Gamma_E$ . Then our representation  $\rho(\pi)$  is isomorphic to  $\rho_1 \otimes \cdots \otimes \rho_k$ . Thus

$$L(s - d'/2, \pi, r) = L(s, \rho(\pi)) = L(s, \rho_1 \otimes \cdots \otimes \rho_k)$$

and we obtain also that

$$L(s - d'/2, \pi, r \otimes \omega) = L(s, \rho(\pi) \otimes \omega) = L(s, \rho_1 \otimes \cdots \otimes \rho_k \otimes \omega).$$

## 4.2 Base change and Brauer's theorem

**Lemma 4.1.** *Let  $\phi$  be an  $l$ -adic representation of  $\Gamma_E$ . Suppose that there exists a Galois extension  $F'$  of  $\mathbb{Q}$ , which contains the field  $K := \mathbb{Q}^{\ker(\omega)}$ , and that the  $L$ -function  $L(s, \phi|_{\Gamma_{F''}} \otimes \chi)$  can be meromorphically continued to the entire complex plane and satisfies a functional equation for any subfield  $F''$  of  $F'$  containing  $E$  such that  $F'$  is a solvable extension of  $F''$  and any finite order character  $\chi$  of  $\Gamma_{F''}$ . Then  $L(s, \phi \otimes \omega)$  can be meromorphically continued to the entire complex plane and satisfies a functional equation.*

**Proof:** From Brauer's theorem (see Theorems 16 and 19 of [SE]), we know that there exist some subfields  $F_i \subset F'$  such that  $\text{Gal}(F'/F_i)$  are solvable, some characters  $\chi_i : \text{Gal}(F'/F_i) \rightarrow \mathbb{Q}^\times$  and some integers  $m_i$ , such that the representation

$$\omega : \text{Gal}(F'/E) \rightarrow \text{Gal}(K/E) \rightarrow \text{GL}_N(\bar{\mathbb{Q}}_l),$$

can be written as  $\omega = \sum_{i=1}^{i=k} m_i \text{Ind}_{\Gamma_{F_i}}^{\Gamma_E} \chi_i$  (a virtual sum). Then

$$\begin{aligned} L(s, \phi \otimes \omega) &= \prod_{i=1}^{i=k} L(s, \phi \otimes \text{Ind}_{\Gamma_{F_i}}^{\Gamma_E} \chi_i)^{m_i} \\ &= \prod_{i=1}^{i=k} L(s, \text{Ind}_{\Gamma_{F_i}}^{\Gamma_E} (\phi|_{\Gamma_{F_i}} \otimes \chi_i))^{m_i} = \prod_{i=1}^{i=k} L(s, \phi|_{\Gamma_{F_i}} \otimes \chi_i)^{m_i}. \end{aligned}$$

We know from our assumption that the  $L$ -functions  $L(s, \phi|_{\Gamma_{F_i}} \otimes \chi_i)$  have a meromorphic continuation to the entire complex plane and verify a functional equation. Thus  $L(s, \phi \otimes \omega)$  can be meromorphically continued to the entire complex plane and satisfies a functional equation. ■

### 4.3 Meromorphic continuation of the zeta functions for curves and surfaces

In this section we prove the second part of Theorem 1.2, that is a consequence of Theorem 4.3 below.

It is known (Theorem M of [RA1]) that:

**Proposition 4.2.** *If  $\pi_1$  and  $\pi_2$  are two cuspidal automorphic representations of  $GL(2)/L$ , where  $L$  is a number field, then  $\pi_1 \otimes \pi_2$  is an automorphic (isobaric) representation of  $GL(4)/L$ .*

We prove now the following result:

**Theorem 4.3.** *If  $K := \bar{\mathbb{Q}}^{Ker(\omega)}$  is a solvable extension of a totally real field and  $d' = 1$  or  $d' = 2$ , then the function  $L(s - d'/2, \pi, r \otimes \omega) = L(s, \rho(\pi) \otimes \omega)$  can be meromorphically continued to the entire complex plane and satisfies a functional equation.*

**Proof:** It is sufficient to show that there exists a Galois extension  $F'$  of  $\mathbb{Q}$ , which contains  $F$  and  $K$ , such that  $\rho(\pi)|_{\Gamma_{F'}}$  satisfies the conditions of Lemma 4.1.

We have two cases:

a)  $d' = 1$ . We assume for simplicity that  $J_F - J'_F = \{1\}$ , where 1 is the trivial embedding of  $F$  in  $\bar{\mathbb{Q}}$ . In this case  $E = F$  and  $\rho(\pi) \cong \rho_{\pi, \lambda}$ . From Theorem 1.1, we deduce that one can find a field  $F'$  which is Galois over  $\mathbb{Q}$ , and contains  $F$  and  $K$ , such that  $\rho_{\pi, \lambda}|_{\Gamma_{F'}}$  is modular. From Langlands base change for  $GL(2)$  for solvable extensions ([L]), we obtain that  $\rho(\pi)|_{\Gamma_{F'}}$  satisfies the conditions of Lemma 4.1.

b)  $d' = 2$ . We assume for simplicity that  $J_F - J'_F = \{1, c\}$ , where 1 is the trivial embedding of  $F$  in  $\bar{\mathbb{Q}}$ . We denote by the same symbol  $c$  the extension of  $c$  to  $\bar{\mathbb{Q}}$ . Then,

$$S = \Gamma_F \cup \Gamma_F c.$$

The stabilizer of  $S$  is  $\Gamma_E$ . It is easy to see that the stabilizer of  $S$  is equal to  $(\Gamma_{Fc} \cap c^{-1}\Gamma_F) \cup (\Gamma_F \cap c^{-1}\Gamma_{Fc})$ . Thus we get

$$\Gamma_E = (\Gamma_{Fc} \cap c^{-1}\Gamma_F) \cup (\Gamma_F \cap c^{-1}\Gamma_{Fc}).$$

We distinguish two cases:

i)  $\Gamma_{Fc} \cap c^{-1}\Gamma_F = \emptyset$ . Then,  $\Gamma_E = \Gamma_F \cap c^{-1}\Gamma_{Fc}$ . Thus,

$$F \subset E \subset F^{gal}$$

where  $F^{gal}$  is the Galois closure of  $F$ . We have

$$S = \Gamma_F \cup \Gamma_{Fc} = \Gamma_F \Gamma_E \cup \Gamma_{Fc} \Gamma_E.$$

If  $\gamma \in \Gamma_E$ , then

$$\tau_{\Gamma_F 1 \Gamma_E}(\gamma)(\omega_1) = \rho_{\pi, \lambda}(\gamma)(\omega_1)$$

and

$$\tau_{\Gamma_F c \Gamma_E}(\gamma)(\omega_1) = \rho_{\pi, \lambda}(c\gamma c^{-1})(\omega_1).$$

Thus

$$\rho(\pi) \cong \rho_{\pi, \lambda}|_{\Gamma_E} \otimes \rho_{\pi, \lambda}|_{\Gamma_E}^c,$$

where

$$\rho_{\pi, \lambda}|_{\Gamma_E}^c(\gamma) = \rho_{\pi, \lambda}(c\gamma c^{-1}).$$

The representation  $\pi$  is one-dimensional or cuspidal and infinite-dimensional. If  $\pi$  is one-dimensional, then  $\pi(g) = \rho_{\pi}(N(g))|N(g)|^{1/2}$ , where  $N$  is the reduced norm map and  $|\cdot|$  denotes the ideles norm and  $\rho_{\pi}$  is a Hecke character.

From Theorem 1.1, we deduce that one can find a solvable extension of a totally real field  $F'$  which is Galois over  $\mathbb{Q}$ , and contains  $F$  and  $K$ , such that  $\rho_{\pi, \lambda}|_{\Gamma_{F'}}$  is modular. Thus

$$\rho(\pi)|_{\Gamma_{F'}} \cong \rho_{\pi, \lambda}|_{\Gamma_{F'}} \otimes \rho_{\pi, \lambda}|_{\Gamma_{F'}}^c,$$

is a tensor product of two automorphic representations and from Langlands base change for  $GL(2)$  for solvable extensions ([L]) and Proposition 4.2, we obtain that  $\rho(\pi)|_{\Gamma_{F'}}$  satisfies the conditions of Lemma 4.1.

ii)  $\Gamma_F c \cap c^{-1} \Gamma_F \neq \emptyset$ . Let  $\Gamma_{E_1} := \Gamma_F \cap c^{-1} \Gamma_F c$ . Thus

$$F \subset E_1 \subset F^{gal}.$$

Since it is obvious now that  $\Gamma_{E_1} \subset \Gamma_E$ ,  $[\Gamma_E : \Gamma_{E_1}] = 2$  and  $\Gamma_E \not\subseteq \Gamma_F$ , we get  $[E_1 : E] = 2$  and  $F \not\subseteq E$ . If  $F_1 := E \cap F$ , then  $[F : F_1] = 2$  and we can easily see that  $c$  when restricted to  $F_1$  is the trivial embedding. Hence  $c$  is the nontrivial automorphism of  $F$  over  $F_1$  and we get that  $\Gamma_{E_1} = \Gamma_F \cap c^{-1} \Gamma_F c = \Gamma_F$ , which means that  $E_1 = F$  and  $E = F_1$  and therefore we have  $[F : E] = 2$  and  $c$  is the nontrivial automorphism of  $F$  over  $E$ .

We have

$$S = \Gamma_F \cup \Gamma_F c = \Gamma_F 1 \Gamma_E.$$

If  $\gamma \in \Gamma_F$ , then

$$\tau_{\Gamma_F 1 \Gamma_E}(\gamma)(\omega_1 \otimes \omega_2) = \rho_{\pi, \lambda}(\gamma)\omega_1 \otimes \rho_{\pi, \lambda}(c\gamma c^{-1})\omega_2.$$

If  $\gamma \in \Gamma_E - \Gamma_F$ , then

$$\tau_{\Gamma_F 1 \Gamma_E}(\gamma)(\omega_1 \otimes \omega_2) = \rho_{\pi, \lambda}(\gamma c^{-1})\omega_2 \otimes \rho_{\pi, \lambda}(c\gamma)\omega_1.$$

Thus  $\rho(\pi)$  is a subrepresentation of

$$\text{Ind}_{\Gamma_F}^{\Gamma_E}(\rho_{\pi, \lambda} \otimes \rho_{\pi, \lambda}^c)$$

that satisfies

$$\rho(\pi)|_{\Gamma_F} \cong \rho_{\pi,\lambda} \otimes \rho_{\pi,\lambda}^c,$$

and from Theorem [D] of [RA2] we know that  $\rho(\pi)$  is automorphic.

As in i) above, we deduce that for some solvable extension of a totally real field  $F'$  which is Galois over  $\mathbb{Q}$ , and contains  $F$  and  $K$ , the representation  $\rho(\pi)|_{\Gamma_{F'}}$  satisfies the conditions of Lemma 4.1. ■

## References

- [AC] J.Arthur, L.Clozel, *Simple algebras, base change and the advanced theory of the trace formula*, Ann. of Math. Studies, Princeton University Press, 1989.
- [B] D.Blasius, *Hilbert modular forms and the Ramanujan conjecture*, Noncommutative geometry and number theory, 35–56, Aspects Math., E37, Vieweg, Wiesbaden, 2006.
- [BL] J.L.Brylinski, J.P.Labesse, *Cohomologie d'intersection et fonctions L de certaines varietes de Shimura*, Annales Scientifiques de l'Ecole Normale Supérieure, 17, 1984, 361-412.
- [BR] D.Blasius, J.D.Rogawski, *Zeta functions of Shimura varieties*, Motives, AMS Proc. Symp. Pure Math. 55, Part 2.
- [C] H.Carayol, *Sur la mauvaise réduction des courbes de Shimura*, Compositio Math., 59, nr.2 (1986), 151-230.
- [D] M.Dimitrov, *Galois representations mod p and cohomology of Hilbert modular varieties*, Ann. Sci. de l'Ecole Norm. Sup. 38, 2005, 505-551.
- [DE] P. Deligne, *Travaux de Shimura*, Sémin. Bourbaki Fév. 71, Exposé 389, Lectures Notes in Math. vol. 244. Berlin-heidelberg-New York; Springer 1971.
- [G] S.S.Gelbart, *Automorphic forms on adeles groups*, Ann. of Math. Studies, Princeton University Press, 1975.
- [HSBT] M.Harris, N.Shepherd-Barron, R.Taylor, *A family of Calabi-Yau varieties and potential automorphy*, to appear in Ann. of Math.
- [L] R. P. Langlands, *Base change for  $GL_2$* , Ann. of Math. Studies 96, Princeton University Press, 1980.
- [R] H.Reimann, *The semi-simple zeta function of quaternionic Shimura varieties*, Lecture Notes in Mathematics 1657, Springer, 1997.
- [RA1] D.Ramakrishnan, *Modularity of the Rankin-Selberg L-series, and multiplicity one for  $SL(2)$* , Ann. of Math., 152(2000), 45-111.

- [RA2] D.Ramakrishnan, *Modularity of solvable Artin representations of  $GO(4)$ -type*, IMRN 2002, No. 1, 1-54.
- [RT] J.D.Rogawski, J.B.Tunnell, *On Artin  $L$ -functions associated to Hilbert modular forms of weight one*, Inv. Math., 74, 1983, 1-43.
- [SE] J.-P.Serre, *Linear representations of finite groups*, Springer 1977.
- [T1] R.Taylor, *On Galois representations associated to Hilbert modular forms*, Inv. Math., 98, 1989, 265-280.
- [T2] R.Taylor, *On the meromorphic continuation of degree two  $L$ -functions*, Documenta Mathematica, Extra Volume: John Coates' Sixtieth Birthday (2006), 729-779.
- [V1] C.Virdol, *Zeta functions of twisted modular curves*, J. Aust. Math. Soc. 80 (2006), 89-103.
- [V2] C.Virdol, *Tate classes and poles of  $L$ -functions of twisted quaternionic Shimura surfaces*, J. of Number Theory 123, Nr. 2 (2007), 315-328.
- [W] A. Wiles, *Modular elliptic curves and Fermat's last theorem*, Ann. of Math. 141, (1995), 443-551.