

MATCHING OF HECKE OPERATORS FOR EXCEPTIONAL DUAL PAIR CORRESPONDENCES

GORDAN SAVIN AND MICHAEL WOODBURY

to Steve Rallis, in memoriam

1. INTRODUCTION

Let F be a p -adic field with ring of integers \mathcal{O} and fixed inverse of uniformizer ϖ . Let q be the order of the residual field. We fix an absolute value on F so that $|\varpi| = q$. We consider exceptional dual pairs $G \times G'$ inside of an adjoint group \mathbf{G} where each group consists of the F points of a split reductive algebraic group. Denote by K and K' hyperspecial maximal compact subgroups of G and G' respectively. Let (Π, \mathbf{V}) be the minimal representation of \mathbf{G} .

If σ' is an irreducible representation of G' , we call an irreducible representation σ of G a Θ -lift of σ' if $\sigma \otimes \sigma'$ is a quotient of Π . If σ, σ' are spherical, we will prove that the correspondence $\sigma \leftrightarrow \sigma'$ is functorial with respect to a natural injection on the dual groups

$$r : \widehat{G}'(\mathbb{C}) \rightarrow \widehat{G}(\mathbb{C}).$$

To be more precise, let C be the centralizer of $r(\widehat{G}')$ in \widehat{G} . Then C is a reductive, possibly finite, group. Let $f : \mathrm{SL}_2(\mathbb{C}) \rightarrow C$ be a map corresponding to the regular unipotent orbit in C by Jacobson-Morozov. Let

$$s = f \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}.$$

Let \mathcal{H} and \mathcal{H}' denote the spherical Hecke algebras of G and G' respectively. Let $T \in \mathcal{H}$ correspond, via the Satake isomorphism, to a finite dimensional representation V of $\widehat{G}(\mathbb{C})$. (We describe this in more detail in Section 2 below.) Write $V = \sum V' \otimes V''$, the restriction of V to $\widehat{G}' \otimes C$. We define a map $\tilde{r} : \mathcal{H} \rightarrow \mathcal{H}'$ by

$$(1) \quad \tilde{r}(V) = \sum_{V'} \mathrm{Tr}_{V''}(s)V'.$$

Note that if C is finite, then $s = 1$, the identity in \widehat{G} , and $\tilde{r}(V)$ is just the restriction of V to \widehat{G}' . We consider the following dual pairs:

| | | | | | |
|--------------|-------|------------------|------------------|-------------------|-------|
| \mathbf{G} | D_4 | D_5 | E_6 | E_7 | E_8 |
| G' | S_3 | PGL_2 | PGL_3 | G_2 | G_2 |
| G | G_2 | G_2 | G_2 | PGSp_6 | F_4 |

TABLE 1. Dual pairs $G \times G' \subset \mathbf{G}$

Partially supported by NSF grant DMS 0852429.

Theorem 1.1. *For the dual pairs $G \times G' \subset \mathbf{G}$ in the above table, $T \in \mathcal{H}_G$ and \tilde{r} given by (1), $\Pi(T) = \Pi(\tilde{r}(T))$ as operators on $\mathbf{V}^{K \times K'}$.*

As a matter of terminology, if the actions of T and $\tilde{r}(T)$ agree on a space V , or, more precisely, on a subset of fixed vectors, for all $T \in \mathcal{H}_G$ we will say there is a *matching of Hecke operators* of \mathcal{H}_G and $\mathcal{H}_{G'}$ on V or, more concisely, *matching* on V . We trust that the precise space of fixed vectors will be clear from context.

An analogue of Theorem 1.1 is well known in the case of classical theta correspondences [5]. For exceptional groups, the first example of matching was obtained by Rallis and Soudry in [6].

The proof of Theorem 1.1 is by induction on the rank of \mathbf{G} . The main tool is Jacquet functors of \mathbf{V} with respect to maximal parabolic subgroups of dual pairs. Most of the needed functors were computed in [4]. One remaining, but rather remarkable case (for $\mathbf{G} = E_8$), is computed in the last section.

2. THE SATAKE ISOMORPHISM

Before giving the proof of Theorem 1.1 we review the facts about the Hecke algebra and the Satake isomorphism (most of which can be found in [2]) which will be relevant, and we give some general lemmas which will be key in the proof of Theorem 1.1.

2.1. The Hecke algebra. Let G be a split reductive group, K a hyperspecial maximal compact subgroup, $B = TU$ a Borel subgroup. There is an Iwasawa decomposition $G = BK$, and the choice of Borel gives a set Φ^+ of positive roots.

We identify the cocharacters of T , $X_\bullet(T)$, with the coweight lattice Λ_c , and we define

$$\Lambda_c^+ = \{\lambda \in \Lambda_c \mid \langle \lambda, \alpha \rangle \geq 0 \text{ for all } \alpha \in \Phi^+\}.$$

Thus, via our identification, each element $\lambda \in \Lambda_c^+$ can be viewed as a map $\lambda : F^\times \rightarrow T$.

Proposition 2.1. *The group G is the disjoint union of double cosets $K\lambda(\varpi)K$ for $\lambda \in \Lambda_c^+$.*

Example. $G = \mathrm{GL}_3(F)$, $\widehat{G} = \mathrm{GL}_3(\mathbb{C})$. If $\alpha_1 = (1, -1, 0)$ and $\alpha_2 = (0, 1, -1)$ are the simple roots, then P^+ consists of $\lambda = (l, m, n)$ such that $l \geq m \geq n$. Then

$$\lambda(t) = \begin{pmatrix} t^l & & \\ & t^m & \\ & & t^n \end{pmatrix}$$

and it is easy to verify directly that

$$\mathrm{GL}_3(F) = \bigsqcup_{l \geq m \geq n} \mathrm{GL}_3(\mathcal{O}) \begin{pmatrix} \varpi^l & & \\ & \varpi^m & \\ & & \varpi^n \end{pmatrix} \mathrm{GL}_3(\mathcal{O}).$$

Recall that the irreducible representations of \widehat{G} , the complex dual group of G , are parametrized by their highest weights $\lambda \in \Lambda_c^+$. Let $R(\widehat{G})$ be the representation ring of \widehat{G} . That is, $R(\widehat{G})$ is the \mathbb{C} -vector space with basis consisting of the irreducible representation of \widehat{G} . We denote the representation of highest weight λ by V_λ , and consider the map

$$(2) \quad R(\widehat{G}) \rightarrow \mathbb{C}[\Lambda_c] \quad V_\mu \mapsto \sum_{\mu} m_\lambda(\mu)[\mu]$$

where $m_\lambda(\mu)$ is the dimension of the μ -weight space in V_λ . Letting W denote the Weyl group, this gives an isomorphism $R(\widehat{G}) \simeq \mathbb{C}[\Lambda_c]^W$.

The Hecke algebra \mathcal{H}_G consists of all locally constant compactly supported K -biinvariant functions $f : G \rightarrow \mathbb{C}$. By Proposition 2.1, \mathcal{H}_G has a basis consisting of the characteristic functions of $K\lambda(\varpi)K$ with $\lambda \in \Lambda_c^+$. We denote these by T_λ .

If (σ, V) is a smooth G -module then the action of $f \in \mathcal{H}_G$ is given by

$$f * v = \int_G f(g)\sigma(g)v dg.$$

We normalize the Haar measure dg so that $\text{vol}(K) = 1$. Let U be the unipotent radical of B as above. By the Iwasawa decomposition, we can write

$$(3) \quad K\lambda(\varpi)K = \bigcup_u \bigcup_t utK$$

for some representatives $t \in T$ and $u \in U$. So if v is a K -fixed vector then

$$(4) \quad T_\lambda * v = \sum_{u,t} \sigma(u)\sigma(t)v.$$

If $r_U : V \rightarrow V_U$ is the natural projection, where V_U is the space of U -coinvariants, then

$$(5) \quad r_U(T_\lambda * v) = \sum_t n(t)r_U(\sigma(t)v)$$

where $n(t)$ is the number of single cosets of type utK appearing in $K\lambda(\varpi)K$.

2.2. The (relative) Satake transform. Let δ_U denote the modular character of B given by

$$d(bub^{-1}) = \delta_U(b)du.$$

The measure du is normalized so that $\text{vol}(K \cap U) = 1$. Obviously, δ_U is trivial on U , and so this defines a character $\delta : T \rightarrow \mathbb{R}_+^\times$. We take $\delta_U^{1/2}(t)$ to be the positive square root of this character. Let Φ^+ denote the positive roots of G (determined by B .) Then if $\mu \in X_\bullet(T)$,

$$(6) \quad \delta^{1/2}(\mu(\varpi)) = q^{\langle \mu, \rho \rangle}, \quad 2\rho = \sum_{\alpha \in \Phi^+} \alpha.$$

The Satake transform $\mathcal{S}_T : \mathcal{H}_G \rightarrow \mathcal{H}_T$ is given by

$$(7) \quad \mathcal{S}_T f(t) = \delta(t)^{1/2} \int_U f(tu)du.$$

It is a fact that \mathcal{S}_T is injective and its image is equal to the Weyl group invariants. Since $\mathcal{H}_T = \mathbb{C}[X_\bullet(T)] = \mathbb{C}[\Lambda_c]$, it follows from our discussion above that this defines an isomorphism $\mathcal{S} : \mathcal{H}_G \rightarrow R(\widehat{G})$.

The Satake transform can be defined analogously for any parabolic $P = MN$. This yields the *relative Satake transform* $\mathcal{S}_M : \mathcal{H}_G \rightarrow \mathcal{H}_M$:

$$(8) \quad \mathcal{S}_M f(m) = \delta_N^{1/2}(m) \int_N f(mn)dn.$$

As above, dn is the measure which gives $N \cap K$ volume 1, and $\delta_N : M \rightarrow \mathbb{R}_+^\times$ is the modular character.

We may assume $P \supset B$, so the composition of \mathcal{S}_M with the Satake transform from \mathcal{H}_M to \mathcal{H}_T is \mathcal{S}_T .

2.3. Some general lemmas. In this section, we prove various simple lemmas which will be used in our proof of Theorem 1.1.

Throughout this paper parabolic induction and the Jacquet functors will be normalized as follows. Let $P = MN$ be a parabolic subgroup of G . Suppose that (σ, W) is a representation of M which we extend trivially to P . Then we define $(\rho, i_P^G(W))$ to be the representation of G consisting of smooth functions $f : G \rightarrow W$ which satisfy,

$$(9) \quad f(mng) = \delta_N^{1/2}(m)\sigma(m)f(g) \quad \text{for all } m \in M, n \in N, g \in G$$

with right regular action $\rho(g)f : h \mapsto f(hg)$. Note that the usual induction functor is

$$(10) \quad \text{Ind}_P^G(W) = i_P^G(\delta_N^{-1/2} \otimes W).$$

If (π, V) is a representation of G , the Jacquet functor with respect to N , $(\pi_M, r_N(V))$, is defined as follows. As usual,

$$V(N) = \langle \pi(n)v - v \mid n \in N, v \in V \rangle,$$

so $V_N = V/V(N)$ is the space of coinvariants. Let $(\pi_M, r_N(V))$ be a representation of M such that $r_N(V) = V_N$, but the M -action is given by

$$(11) \quad \pi_M(m)v = \delta_N^{-1/2}(m)\pi(m)v.$$

Since N is normal, it is trivial to see that this is well defined.

Since the induction and the Jacquet functor are normalized, the statement of Frobenius reciprocity is quite simple:

$$(12) \quad \text{Hom}_G(V, i_P^G(W)) = \text{Hom}_M(r_N(V), W).$$

Lemma 2.2. *Suppose that G is a split reductive group, $K \subset G$ is a maximal compact subgroup and $MN = P \subset G$ is a parabolic subgroup. Let $K_M = K \cap M$. For (π, V) any smooth representation of M , the following statements hold.*

- (i) *The map $\varphi : (i_P^G(V))^K \rightarrow V^{K_M}$ given by $f \mapsto f(1)$ is an isomorphism.*
- (ii) *If $T \in \mathcal{H}$ and $v \in (i_P^G(V))^K$ then $\varphi(T * v) = \mathcal{S}_M(T) * \varphi(v)$.*

Proof. Since $G = PK$ any element g in G can be written as $g = mnk$, where $k \in K, mn \in P$, and we can define

$$\psi : V^{K_M} \rightarrow (i_P^G(V))^K$$

by specifying that $\psi(v)(mnk) = \delta_N(m)^{1/2}\pi(m)v$. This is well defined precisely because $v \in V^{K_M}$. Now, (i) follows by computing that ψ is the inverse of φ .

For remainder of the proof let $\mathcal{S} = \mathcal{S}_M$. Let T_λ be the characteristic function of $K\lambda(\varpi)K$. Note that $\mathcal{S}(T_\lambda)$ is determined by its values on a set of coset representatives for M/K_M . We fix such a set. Using the Iwasawa decomposition $G = PK$, we may write (in analogy to (3))

$$(13) \quad K\lambda(\varpi)K = \bigcup_i \bigcup_j m_i n_i K$$

with the m_i chosen from the given set of coset representatives. If $m \notin K\lambda(\pi)K$, then obviously $\mathcal{S}(T_\lambda)(m) = 0$. Otherwise, $m = m_{j_0}$ for some m_{j_0} appearing in the decomposition (13). Let

$$n(i, j) := \#\{n_i \mid m_j n_i K \text{ appears in (13)}\}.$$

Then

$$\begin{aligned}
\mathcal{S}(T_\lambda)(m_{j_0}) &= \delta^{1/2}(m_{j_0}) \int_N T_\lambda(m_{j_0}n) dn \\
&= \delta^{1/2}(m_{j_0}) \sum_{i,j} T_\lambda(m_{j_0}n_i) \\
&= \delta^{1/2}(m_{j_0}) n(i, j_0),
\end{aligned}$$

since $\text{vol}(N/(K \cap N)) = 1$.

Let $f \in (\text{Ind}_P^G V)^K$. By (i), $\varphi(f) = f(1) = v \in V^{K_M}$. So, by the previous calculation,

$$\begin{aligned}
\mathcal{S}(T_\lambda) * v &= \int_M \mathcal{S}(T_\lambda)(m) \pi(m) v dm \\
&= \sum_j \mathcal{S}(T_\lambda)(m_j) \pi(m_j) v \\
&= \sum_j \delta^{1/2}(m_j) n(i, j) \pi(m_j) v.
\end{aligned}$$

On the other hand, since f is fixed by K ,

$$\begin{aligned}
\varphi(T_\lambda * f) &= (T_\lambda * f)(1) = \int_G T_\lambda(g) \sigma(g) f(1) dg \\
&= \sum_{i,j} f(1m_j n_i) \\
&= \sum_j n(i, j) \delta^{1/2}(m_j) \pi(m_j) v.
\end{aligned}$$

By Proposition 2.1, $\{T_\lambda \mid \lambda \in \Lambda_c^+\}$ forms a basis of \mathcal{H} . Therefore, (ii) is proved. \square

Lemma 2.3. *Let G be a reductive group with $P = MN$ the Levi decomposition of a parabolic. If V is any G -module, the map $V^K \rightarrow V_N^{K_M}$ is injective.*

Proof. For $I \subset G$ the Iwahori subgroup, Borel (see [1]) proved that $V^I \hookrightarrow V_N^{I_M}$. As the Jacquet functor is intertwining for the action of P , the image of V^K is clearly fixed by K_M . Since $V^K \subset V^I$, this gives the desired result. \square

Let G be a split reductive group defined over F . If $\chi : G \rightarrow \text{GL}_1$ is a character then $\chi^* : \mathbb{C}^\times \rightarrow \widehat{G}$ will denote the corresponding co-character. The group $\chi^*(\mathbb{C}^\times)$ is in the center of \widehat{G} . For example, if $G = \text{GL}_n$ and $\chi = \det$, then $\chi^*(z)$ is the scalar matrix in $\widehat{G} = \text{GL}_n(\mathbb{C})$.

Lemma 2.4. *Let $\chi : G \rightarrow \text{GL}_1$ be a character. Let V be a finite dimensional irreducible representation of \widehat{G} (i.e. a Hecke operator for G). Let m be a half integer. Then $\chi^*(q^m)$ acts on V as q^n for some half integer n . Let π be a representation of G . Then V acts on $\pi \otimes |\chi|^m$ as $q^n \cdot V$ acts on π .*

Lemma 2.5. *Let π be a representation of $G = G' \times G''$ obtained as the pullback of π' , a representation of G' . Let $s \in \widehat{G}''$ be the image of $\begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}$ under the principal $\text{SL}_2 \rightarrow \widehat{G}''$.*

For V a finite dimensional representation of \widehat{G} (i.e. a Hecke operator for G), write $V = \sum V' \otimes V''$, the restriction of V to $\widehat{G}' \times \widehat{G}''$. Then V acts on π as $\sum \text{Tr}_{V''}(s) V'$ acts on π' .

Proof. Since the Satake parameter of the trivial representation of a group G'' is s , the result is clear. \square

Lemma 2.6. *Let $\chi : G \rightarrow \mathrm{GL}_1$ be a character, and let $C = \chi^*(\mathbb{C}^\times) \subseteq \widehat{G}$. Let π' be a representation of GL_1 . Then $\pi = \pi' \circ \chi$ is a representation of G . Let $s \in \widehat{G}$ be the image of $\begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}$ under the principal $\mathrm{SL}_2 \rightarrow \widehat{G}$. For V a finite dimensional representation of \widehat{G} (i.e. a Hecke operator for G), write $V = \sum V' \otimes V''$, the restriction of V to $C \times \mathrm{SL}_2$. Then V acts on π as $\sum \mathrm{Tr}_{V''}(s)V'$ acts on π' .*

Proof. We prove this for $G = \mathrm{GL}_n$ and $\chi = \det$ is the determinant. In this case C is the center. It suffices to prove the statement for the fundamental representations $V_{\lambda_i} = V_i = \wedge^i \mathbb{C}^n$ where

$$\lambda_i = (\underbrace{1, \dots, 1}_{i \text{ times}}, 0, \dots, 0).$$

Since V_i is miniscule, the Satake isomorphism gives $\mathcal{S}(T_{\lambda_i}) = q^{i(n-i)/2}V_i$. Therefore, the action of $\pi(V_i)$ is

$$\begin{aligned} q^{i(i-n)/2} \int_{K\lambda_i(\varpi)K} \pi(g)dg &= q^{i(i-n)/2} \mathrm{vol}(K\lambda_i(\varpi)K) \pi'(\det(\lambda_i(\varpi))) \\ &= q^{i(i-n)/2} \mathrm{vol}(K\lambda_i(\varpi)K) \pi'(\varpi)^i. \end{aligned}$$

The center C acts on V_i by the character $z \mapsto z^i$. Note that

$$s = \begin{pmatrix} q^{(n-1)/2} & & & \\ & q^{(n-3)/2} & & \\ & & \ddots & \\ & & & q^{(1-n)/2} \end{pmatrix},$$

is the image of $\begin{pmatrix} q^{1/2} & \\ & q^{-1/2} \end{pmatrix}$ under the principal $\mathrm{SL}_2 \rightarrow \widehat{G}$. So, to complete the proof we just need to show that

$$q^{i(i-n)/2} \mathrm{vol}(K\lambda_i(\varpi)K) = \mathrm{Tr}_{V_i}(s).$$

This is immediate from the discussion in [2, Section 3] and the fact that V_i is miniscule. \square

Lemma 2.7. *Let G be a reductive group. Let $C_c^\infty(G)$ denote the space of smooth, compactly supported functions on G . This is a $G \times G$ module for the left and right action of G called the regular representation. On $C_c^\infty(G)^{K \times K}$ we have a matching of the Hecke algebras for the left and right action.*

Proof. This is obvious since $C_c^\infty(G)^{K \times K}$ is nothing else but the Hecke algebra itself. To be precise, since the left action on f in $C_c^\infty(G)$ is by $\lambda_g(f)(x) = f(g^{-1}x)$, a Hecke operator R , acting from the right, is matched with R^* defined by $R^*(x) = R(x^{-1})$. \square

Remark. Notice that this matching of Hecke operators, when considered as a matching of virtual representations, matches $V \in R(G)$ with its dual \widetilde{V} . In particular, if V is self-dual then it is matched with itself.

3. OUR GROUPS

3.1. Octonions. Let \mathbb{O} denote the non-associative division algebra of rank 8 over F . There is an F -linear anti-involution $x \mapsto \bar{x}$ on \mathbb{O} , hence norm and trace maps

$$\mathbb{N} : \mathbb{O} \rightarrow F \quad x \mapsto x\bar{x} = \bar{x}x, \quad \text{Tr} : \mathbb{O} \rightarrow F, \quad x \mapsto x + \bar{x}$$

satisfying

$$\mathbb{N}(x \cdot y) = \mathbb{N}(x)\mathbb{N}(y), \quad \text{Tr}(x \cdot y) = \text{Tr}(y \cdot x), \quad \text{Tr}(x \cdot (y \cdot z)) = \text{Tr}((x \cdot y) \cdot z).$$

On the set \mathbb{O}^0 of trace zero elements, we have $\bar{x} = -x$. The group G_2 is the automorphism group of \mathbb{O} .

The quadratic form $\mathbb{N} : \mathbb{O} \rightarrow F$ has signature $(4, 4)$, which means that \mathbb{O} has a basis $\{1, i, j, k, l, li, lj, lk\}$. (Note that $l^2 = 1$.) The following basis is particularly useful.

$$(14) \quad \begin{aligned} s_1 &= \frac{1}{2}(i + li), & s_2 &= \frac{1}{2}(j + lj), & s_3 &= \frac{1}{2}(k + lk), & s_4 &= \frac{1}{2}(1 + l), \\ t_1 &= \frac{1}{2}(i - li), & t_2 &= \frac{1}{2}(j - lj), & t_3 &= \frac{1}{2}(k - lk), & t_4 &= \frac{1}{2}(1 - l). \end{aligned}$$

The multiplication table for this basis is given in Table 2.

| | s_1 | s_2 | s_3 | t_1 | t_2 | t_3 | s_4 | t_4 |
|-------|--------|--------|--------|--------|--------|--------|-------|-------|
| s_1 | 0 | $-t_3$ | t_2 | s_4 | 0 | 0 | 0 | s_1 |
| s_2 | t_3 | 0 | $-t_1$ | 0 | s_4 | 0 | 0 | s_2 |
| s_3 | $-t_2$ | t_1 | 0 | 0 | 0 | s_4 | 0 | s_3 |
| t_1 | t_4 | 0 | 0 | 0 | s_3 | $-s_2$ | t_1 | 0 |
| t_2 | 0 | t_4 | 0 | $-s_3$ | 0 | s_1 | t_2 | 0 |
| t_3 | 0 | 0 | t_4 | s_2 | $-s_1$ | 0 | t_3 | 0 |
| s_4 | s_1 | s_2 | s_3 | 0 | 0 | 0 | s_4 | 0 |
| t_4 | 0 | 0 | 0 | t_1 | t_2 | t_3 | 0 | t_4 |

TABLE 2. Multiplication Table for Octonions

Remark. From this basis it is evident that a subspace $V \subset \mathbb{O}^0$ on which multiplication is trivial is at most 2-dimensional. (We call such a subspace a *null space* or a *null subspace*.) Indeed, let $\{i, j, k\} = \{1, 2, 3\}$. Then from the multiplication table we see that $s_i^\perp = \langle s_i, t_j, t_k \rangle$, and the null spaces of \mathbb{O}^0 which contain s_i are all of the form $\langle s_i, at_j + bt_k \rangle$ for fixed $a, b \in F$. Since G_2 acts transitively on (nonzero) elements of trace zero and norm zero, this phenomenon is generic.

3.1.1. Maximal parabolic subgroups in G_2 . They are described as the stabilizers of null subspaces $V \subset \mathbb{O}^0$. Let V_1 be spanned by s_1 and V_2 by s_1 and t_2 . Then $V_3 = V_1^\perp$ is spanned by s_1, t_2 and t_3 . Let $P_1 = M_1N_1$ and $P_2 = M_2N_2$ be the stabilizers of V_1 and V_2 , respectively. The Levi factor M_2 acts on V_2 . The choice of the basis in V_2 gives an isomorphism $M_2 \cong \text{GL}_2$. The Levi factor M_1 acts on V_3/V_1 and we have an isomorphism $M_1 \cong \text{GL}_2$. It is not difficult to see that $g \in \text{GL}_2 \cong M_1$ acts on V_1 by $\det(g)$. The set of all g in G_2 such that all s_i and t_i are eigenvectors is a maximal split torus T in G_2 .

The stabilizer of $s_4 - t_4$ in G_2 is a group isomorphic to SL_3 . Under the action of this group we have a decomposition

$$\mathbb{O}^0 = \langle s_1, s_2, s_3 \rangle \oplus \langle t_1, t_2, t_3 \rangle \oplus \langle s_4 - t_4 \rangle.$$

We can identify the stabilizer of $s_4 - t_4$ with SL_3 so that the action on $\langle s_1, s_2, s_3 \rangle$ is standard. The torus T of G_2 sits in SL_3 . In this way, we can represent elements in T by 3×3 matrices. For example, if α_l is a long root and α_s a short root perpendicular to α_l then, up to Weyl group conjugation,

$$(15) \quad \alpha_l^*(t) = \begin{pmatrix} t & & \\ & 1 & \\ & & t^{-1} \end{pmatrix} \text{ and } \alpha_s^*(t) = \begin{pmatrix} t & & \\ & t^{-2} & \\ & & t \end{pmatrix}.$$

3.2. Description of groups. Let $P = MN$ be a maximal parabolic subgroup of \mathbf{G} as in the table below. The group N is abelian, except in the case E_8 , where N has one-dimensional center Z . In order to give a uniform notation, let Z be trivial if N is abelian. Let d denote the dimension of N/Z . Let $C \cong \mathrm{GL}_1$ be the center of M . Fix an isomorphism $\lambda_* : \mathrm{GL}_1 \rightarrow C$ such that the adjoint action of $\lambda_*(z)$ on N/Z is given by multiplication by z . We have a dual pair $G_2 \times H \subset \mathbf{G}$ such that $Q = LU = H \cap P$ is a maximal parabolic of H . Let $N_0 \subseteq N/Z$ be the complement of \bar{U}/\bar{Z} under the invariant pairing induced by the Killing form. We fix an isomorphism of L with a classical group so that the action of $G_2 \times L$ on N_0 is isomorphic to $\mathbb{O}_0 \otimes F^n$, the space of n -tuples in \mathbb{O}_0 , and $h \in L$ acts on an n -tuple (x_1, \dots, x_n) by $(x_1, \dots, x_n)h^{-1}$. Since the scalar matrix z^{-1} in L acts on N/Z as z , the center of L coincides with the center of M . In the case of $\mathbf{G} = E_8$, let i be the isogeny character of GSp_6 .

| | | | | |
|--------------------|-----------------|-----------------|-----------------|------------------|
| \mathbf{G} | D_5 | E_6 | E_7 | E_8 |
| M | D_4 | D_5 | E_6 | E_7 |
| d | 8 | 16 | 27 | 56 |
| L | GL_1 | GL_2 | GL_3 | GSp_6 |
| $\delta_{\bar{U}}$ | $ \det $ | $ \det $ | $ \det ^2$ | $ i ^8$ |
| $\delta_{\bar{N}}$ | $ \det ^8$ | $ \det ^8$ | $ \det ^9$ | $ i ^{29}$ |

TABLE 3. Maximal parabolic subgroups

The last row of the table is the restriction of the character $\delta_{\bar{N}}$ to L . The group GSp_6 acts by the isogeny character i on \bar{Z} .

3.2.1. *Example:* $\mathbf{G} = E_7$. Let J_{27} be the 27-dimensional Jordan algebra over F given by

$$(16) \quad J_{27} = \left\{ A = \begin{pmatrix} a & z & \bar{y} \\ \bar{z} & b & x \\ y & \bar{x} & c \end{pmatrix} \middle| a, b, c \in F, \quad x, y, z \in \mathbb{O} \right\}.$$

The determinant on J_{27} gives an F -valued cubic form on J_{27} . The adjoint group \mathbf{G} has a maximal parabolic $P = MN$ such that

$$(17) \quad M \cong \{g \in \mathrm{GL}(J_{27}) \mid \det(g(A)) = \lambda(g) \det(A) \text{ for some similitude } \lambda(g) \in F^\times\},$$

a reductive group of type E_6 , and $N \cong J_{27}$ as M -modules. Moreover, $\bar{N} \cong J_{27}$, and the natural pairing of \bar{N} and N can be identified with the trace on J_{27} .

Evidently, $G_2 \subset M$ acts term by term on the elements of A , and since $g \in \mathrm{GL}_3$ acts via

$$(18) \quad g \cdot A = (\det g)^{-1} g A g^t,$$

the action of G_2 and GL_3 obviously commute, hence $G_2 \times \mathrm{GL}_3 \subset M$. As described in [4, Section 5], $G_2 \times \mathrm{PGSp}_6 \subset \mathbf{G}$ is a dual pair, and $Q = LU = \mathrm{PGSp}_6 \cap P$ where U can be identified with J_6 , the subalgebra of J_{27} consisting of (symmetric) matrices with entries in the field F , and $L \simeq \mathrm{GL}_3$ with action on $J_{27} \simeq N$ given by (18). Thus, the orthogonal complement N_0 of \bar{U} in N is identified with the subspace of J_{27} consisting of matrices with 0 on the diagonal and traceless octonions off the diagonal, that is, N_0 is identified with the set of triples (x, y, z) of traceless octonions. Moreover, $(g, h) \in G_2 \times \mathrm{GL}_3$ acts on (x, y, z) by $(gx, gy, gz)h^{-1}$. (The action of h follows from Cramer's rule.)

4. THE PROOF

4.1. The base case. Let \mathbf{V} be the minimal representation of D_4 . Let $G' = S_3$ be the group of permutations of 3 letters. Then S_3 acts on D_4 , by outer automorphisms, fixing G_2 . Since \mathbf{V} can be extended to a representation of a semi-direct product of D_4 and S_3 , we have a dual pair $G \times G' = G_2 \times S_3$ acting on \mathbf{V} . We let $\widehat{G}' = S_3$ and

$$r : S_3 \rightarrow G_2(\mathbb{C})$$

such that the centralizer C of $r(S_3)$ in $G_2(\mathbb{C})$ is $\mathrm{SO}(3) \subset \mathrm{SL}_3(\mathbb{C}) \subset G_2(\mathbb{C})$. (The group $\mathrm{SO}(3) \simeq \mathrm{PGL}_2(\mathbb{C})$ corresponds to the subregular unipotent orbit by the Jacobson-Morozov theorem.)

Let $K' = G'$. Then the Hecke algebra \mathcal{H}' is one-dimensional. With these choices, Theorem 1.1 asserts that the S_3 -invariants of the minimal representation of D_4 is the unramified representation π_{sr} of G_2 whose Satake parameter corresponds to the subregular orbit. This is proved in [3].

4.2. The general case. Assume that $\mathbf{G} \neq D_4$. Then we have a maximal parabolic $P = MN$ in \mathbf{G} and the corresponding maximal parabolic $Q = LU$ in H as in Table 3. For simplicity, let K be the maximal compact subgroup of G_2 , and K' that of H . Assume that we want to show matching of two operators T and T' . By Lemma 2.3,

$$\mathbf{V}^{K \times K'} \hookrightarrow r_{\bar{U}}(\mathbf{V})^{K \times K'_L}$$

where $K'_L = K' \cap L$. Thus Theorem 1.1 holds if we can show matching of T and $\mathcal{S}_L(T')$ on $r_{\bar{U}}(\mathbf{V})^{K \times K'_L}$. If $\mathbf{G} \neq E_8$, the unnormalized Jacquet functor $\mathbf{V}_{\bar{U}}$ was computed in [4]. In the context of the present work, we find it convenient to describe these results in terms of maximal parabolic subgroups of $L = \mathrm{GL}_n$.

4.2.1. Maximal parabolic subgroups in L . Recall that GL_n acts on F^n , the space of row vectors. For $m \leq n$ let $Q_m = L_m U_m$ be the stabilizer of the subspace consisting of the row vectors whose last $n - m$ entries are 0. Then $L_m = \mathrm{GL}_m \times \mathrm{GL}_{n-m}$. Let $g = (g_1, g_2) \in \mathrm{GL}_{n-m} \times \mathrm{GL}_m$. The modular character δ_{U_m} is

$$(19) \quad \delta_{U_m}(g) = |\det(g_1)|^{m-n} \cdot |\det(g_2)|^m.$$

Let s, t be a pair of real numbers. Let $C_c^\infty(\mathrm{GL}_m)[s, t]$ be the vector space $C_c^\infty(\mathrm{GL}_m)$ with an L_m -module structure defined by

$$(g_1, g_2) \cdot f(h) = |\det g_2|^t |\det g_1|^s f(hg_1).$$

We shall omit $[s, t]$ in the notation if $s = t = 0$.

Let $P_1 = M_1 N_1$ and $P_2 = M_2 N_2$ by the maximal parabolic subgroups of G_2 (as defined in Section 3.1.1.) Recall that $M_1 \cong \mathrm{GL}_2$ and $M_2 \cong \mathrm{GL}_2$. The modular characters are

$$(20) \quad \delta_{N_1}(g) = |\det(g)|^5 \text{ and } \delta_{N_2}(g) = |\det(g)|^3.$$

Let $g \in M_m$. The action of g on $C_c^\infty(\mathrm{GL}_m)[s, t]$ is given by

$$g \cdot f(h) = \begin{cases} f(\det g^{-1}h) & \text{if } m = 1, \\ f(g^{-1}h) & \text{if } m = 2. \end{cases}$$

With this notation in hand, we now describe the Jacquet module $r_{\bar{U}}(\mathbf{V})$ for each of our cases.

4.2.2. $\mathbf{G} = D_5$.

Proposition 4.1. *As a $G_2 \times \mathrm{GL}_1$ -module, $r_{\bar{U}}(\mathbf{V})$ has a filtration with two successive subquotients*

- (1) $i_{P_1}^{G_2}(C_c^\infty(\mathrm{GL}_1))$.
- (2) $\mathbf{V}(M) \otimes |\det|^{\frac{1}{2}} \oplus 1 \otimes |\det|^{\frac{5}{2}}$.

Here $\mathbf{V}(M)$ is the minimal representation of M/C .

Proof. This is simply a normalized version of Proposition 2.3 of [4]. Indeed, $\mathbf{V}_{\bar{U}}$ has a filtration with two successive quotients

- $\mathrm{Ind}_{P_1}^{G_2}(C_c^\infty(\mathrm{GL}_1)) \otimes |\det|^3$.
- $\mathbf{V}(M) \otimes |\det| \oplus 1 \otimes |\det|^3$.

where induction is not normalized. Since $r_{\bar{U}}(\mathbf{V}) = \mathbf{V}_{\bar{U}} \otimes \delta_{\bar{U}}^{-\frac{1}{2}}$, and $\delta_{\bar{U}} = |\det|$ by Table 3, $r_{\bar{U}}(\mathbf{V})$ has a filtration with two successive quotients

- $\mathrm{Ind}_{P_1}^{G_2}(C_c^\infty(\mathrm{GL}_1)) \otimes |\det|^{\frac{5}{2}}$.
- $\mathbf{V}(M) \otimes |\det|^{\frac{1}{2}} \oplus 1 \otimes |\det|^{\frac{5}{2}}$.

and (2) follows. Since, for any s , $C_c^\infty(\mathrm{GL}_m) \cong C_c^\infty(\mathrm{GL}_m) \otimes |\det|^s$ as $\mathrm{GL}_m \times \mathrm{GL}_m$ -modules, we can replace $C_c^\infty(\mathrm{GL}_1)$ in the first bullet by $C_c^\infty(\mathrm{GL}_1) \otimes |\det|^{-\frac{5}{2}}$. By (10), this normalizes the induction for G_2 and, at the same time, removes the character $|\det|^{\frac{5}{2}}$ of GL_1 . Hence (1) follows. \square

Proof of Theorem 1.1 in case $\mathbf{G} = D_5$. The dual group of $H = \mathrm{PGL}_2$ is $\mathrm{SL}_2(\mathbb{C})$. The map $r : \mathrm{SL}_2(\mathbb{C}) \rightarrow G_2(\mathbb{C})$ corresponds to a long root α_l of G_2 : $r(\mathrm{SL}_2(\mathbb{C})) = \mathrm{SL}_{2,l}(\mathbb{C}) \subset G_2(\mathbb{C})$. The centralizer C of $\mathrm{SL}_{2,l}(\mathbb{C})$ in $G_2(\mathbb{C})$ is $\mathrm{SL}_{2,s}(\mathbb{C})$ corresponding to a short root α_s perpendicular to α_l .

Let V be a finite dimensional representation of $G_2(\mathbb{C})$ and T_2 the corresponding Hecke operator for G_2 . Let

$$(21) \quad s = \alpha_s^*(q^{1/2}) = \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix} \in \mathrm{SL}_{2,s}(\mathbb{C}).$$

If the restriction of V to $\mathrm{SL}_{2,l}(\mathbb{C}) \times \mathrm{SL}_{2,s}(\mathbb{C})$ is $\sum V' \otimes V''$, we define T_1 as corresponding to $\sum \mathrm{Tr}_{V''}(s)V'$. We want to show that T_2 matches with $\mathcal{S}_L(T_1)$ on $r_{\bar{U}}(\mathbf{V})$. Since \widehat{L} is the torus $\alpha_l^*(\mathbb{C}^\times) \subseteq \mathrm{SL}_{2,l}(\mathbb{C})$, the operator $\mathcal{S}_L(T_1)$ corresponds to the representation $\sum \mathrm{Tr}_{V''}(s)V'$ of $\alpha_l^*(\mathbb{C}^\times)$ obtained by restricting each V' to the torus $\alpha_l^*(\mathbb{C}^\times)$.

First, we show matching on (1) in Proposition 4.1. By Lemma 2.2, we need to show matching of $\mathcal{S}_{M_1}(T_2)$ and $\mathcal{S}_L(T_1)$ on $C_c^\infty(\mathrm{GL}_1)$. The operator $\mathcal{S}_{M_1}(T_2)$ corresponds to the restriction of V to \widehat{M}_1 . Since M_1 corresponds to a long root, $\widehat{M}_1 \simeq \mathrm{GL}_{2,s}(\mathbb{C})$. The center of $\mathrm{GL}_{2,s}(\mathbb{C})$ is the torus $\alpha_l^*(\mathbb{C}^\times) \subseteq \mathrm{SL}_{2,l}(\mathbb{C})$. Let s be as in (21) and let $V = \sum V' \otimes V''$ be the restriction of V to $\mathrm{SL}_{2,l}(\mathbb{C}) \times \mathrm{SL}_{2,s}(\mathbb{C})$, as before. By Lemma 2.6, $\mathcal{S}_{M_1}(T_2)$ acts on $C_c^\infty(\mathrm{GL}_1)$ as the Hecke operator for GL_1 that corresponds to the representation $\sum \mathrm{Tr}_{V''}(s)V'$ of $\alpha_l^*(\mathbb{C}^\times)$, the center of $\mathrm{GL}_{2,s}(\mathbb{C})$. In particular, this is the same GL_1 -operator as $\mathcal{S}_L(T_1)$. Matching now follows from Lemma 2.7.

Using the previously proved base case, matching on (2) in Proposition 4.1 reduces to a simple check on two $G_2 \times L$ modules: $\pi_{sr} \otimes |\det|^{\frac{1}{2}}$ and $1 \otimes |\det|^{\frac{5}{2}}$. \square

4.2.3. $\mathbf{G} = E_6$.

Proposition 4.2. *As a $G_2 \times \mathrm{GL}_2$ -module, $r_{\bar{U}}(\mathbf{V})$ has a filtration with three successive subquotients*

- (1) $i_{P_2}^{G_2}(C_c^\infty(\mathrm{GL}_2))$.
- (2) $i_{P_1 \times Q_1}^{G_2 \times \mathrm{GL}_2}(C_c^\infty(\mathrm{GL}_1)[- \frac{1}{2}, 1])$.
- (3) $\mathbf{V}(M) \otimes |\det|^{\frac{1}{2}} \oplus 1 \otimes |\det|^{\frac{3}{2}}$.

Here $\mathbf{V}(M)$ is the minimal representation of M/C .

This is a normalized version of Theorem 4.3 of [4] which states that $\mathbf{V}_{\bar{U}}$ has a filtration with three successive quotients

- $\mathrm{Ind}_{P_2}^{G_2}(C_c^\infty(\mathrm{GL}_2)) \otimes |\det|^2$.
- $\mathrm{Ind}_{P_1 \times Q_1}^{G_2 \times \mathrm{GL}_2}(C_c^\infty(\mathrm{GL}_1)) \otimes |\det|^2$.
- $\mathbf{V}(M) \otimes |\det| \oplus 1 \otimes |\det|^2$.

Proof of Theorem 1.1 in case $\mathbf{G} = E_6$. The dual group of $H = \mathrm{PGL}_3$ is $\mathrm{SL}_3(\mathbb{C})$. We have an inclusion $r : \mathrm{SL}_3(\mathbb{C}) \rightarrow G_2(\mathbb{C})$ where $\mathrm{SL}_3(\mathbb{C}) \subset G_2(\mathbb{C})$ is given by the long roots.

Let V be a finite dimensional representation of $G_2(\mathbb{C})$ and T_2 the corresponding Hecke operator for G_2 . We restrict V to $\mathrm{SL}_3(\mathbb{C})$, and let T_2 be the corresponding Hecke operator for PGL_3 . We want to show that T_2 matches with $\mathcal{S}_L(T_1)$ on $r_{\bar{U}}(\mathbf{V})$.

First, we consider matching on $\mathbf{V}(M) \otimes |\det|^{\frac{1}{2}}$. The dual group of $L = \mathrm{GL}_2$ is $\mathrm{GL}_{2,l}(\mathbb{C}) \subseteq \mathrm{SL}_3(\mathbb{C})$. The group L acts on $\mathbf{V}(M)$ by its quotient PGL_2 . The dual group of PGL_2 is $\mathrm{SL}_{2,l}(\mathbb{C})$. The operator $\mathcal{S}_L(T_1)$ acts on $\mathbf{V}(M)$ as the Hecke operator for PGL_2 that corresponds to the restriction of V to $\mathrm{SL}_{2,l}(\mathbb{C})$. We need to take into account the twist by $|\det|^{\frac{1}{2}}$. Let χ be the determinant character of L . Let $\chi^* : \mathbb{C}^\times \rightarrow \mathrm{GL}_{2,l}(\mathbb{C})$ be the corresponding co-character. Note that $\chi^* = \alpha_s^*$. Let $s = \alpha_s^*(q^{1/2}) \in \mathrm{SL}_{2,s}(\mathbb{C})$. Let $V = \sum V' \otimes V''$ be the restriction of V to $\mathrm{SL}_{2,l}(\mathbb{C}) \times \mathrm{SL}_{2,s}(\mathbb{C})$. Let T be the Hecke operator for PGL_2 that corresponds to the representation $\sum \mathrm{Tr}_{V''}(s)V'$ of $\mathrm{SL}_{2,l}(\mathbb{C})$. By Lemma 2.4, $\mathcal{S}_L(T_1)$ acts on $\mathbf{V}(M) \otimes |\det|^{\frac{1}{2}}$ as T acts on $\mathbf{V}(M)$. But T is matched with T_2 on $\mathbf{V}(M)$, by the case $\mathbf{G} = D_5$.

To prove matching on (1) in Proposition 4.2 it suffices to show that $\mathcal{S}_{M_2}(T_2)$ and $\mathcal{S}_L(T_1)$ are matching on $C_c^\infty(\mathrm{GL}_2)$, by Lemma 2.2. Since M_2 corresponds to a short root, $\widehat{M}_2 \simeq \mathrm{GL}_{2,l}(\mathbb{C})$. Thus, the dual groups of M_2 and L are conjugate in $G_2(\mathbb{C})$ and matching follows from Lemma 2.7. (See also the remark following Lemma 2.7.)

Matching on (2) is similar to (1), albeit slightly more complicated to write down, so we omit details. \square

4.2.4. $\mathbf{G} = E_7$.

Proposition 4.3. *As a $G_2 \times \mathrm{GL}_3$ -module, $r_{\bar{U}}(\mathbf{V})$ has a filtration with three successive subquotients*

- (1) $i_{P_2 \times Q_2}^{G_2 \times \mathrm{GL}_3}(C_c^\infty(\mathrm{GL}_2))$.
- (2) $i_{P_1 \times Q_1}^{G_2 \times \mathrm{GL}_3}(C_c^\infty(\mathrm{GL}_1)[- \frac{1}{2}, \frac{1}{2}])$.
- (3) $\mathbf{V}(M) \oplus 1 \otimes |\det|$.

Here $\mathbf{V}(M)$ is the minimal representation of M/C .

This is a normalized version of Theorem 5.3 of [4] which states that $\mathbf{V}_{\bar{U}}$ has a filtration with three successive quotients

- $\mathrm{Ind}_{P_2 \times Q_2}^{G_2 \times \mathrm{GL}_3}(C_c^\infty(\mathrm{GL}_2)) \otimes |\det|^2$.
- $\mathrm{Ind}_{P_1 \times Q_1}^{G_2 \times \mathrm{GL}_3}(C_c^\infty(\mathrm{GL}_1)) \otimes |\det|^2$.
- $\mathbf{V}(M) \otimes |\det| \oplus 1 \otimes |\det|^2$.

Proof of Theorem 1.1 in case $\mathbf{G} = E_7$. The dual group of $H = \mathrm{PGSp}_6$ is $\mathrm{Spin}_7(\mathbb{C})$. Let \mathbb{Z}^3 be the root lattice of $\mathrm{Spin}_7(\mathbb{C})$ so that the short roots correspond to the standard basis vectors e_1, e_2, e_3 in \mathbb{Z}^3 . Let V_8 be the spin representation of $\mathrm{Spin}_7(\mathbb{C})$. The weights of V_8 are $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$. Under the action of $\widehat{L} \cong \mathrm{GL}_3(\mathbb{C})$ the spin representation decomposes as

$$V_8 = V_1 \oplus V_3 \oplus V_3^* \oplus V_1^*$$

where V_3 is the standard representation of $\mathrm{GL}_3(\mathbb{C})$ and V_1 is the determinant character. The weights these 4 summands are (x, y, z) such that $x + y + z = \frac{3}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{3}{2}$, respectively. We have an injection

$$r : G_2(\mathbb{C}) \rightarrow \mathrm{Spin}_7(\mathbb{C})$$

where $G_2(\mathbb{C})$ is defined as the stabilizer of a non-zero vector in V_1 , for example. In particular, $G_2(\mathbb{C}) \cap \mathrm{GL}_3(\mathbb{C}) = \mathrm{SL}_3(\mathbb{C})$.

Let V be a finite dimensional representation of $\mathrm{Spin}_7(\mathbb{C})$ and T_1 the corresponding Hecke operator for PGSp_6 . We restrict V to $G_2(\mathbb{C})$, and let T_2 be the corresponding Hecke operator for G_2 . We want to show that T_2 matches with $\mathcal{S}_L(T_1)$ on $r_{\bar{U}}(\mathbf{V})$. Matching on (3) in Proposition 4.3 trivially follows from the previously proved case $\mathbf{G} = E_6$.

To prove matching on (1) it suffices to show that $\mathcal{S}_{M_2}(T_2)$ matches with $\mathcal{S}_{L_2} \circ \mathcal{S}_L(T_1)$ on $C_c^\infty(\mathrm{GL}_2)$. Since M_2 corresponds to a short root, $\widehat{M}_2 \simeq \mathrm{GL}_{2,l}(\mathbb{C}) \subseteq \mathrm{SL}_3(\mathbb{C})$ where, for definiteness, $g \in \mathrm{GL}_{2,l}(\mathbb{C})$ sits in $\mathrm{SL}_3(\mathbb{C})$ as a block diagonal matrix $(g, \det g^{-1})$. The operator $\mathcal{S}_{M_2}(T_2)$ corresponds to the restriction of V to $\mathrm{GL}_{2,l}(\mathbb{C})$. Since $L_2 = \mathrm{GL}_2 \times \mathrm{GL}_1$ and GL_1 acts trivially on $C_c^\infty(\mathrm{GL}_2)$, the operator $\mathcal{S}_{L_2} \circ \mathcal{S}_L(T_1)$ acts on $C_c^\infty(\mathrm{GL}_2)$ as the Hecke operator for GL_2 that corresponds to the restriction of V to $\mathrm{GL}_2(\mathbb{C})$, the first factor of

$$\widehat{L}_2 = \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C}) \subseteq \mathrm{GL}_3(\mathbb{C}) \subseteq \mathrm{Spin}_7(\mathbb{C}).$$

This $\mathrm{GL}_2(\mathbb{C})$ is conjugated to $\mathrm{GL}_{2,l}(\mathbb{C})$ in Spin_7 by the reflection corresponding to the short root e_3 . Matching on (1) now follows from Lemma 2.7.

To prove matching on (2) it suffices to show that $\mathcal{S}_{M_1}(T_2)$ matches with $\mathcal{S}_{L_1} \circ \mathcal{S}_L(T_1)$ on $C_c^\infty(\mathrm{GL}_1)$. Since M_1 corresponds to a short root, $\widehat{M}_1 \simeq \mathrm{GL}_{2,s}(\mathbb{C})$. The center of $\mathrm{GL}_{2,s}(\mathbb{C})$ is the torus $\alpha_l^*(\mathbb{C}^\times) \subseteq \mathrm{SL}_{2,l}(\mathbb{C})$. By Lemma 2.6, $\mathcal{S}_{M_2}(T_2)$ acts on $C_c^\infty(\mathrm{GL}_1)$ as the Hecke operator for GL_1 that corresponds to the restriction of V to $\alpha_l^*(\mathbb{C}^\times)$, weighted by the eigenvalues of $\alpha_s^*(q^{1/2})$. On the other hand, the operator $\mathcal{S}_{L_1} \circ \mathcal{S}_L(T_1)$ acts on $C_c^\infty(\mathrm{GL}_1)$ as the Hecke operator for GL_1 that corresponds to the restriction of V to $\mathrm{GL}_1(\mathbb{C})$, the first factor of

$$\widehat{L}_1 = \mathrm{GL}_1(\mathbb{C}) \times \mathrm{GL}_2(\mathbb{C}) \subseteq \mathrm{GL}_3(\mathbb{C}) \subseteq \mathrm{Spin}_7(\mathbb{C}),$$

weighted by the eigenvalues of

$$s = \begin{pmatrix} q^{-\frac{1}{2}} & & \\ & q^{\frac{1}{2}} & \\ & & q^{\frac{1}{2}} \end{pmatrix} \cdot \begin{pmatrix} 1 & & \\ & q^{\frac{1}{2}} & \\ & & q^{-\frac{1}{2}} \end{pmatrix} \in \mathrm{GL}_3(\mathbb{C}) \subseteq \mathrm{Spin}_7(\mathbb{C}),$$

where the first matrix in the above product reflects the twisting $[-\frac{1}{2}, \frac{1}{2}]$ in (2), and the second comes from Lemma 2.5. The pairs $(\alpha_l^*(\mathbb{C}^\times), \alpha_s^*(q^{1/2}))$ (see (15)) and $(\mathrm{GL}_1(\mathbb{C}), s)$ are conjugated in $\mathrm{Spin}_7(\mathbb{C})$ by the reflection corresponding to the short root e_3 . Matching on (2) now follows from Lemma 2.7. \square

4.2.5. $\mathbf{G} = E_8$. We start with a description of maximal parabolic subgroups in GSp_{2n} . This is a group of isogenies of a symplectic form (\cdot, \cdot) on a $2n$ dimensional space. Let $e_1, \dots, e_n, f_1, \dots, f_n$ be a symplectic basis, i.e.

$$(e_i, f_j) = -(f_j, e_i) = \delta_{ij} \text{ and } (e_i, e_j) = (f_i, f_j) = 0.$$

Using this basis we identify the symplectic space with F^{2n} , the space of $2n$ -tuples. We identify GSp_{2n} with the group of $2n \times 2n$ matrices action on F^{2n} from the right, and preserving the symplectic form, up to an isogeny character :

$$(vg^{-1}, ug^{-1}) = i(g)^{-1}(v, u).$$

For every $m \leq n$, let $Q_m = L_m U_m$ be the subgroup of GSp_{2n} preserving the subspace spanned by e_1, \dots, e_m . Then $Q_m \cong \mathrm{GL}_m \times \mathrm{GSp}_{2(n-m)}$ where $g = (g_1, g_2) \in \mathrm{GL}_m \times \mathrm{GSp}_{2(n-m)}$ acts as follows: $e_i g_1^{-1}$, for $1 \leq i \leq m$, $f_i g_1^{-1}$ for $1 \leq i \leq m$ and trivially on other basis elements; g_2^{-1} is an isogeny on the symplectic space spanned by e_{m+1}, \dots, e_n and f_{m+1}, \dots, f_n , $e_i g_2^{-1} = e_i$, for $1 \leq i \leq m$, and $f_i g_2^{-1} = i(g_2)^{-1} \cdot f_i$ for $1 \leq i \leq m$. The modular character δ_{U_m} is

$$(22) \quad \delta_{U_m}(g) = |\det(g_1)|^{-(2n-m+1)} |i(g_2)|^{\frac{m(2n-m+1)}{2}}$$

Let s, t be a pair of real numbers. Let $C_c^\infty(\mathrm{GL}_m)[s, t]$ be an L_m -module defined as follows. As a vector space $C_c^\infty(\mathrm{GL}_m)[s, t] = C_c^\infty(\mathrm{GL}_m)$, and $(g_1, g_2) \in \mathrm{GL}_m \times \mathrm{GSp}_{2(n-m)}$ acts by

$$(g_1, g_2) \cdot f(h) = |\det g_1|^s |i(g_2)|^t f(hg_1).$$

Proposition 4.4. *As a $G_2 \times \mathrm{GSp}_6$ -module, $r_{\bar{V}}(\mathbf{V})$ has a filtration with three successive subquotients*

- (1) $i_{P_2 \times Q_2}^{G_2 \times \mathrm{GSp}_6}(C_c^\infty(\mathrm{GL}_2)[1, -\frac{3}{2}])$.
- (2) $i_{P_1 \times Q_1}^{G_2 \times \mathrm{GSp}_6}(C_c^\infty(\mathrm{GL}_1)[\frac{1}{2}, -\frac{1}{2}])$.
- (3) $\mathbf{V}(M) \otimes |i|^{-1} \oplus 1 \otimes |i|$.

Here $\mathbf{V}(M)$ is the minimal representation of M/C .

The proof of this proposition is given in Section 5.

Proof of Theorem 1.1 in case $\mathbf{G} = E_8$. The map $r : G_2(\mathbb{C}) \rightarrow F_4(\mathbb{C})$ is described as follows. A split, simply connected group G_{sc} of type E_6 can be realized as a subgroup of $\mathrm{GL}(J_{27})$ fixing $\det : J_{27} \rightarrow F$. As described in 3.2.1, there is a dual pair $G_2 \times \mathrm{SL}_3 \subseteq G_{sc}$. The group F_4 is the subgroup of G_{sc} consisting of elements fixing the identity matrix in J_{27} . It is easy to check that $(G_2 \times \mathrm{SL}_3) \cap F_4 = G_2 \times \mathrm{SO}_3$. This defines r , and the centralizer C of $r(G_2(\mathbb{C}))$ is $\mathrm{SO}_3(\mathbb{C})$. The dual group \widehat{L} of $L = \mathrm{GSp}_6$ is a Levi factor of type B_3 . Let $i^* : \mathbb{C}^\times \rightarrow \widehat{L}$ be the co-character corresponding to the isogeny character i of GSp_6 . Then $i^*(\mathbb{C}^\times)$ is the center of \widehat{L} . We can conjugate $G_2(\mathbb{C})$ in $F_4(\mathbb{C})$ so that

$$G_2(\mathbb{C}) \subseteq \mathrm{Spin}_7(\mathbb{C}) = [\widehat{L}, \widehat{L}].$$

In this way, $i^*(\mathbb{C}^\times)$ is a maximal torus in $\mathrm{SO}_3(\mathbb{C})$, the centralizer of $G_2(\mathbb{C})$. The image of $\begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$ in $\mathrm{SO}_3(\mathbb{C})$ is $s = i^*(q)$.

Let V be a finite dimensional representation of $F_4(\mathbb{C})$ and T_1 the corresponding Hecke operator for F_4 . If $V = \sum V' \otimes V''$ is the restriction of V to $G_2(\mathbb{C}) \times \mathrm{SO}_3(\mathbb{C})$, we have defined T_2 as corresponding to $\sum \mathrm{Tr}_{V''}(s)V'$. We want to show that T_2 matches with $\mathcal{S}_L(T_1)$ on $r_{\bar{U}}(\mathbf{V})$. The operator $\mathcal{S}_L(T_1)$ corresponds to the representation \widehat{L} obtained by restricting V to \widehat{L} . We now show matching on (3). Let $V = \sum V_n$ be the restriction of V to $\mathrm{Spin}_7(\mathbb{C}) = [\widehat{L}, \widehat{L}]$, where $i^*(q)$ act as q^n on V_n . Let T be the Hecke operator for PGSp_6 that corresponds to the representation $\sum q^{-n}V_n$ of $\mathrm{Spin}_7(\mathbb{C})$. Then, by Lemma 2.4, $\mathcal{S}_L(T_1)$ acts on $\mathbf{V}(M) \otimes |i|^{-1}$ as T acts on $\mathbf{V}(M)$. Matching on $\mathbf{V}(M) \otimes |i|^{-1}$ now follows from the previously proved case $\mathbf{G} = E_7$. Let $s_p \in G_2(\mathbb{C})$ be the image of $\begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$ under the principal

$$f : \mathrm{SL}_2(\mathbb{C}) \rightarrow G_2(\mathbb{C}).$$

Then T_2 acts on $1 \otimes |i|$ as the scalar $\sum \mathrm{Tr}_{V''}(s) \cdot \mathrm{Tr}_{V'}(s_p)$. Under the inclusion $G_2(\mathbb{C}) \subseteq \mathrm{Spin}_7(\mathbb{C})$, the composite $f : \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{Spin}_7(\mathbb{C})$ is the principal $\mathrm{SL}_2(\mathbb{C})$ in $\mathrm{Spin}_7(\mathbb{C})$. By Lemma 2.4, $\mathcal{S}_L(T_1)$ acts on $1 \otimes |i|$ as the scalar $\sum q^n \mathrm{Tr}_{V_n}(s_p)$. Since

$$\sum q^n \mathrm{Tr}_{V_n}(s_p) = \sum \mathrm{Tr}_{V''}(s) \cdot \mathrm{Tr}_{V'}(s_p),$$

matching is now proved on (3). The remaining cases are similar to $\mathbf{G} = E_7$, so we leave them as an exercise. \square

5. A JACQUET MODULE FOR E_8

In this section we prove Proposition 4.4. In this case N is a Heisenberg group. A starting point is the following (Theorem 6.1 [4]).

Proposition 5.1. *Let Ω be the M -orbit of the highest weight vector in N/Z . We have the following exact sequence of \bar{P} -modules,*

$$0 \rightarrow C_c^\infty(\Omega) \rightarrow \mathbf{V}_{\bar{Z}} \rightarrow \mathbf{V}_{\bar{N}} \rightarrow 0.$$

The action of \bar{P} on $f \in C_c^\infty(\Omega)$ is given by:

- For every $\bar{n} \in \bar{N}$

$$\Pi(\bar{n})f(x) = \psi(\langle x, \bar{n} \rangle)f(x).$$

- For every $m \in M$

$$\Pi(m)f(x) = |i(m)|^5 f(m^{-1}xm).$$

Here ψ is a non-trivial additive character, $\langle \cdot, \cdot \rangle$ is a pairing between N/Z and \bar{N}/\bar{Z} induced by the Killing form, and $i : M \rightarrow \text{GL}_1$ is the character obtained by acting on \bar{Z} . Moreover, $\mathbf{V}_{\bar{N}} \cong \mathbf{V}(M) \otimes |i|^3 \oplus |i|^5$ where $\mathbf{V}(M)$ is the minimal representation of M with the center acting trivially.

By Section 7 in [4] we have an identification of vector spaces

$$N/Z \cong \bar{N}/\bar{Z} \cong F \oplus J_{27} \oplus J_{27} \oplus F$$

so that the pairing $\langle \cdot, \cdot \rangle$ is given by

$$\langle (a, B, C, d), (a', B', C', d') \rangle = aa' + \text{Tr}(BB') + \text{Tr}(CC') + dd'.$$

Under these identifications, the action of G_2 on N/Z is the obvious one, and the centralizer of G_2 in \bar{N}/\bar{Z} is

$$\bar{U}/\bar{Z} \cong F \oplus J_6 \oplus J_6 \oplus F$$

where J_6 is the space of 3×3 matrices with coefficients in F . It follows that the orthogonal complement of \bar{U}/\bar{Z} in N/Z can be identified by $J^0 \oplus J^0$ where J^0 is the set of $B \in J$ of the form

$$B = \begin{pmatrix} 0 & z & -y \\ -z & 0 & x \\ y & -x & 0 \end{pmatrix}$$

with $x, y, z \in \mathbb{O}^0$. Given this, we may denote an element $(B, B') \in J^0 \oplus J^0$ by a six-tuple

$$(u, u') = ((x, y, z), (x', y', z'))$$

of traceless octonions. The action of $G_2 \times \text{GSp}_6$ on these elements is simple to describe. First, $g \in G_2$ acts on every component of the six-tuple (u, u') from the left, and $h \in \text{GSp}_6$ acts by $(u, u')h^{-1}$. In this way, the character i of M restricts to the isogeny character of GSp_6 . We highlight the action of certain subgroups: $\text{SL}_2 \times \text{SL}_2 \times \text{SL}_2 \subseteq \text{Sp}_6$ acting on the pairs (x, x') , (y, y') and (z, z') respectively in the obvious way, and $h \in \text{GL}_3$ by

$$(uh^{-1}, u'h^t).$$

Note that GSp_6 preserves, up to the isogeny character,

$$J^0 \oplus J^0 \rightarrow \wedge^2 \mathbb{O}^0 \quad ((x, y, z), (x', y', z')) \mapsto x \wedge x' + y \wedge y' + z \wedge z'.$$

Let Ω_0 be the intersection of Ω with $J^0 \oplus J^0$, the orthogonal complement of \bar{U}/\bar{Z} . It follows, from Proposition 5.1, that there is an exact sequence of $G_2 \times \text{GSp}_6$ -modules

$$0 \rightarrow C_c^\infty(\Omega_0) \rightarrow \mathbf{V}_{\bar{U}} \rightarrow \mathbf{V}_{\bar{N}} \rightarrow 0,$$

where $(g, h) \in G_2 \times \text{GSp}_6$ acts on $f \in C_c^\infty(\Omega_0)$ by

$$\Pi(g, h)f(x) = |i(h)|^5 f(g^{-1}xh).$$

In order to understand $C_c^\infty(\Omega_0)$, we need to compute $G_2 \times \text{GSp}_6$ -orbits on Ω_0 .

Proposition 5.2. *The set Ω_0 consists of pairs of $((x, y, z), (x', y', z')) \in J^0 \times J^0$ such that $\langle x, y, z, x', y', z' \rangle$ is a non-zero null subspace of \mathbb{O}^0 , and such that*

$$x \wedge x' + y \wedge y' + z \wedge z' = 0.$$

Moreover, Ω_0 consists of two $G_2 \times \text{GSp}_6$ orbits Ω_1 and Ω_2 where

$$\Omega_m = \{((x, y, z), (x', y', z')) \in \Omega \mid \dim(\langle x, y, z, x', y', z' \rangle) = m\}.$$

Proof. If $(0, B, B', 0) \in \Omega_0$ then, by Lemma 7.5 in [4], B is a rank one matrix, $B^2 = \text{Tr}(B)B$. Since $\text{Tr}(B) = 0$, we have $B^2 = 0$, and this is equivalent to

$$x^2 = y^2 = z^2 = xy = yz = 0,$$

i.e. the entries of B span a null subspace of \mathbb{O}^0 . Acting by $\text{SL}_2 \times \text{SL}_2 \times \text{SL}_2$, we can replace x, y and z (all or some) by x', y' and z' , respectively. Hence $\langle x, y, z, x', y', z' \rangle$ is a non-zero null subspace of \mathbb{O}^0 . If the dimension of this null space is 1. Without loss of generality, we can assume that $x \neq 0$. Then (u, u') is in the GSp_6 orbit of $((x, 0, 0), (0, 0, 0))$. Since G_2 acts transitively on 1-dimensional null subspaces, we have one $G_2 \times \text{GSp}_6$ orbit. Assume that the dimension is 2. Without loss of generality, we can assume that x and z are a basis of this space. Using the action of GL_3 we can arrange that $y = 0$. Since

$$x' = ax + cz, y' = ex + fz, z' = bz + dx$$

for some $a, b, c, d, e, f \in F$,

$$x \wedge (ax + cz) + 0 \wedge (ex + fz) + z \wedge (bz + cx) = (c - d)(x \wedge z)$$

and this is 0 if and only if $c = d$. If $c = d$ then it is not too difficult to see that (u, u') is in the GSp_6 orbit of $((x, 0, z), (0, 0, 0))$. Since G_2 acts transitively on 2-dimensional null subspaces, we have one $G_2 \times \text{GSp}_6$ orbit. Thus, to finish the proof we must show that $c = d$. This is done in Lemma 5.3, using that $B' = A \times B$ (the cross product) for some $A \in J_{27}$, by Lemma 7.5 in [4]. \square

Lemma 5.3. *Suppose that $x, z \in \mathbb{O}^0$ be linearly independent such that $x^2 = z^2 = xz = 0$. Let $x_1, y_1, z_1 \in \mathbb{O}^0$, and set*

$$A = A_0 + \begin{pmatrix} 0 & z_1 & -y_1 \\ -z_1 & 0 & x_1 \\ y_1 & -x_1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & z & 0 \\ -z & 0 & x \\ 0 & -x & 0 \end{pmatrix}$$

where $A_0 \in J_6$. If

$$A \times B = \begin{pmatrix} 0 & z' & -y' \\ -z' & 0 & x' \\ y' & -x' & 0 \end{pmatrix}$$

is such that $x', y', z' \in \mathbb{O}^0$ then $x', y', z' \in Fx + Fz$. Moreover,

$$(23) \quad x' = bx + cz, \quad \text{and} \quad z' = az + cx$$

for some constants $a, b, c \in F$.

Proof. Since G_2 acts transitively on the set of 2-dimensional null spaces of \mathbb{O}^0 , by the G_2 action (which commutes with the cross product), we may assume that $x = s_1$ and $z = t_2$,

and let

$$\begin{aligned}x_1 &= a_1^x s_1 + a_2^x s_2 + a_3^x s_3 + b_1^x t_1 + b_2^x t_2 + b_3^x t_3 + c^x (s_4 - t_4), \\y_1 &= a_1^y s_1 + a_2^y s_2 + a_3^y s_3 + b_1^y t_1 + b_2^y t_2 + b_3^y t_3 + c^y (s_4 - t_4), \\z_1 &= a_1^z s_1 + a_2^z s_2 + a_3^z s_3 + b_1^z t_1 + b_2^z t_2 + b_3^z t_3 + c^z (s_4 - t_4)\end{aligned}$$

where the elements $s_i, t_j \in \mathbb{O}$ are the basis elements given in (14).

The cross product is given by

$$A \times B = A \circ B - \frac{1}{2}A \operatorname{Tr} B - \frac{1}{2}B \operatorname{Tr} A + \frac{1}{2}(\operatorname{Tr} A \operatorname{Tr} B - \operatorname{Tr}(A \circ B)).$$

From this a simple calculation shows that if $A = A_0$ then the condition (23) is satisfied¹. We may therefore assume that $A_0 = 0$. We find that

$$A \times \begin{pmatrix} 0 & z & 0 \\ -z & 0 & x \\ 0 & -x & 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -\operatorname{Tr}(z_1 z) & y_1 x & z x_1 + z_1 x \\ x y_1 & 0 & z y_1 \\ x z_1 + x_1 z & y_1 z & -\operatorname{Tr}(x x_1) \end{pmatrix}.$$

Note that the factor of $1/2$ can be safely ignored since it can be absorbed into x_1, y_1, z_1 .

This imposes several conditions:

- (A) $\operatorname{Tr}(z_1 z) = \operatorname{Tr}(x x_1) = 0$
- (B) $x y_1, y_1 z \in \mathbb{O}^0$
- (C) $x z_1 + x_1 z \in \mathbb{O}^0$

Notice that for $w = a_1 s_1 + a_2 s_2 + a_3 s_3 + b_1 t_1 + b_2 t_2 + b_3 t_3 + c(s_4 - t_4) \in \mathbb{O}^0$, we have

$$wz = wt_2 = a_2 s_4 + b_1 s_3 - b_3 s_1 - ct_2,$$

$$xw = s_1 w = -a_2 t_3 + a_3 t_2 + b_1 s_4 - cs_1.$$

Combining this calculation with condition (A) shows that $a_2^z = b_1^x = 0$. With (B) it implies that $a_2^y = b_1^y = 0$, and with condition (C) we get that $b_1^z = -a_2^x$. Putting this all together, we have

$$\begin{aligned}x' &= -y_1 z = b_3^y s_1 + c^y t_2, \\y' &= x z_1 + x_1 z = (a_3^z - c^x) t_2 - (c^z + b_3^y) s_1, \\z' &= -x y_1 = -a_3^y t_2 + c^y s_1.\end{aligned}$$

This proves the Lemma. □

Proposition 5.2 implies that $C_c^\infty(\Omega_2)$ is a submodule of $C_c^\infty(\Omega_0)$ and $C_c^\infty(\Omega_1)$ is a quotient. Let S_1 and S_2 be the stabilizers of $((s_1, 0, 0), (0, 0, 0))$ and $((s_1, t_2, 0), (0, 0, 0))$ in $G_2 \times \operatorname{GSp}_6$, respectively. Let P_m and Q_m , $m = 1, 2$, be the maximal parabolic subgroups in G_2 and GSp_6 , respectively, as introduced in 3.1.1 and 4.2.5. In particular, these parabolic groups come with maps $P_m \times Q_m \rightarrow \operatorname{GL}_m \times \operatorname{GL}_m$. Then S_m is the inverse image of $\Delta(\operatorname{GL}(m))$, the diagonally embedded GL_m into $\operatorname{GL}_m \times \operatorname{GL}_m$. Now one can easily deduce that $\mathbf{V}_{\bar{U}}$, as a representation of $G_2 \times \operatorname{GSp}_6$, has the following three sub quotients, here induction is not normalized.

- (1) $C_c^\infty(\Omega_2) \cong \operatorname{Ind}_{P_2 \times Q_2}^{G_2 \times \operatorname{GSp}_6}(C_c^\infty(\operatorname{GL}_2)) \otimes |i|^5$.
- (2) $C_c^\infty(\Omega_1) \cong \operatorname{Ind}_{P_1 \times Q_1}^{G_2 \times \operatorname{GSp}_6}(C_c^\infty(\operatorname{GL}_1)) \otimes |i|^5$.

¹This is the action of Sp_6

$$(3) \mathbf{V}_{\bar{N}} \cong \mathbf{V}(M) \otimes |i|^3 \oplus 1 \otimes |i|^5.$$

Proposition 4.4 is simply a normalized version of this result.

REFERENCES

- [1] Armand Borel. Admissible representations of a semi-simple group over a local field with vectors fixed under an Iwahori subgroup. *Invent. Math.*, 35:233–259, 1976.
- [2] Benedict H. Gross. On the Satake isomorphism. In *Galois representations in arithmetic algebraic geometry (Durham, 1996)*, volume 254 of *London Math. Soc. Lecture Note Ser.*, pages 223–237. Cambridge Univ. Press, Cambridge, 1998.
- [3] Jing-Song Huang, Kay Magaard, and Gordan Savin. Unipotent representations of G_2 arising from the minimal representation of D_4^E . *J. Reine Angew. Math.*, 500:65–81, 1998.
- [4] K. Magaard and G. Savin. Exceptional Θ -correspondences. I. *Compositio Math.*, 107(1):89–123, 1997.
- [5] Colette Mœglin, Marie-France Vignéras, and Jean-Loup Waldspurger. *Correspondances de Howe sur un corps p -adique*, volume 1291 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.
- [6] Stephen Rallis and David Soudry. Cubic correspondences arising from G_2 . *Amer. J. Math.*, 119(2):251–335, 1997.
- [7] Gordan Savin and Michael Woodbury. Structure of internal modules and a formula for the spherical vector of minimal representations. *J. Algebra*, 312(2):755–772, 2007.

G. S.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT
E-mail address: savin@math.utah.edu

M. W.: DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY
E-mail address: woodbury@math.columbia.edu