

Mathematics G4402. Modern Geometry
Assignment 5

Fall 2011

Due on Monday, October 17, 2011

[dC]= do Carmo, *Riemannian Geometry*.

- (1) Define a smooth function Q on \mathbb{R}^{n+1} by

$$Q(x_0, x_1, \dots, x_n) = -x_0^2 + x_1^2 + \dots + x_n^2.$$

Define a smooth $(0, 2)$ symmetric tensor q on \mathbb{R}^{n+1} by

$$q = -dx_0^2 + dx_1^2 + \dots + dx_n^2.$$

- (a) (Hyperbolic space) Note that -1 is a regular value of the smooth function Q , so

$$H^n = \{(x_0, x_1, \dots, x_n) \in \mathbb{R}^{n+1} \mid Q(x_0, \dots, x_n) = -1, x_0 > 0\}$$

is an n -dimensional submanifold of \mathbb{R}^{n+1} . Let $i : H^n \hookrightarrow \mathbb{R}^{n+1}$ be the inclusion map, and define $g = i^*q \in C^\infty(H^n, S^2T^*H^n)$. Show that g is positive definite, so it is a Riemannian metric on H^n .

- (b) (Poincaré disk) Show that

$$(x_0, \dots, x_n) \mapsto \frac{1}{x_0 + 1}(x_1, \dots, x_n).$$

defines a diffeomorphism f from H^n onto the unit disk

$$D^n = \{(y_1, \dots, y_n) \in \mathbb{R}^n \mid y_1^2 + \dots + y_n^2 < 1\}.$$

Show that $(f^{-1})^*g = \rho \sum_{i=1}^n dy_i^2$ for some smooth, positive function ρ on D^n , and find $\rho(y_1, \dots, y_n)$.

- (c) (Poincaré upper half space) Let $\mathcal{H}^n = \{(y_1, \dots, y_n) \in \mathbb{R}^n \mid y_n > 0\}$ be the n -dimensional upper half space. Define a Riemannian metric h on \mathcal{H}^n by

$$h = \frac{dy_1^2 + \dots + dy_n^2}{y_n^2}.$$

Prove that (\mathcal{H}^n, h) is isometric to $(D^n, (f^{-1})^*g)$. (Hint: see [GHL] page 57.)

- (2) Let T^2 be embedded in \mathbb{R}^3 as image of \mathbb{R}^2 by the map Φ defined by

$$\Phi(\theta, \phi) = ((a + b \cos \theta) \cos \phi, (a + b \cos \theta) \sin \phi, b \sin \theta)$$

where $a > b > 0$. Let g the Riemannian metric induced on T^2 by the Euclidean metric of \mathbb{R}^3 .

(a) Write g in the form $g = E d\theta^2 + F(d\theta d\phi + d\phi d\theta) + G d\phi^2$.

(b) Compute the volume of (T^2, g) .

- (3) Show that any isometry of the Euclidean space \mathbb{R}^n must take straight lines to straight lines. Show that the only isometries of \mathbb{R}^n are those of the form $\mathbf{x} \mapsto A\mathbf{x} + \mathbf{b}$ for constant $A \in O(n)$, $\mathbf{b} \in \mathbb{R}^n$.