

Mathematics G4402. Modern Geometry
Assignment 8

Fall 2015

Due on Thursday, November 12, 2015

- (1) Let H be a closed Lie subgroup of a Lie group G , and let $G/H = \{aH \mid a \in G\}$ be the set of *left* cosets of H in G . (In other words, let H act on G by *right* multiplication and let G/H be the quotient.) G acts on G/H on the *left* by $G \times G/H \rightarrow G/H$, $(a, bH) \mapsto abH$. Let g be a *right* invariant Riemannian metric on G . By the theorems stated in class, there is a unique Riemannian metric \hat{g} on G/H such that $\pi : (G, g) \rightarrow (G/H, \hat{g})$ is a Riemannian submersion. Prove that if g is left invariant then G acts isometrically on $(G/H, \hat{g})$.
- (2) Let g_n be the bi-invariant metric on $SO(n)$ defined in Problem 3 of Assignment 6. We have seen in class that there is a diffeomorphism $f : S^n \rightarrow SO(n+1)/SO(n)$.

Let \hat{g} be the unique Riemannian metric on $SO(n+1)/SO(n)$ such that

$$\pi : (SO(n+1), g_{n+1}) \rightarrow (SO(n+1)/SO(n), \hat{g})$$

is a Riemannian submersion. Prove that $f^*\hat{g} = \lambda g_{\text{can}}$ for some $\lambda > 0$, and find λ . (Hint: What is the horizontal space $H_{I_{n+1}} \subset T_{I_{n+1}}SO(n+1)$?)

- (3) Let $\mathcal{H} = \{(y_1, y_2) \in \mathbb{R}^2 \mid y_2 > 0\}$ be the upper half plane, and define a Riemannian metric on \mathcal{H} by

$$g = \frac{dy_1^2 + dy_2^2}{y_2^2}.$$

- (a) Compute the Christoffel symbols Γ_{ij}^k , $i, j, k \in \{1, 2\}$, for the Levi-Civita connection ∇ on (\mathcal{H}, g) .
- (b) Define $\gamma : \mathbb{R} \rightarrow \mathcal{H}$ by $\gamma(t) = (t, 1)$. Then γ is a smooth curve in \mathcal{H} . Let

$$V(t) = a(t) \frac{\partial}{\partial y_1} + b(t) \frac{\partial}{\partial y_2}$$

be the unique parallel (w.r.t. ∇) vector field along γ such that $V(0) = \frac{\partial}{\partial y_2}$. Find $a(t), b(t)$ for $t \in \mathbb{R}$.

- (4) Let $F : (M, g) \rightarrow (N, h)$ be an isometric immersion. For any $p \in M$, let π_p be the orthogonal projection from $T_{F(p)}N$ to the image of $dF_p : T_pM \rightarrow T_{F(p)}N$. Let X, Y be C^∞ vector fields on M which are F -related to C^∞ vector fields \tilde{X}, \tilde{Y} on N , respectively. Let ∇ and $\tilde{\nabla}$ be the Levi-Civita connections on (M, g) and on (N, h) , respectively. Prove that for any $p \in M$,

$$dF_p((\nabla_X Y)(p)) = \pi_p((\tilde{\nabla}_{\tilde{X}} \tilde{Y})(F(p)))$$