

Q: When do we have $[M(\lambda):L(\mu)] \neq 0$?

Last time (Verma): It suffices to put $\mu = s_{\alpha} \cdot \lambda$ s.t. $\lambda \geq \mu$ because then $M(\mu) \hookrightarrow M(\lambda)$.

Cor: if we have $\mu = s_{\alpha_1} \dots s_{\alpha_r} \cdot \lambda \leq s_{\alpha_2} \dots s_{\alpha_r} \cdot \lambda \leq \dots \leq s_{\alpha_r} \cdot \lambda \leq \lambda$ (*)
we get $M(\mu) \hookrightarrow M(\lambda)$ also.

When (*) is satisfied we say μ is strongly linked to λ and we write $\mu \uparrow \lambda$

BGG Theorem: If $[M(\lambda):L(\mu)] \neq 0$, then $\mu \uparrow \lambda$.
We will prove this shortly.

Recall that \mathcal{O} is split into categories \mathcal{O}_{λ} , indexed by \mathfrak{g} -antidominant weights λ . Assume λ is integral and dot-regular, so in particular λ is the unique \mathfrak{g} -antidominant weight in $W \cdot \lambda$. Then we can rephrase the condition $w' \cdot \lambda \uparrow w \cdot \lambda$ exclusively in terms of the Weyl group.

The Bruhat order: the length function $l: W \rightarrow \mathbb{Z}_{\geq 0}$ is defined equivalently as

$$l(w) = \min \{ n \mid w = s_{\alpha_1} \dots s_{\alpha_n}, \alpha_i \in \Delta^+ \}$$

$$= \# \{ \alpha > 0 \mid w\alpha < 0 \} \quad \text{"reduced expression"}$$

Whenever we have $w' = s_{\alpha} w$ ($\alpha \in \Delta^+$), we write $w' \xrightarrow{\alpha} w$.

The Bruhat order is then defined as $w' \leq w$ iff $\exists \alpha_1, \dots, \alpha_n \in \Delta^+$ s.t.
 $w' \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} w$

Equivalently: Some (every) reduced expression for w has a subexpression which is a reduced expression for w' .

Prop: If λ is integral, dot-regular and \mathfrak{g} -antidominant, then
 $w' \cdot \lambda \uparrow w \cdot \lambda \iff w' \leq w$




Fig: The highlighted dots are the elements of $W \cdot \lambda$. Here λ is the lowest weight in $W \cdot \lambda$ and is therefore \mathfrak{g} -antidominant.

Proof: It suffices to prove $s_{\alpha} w \uparrow w \cdot \lambda \iff s_{\alpha} w \xrightarrow{\alpha} w$
These are equivalent since

$$s_{\alpha} (w \cdot \lambda + \rho) < w \cdot \lambda + \rho$$

$$\iff s_{\alpha} w (\lambda + \rho) < w (\lambda + \rho)$$

$$\iff \langle w (\lambda + \rho), \alpha^\vee \rangle > 0$$

$$\iff \langle \lambda + \rho, w^{-1} \alpha^\vee \rangle > 0$$

$$\stackrel{\lambda \text{ antidom}}{\iff} w^{-1} \alpha < 0$$

$$\iff w^{-1} s_{\alpha} \alpha > 0$$

$$\iff l(w^{-1} s_{\alpha}) = l(w^{-1}) - 1$$

$$\iff l(s_{\alpha} w) = l(w) - 1$$

$$\iff s_{\alpha} w \xrightarrow{\alpha} w \quad \square$$

Remark: The condition $w' \leq w$ is independent of λ ! (So we may only look at \mathcal{O}_0)

Summary: $[M(w \cdot \lambda):L(w' \cdot \lambda)] \neq 0 \iff w' \cdot \lambda \uparrow w \cdot \lambda \iff w' \leq w$.
(if λ dot-reg, \mathfrak{g} -antidom, integral)

Reminder on the Bruhat order: $w' \xrightarrow{\alpha} w$ means $w' = s_{\alpha} w$ and $l(w') = l(w) - 1$
 $w' \leq w$ means $w' \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} w$

Also $l(w) = \min \{ k \mid s_{\alpha_1} \dots s_{\alpha_k} = w \} = \# \{ \alpha > 0 : w\alpha < 0 \}$
(simple)
 $l(w^{-1}) = l(w)$

Rmk: it is possible to have $w \cdot \lambda < \lambda$ yet $w \cdot \lambda \not\leq \lambda$ (only in rank ≥ 3)

Example: $\mathfrak{g} = \mathfrak{sl}_4 \mathbb{C}$, $\lambda = s_{\alpha_2} s_{\alpha_3} \cdot (-\lambda_2)$ Computation shows $\lambda - \mu = \alpha_1 + \alpha_3$. However,
 $\mu = s_{\alpha_3} s_{\alpha_2} s_{\alpha_1} \cdot (-\lambda_2)$




Fig 2: $s_{\alpha_2}s_{\alpha_3}$ and $s_{\alpha_3}s_{\alpha_2}s_{\alpha_1}$ are unrelated. (Bruhat order)

The BGG Theorem is a corollary of:

Theorem (Jantzen Filtration): let $\lambda \in \mathfrak{h}^*$ arbitrary. Then $M(\lambda)$ has a filtration
 $M(\lambda) = M(\lambda)^0 \supset M(\lambda)^1 \supset \dots \quad (M(\lambda)^{\infty} = 0)$

such that

(a) Each nonzero quotient $M(\lambda)^i / M(\lambda)^{i+1}$ has a nondegenerate contravariant form

(b) $M(\lambda)^i / M(\lambda)^{i+1} = L(\lambda)$ (i.e. $M(\lambda)^1 = N(\lambda)$)

(c) $\sum_{i \geq 0} \text{ch } M(\lambda)^i = \sum_{\alpha > 0, s_{\alpha} \cdot \lambda < \lambda} \text{ch } M(s_{\alpha} \cdot \lambda)$ ("Jantzen sum formula")

Cor: (BGG Theorem): By induction on $k = \#\{\mu \in W \cdot \lambda \mid \mu \leq \lambda\}$. If $k=1$ then λ is minimal so $M(\lambda)$ is simple. Otherwise assume $[M(\lambda) : L(\mu)] > 0$, $\mu < \lambda$. Then $[M(\lambda)^1 : L(\mu)] > 0$ so $[M(s_{\alpha} \cdot \lambda) : L(\mu)] > 0$ for some $\alpha > 0$ s.t. $s_{\alpha} \cdot \lambda < \lambda$, so $s_{\alpha} \cdot \lambda \uparrow \lambda$. But by induction hypothesis $\mu \uparrow s_{\alpha} \cdot \lambda$, so $\mu \uparrow \lambda$. \bullet

Application of Jantzen's Filtration

For the next application of the theorem, we need a result that was mentioned last time:

(4.10)

Prop: let λ be integral, dot-regular, ρ -antidominant. Then for any $w \cdot \lambda \in W \cdot \lambda$, $M(\lambda) = L(\lambda)$ is the unique simple submodule of $M(w \cdot \lambda)$. Furthermore, $[M(w \cdot \lambda) : L(\lambda)] = 1$.

Proof: We saw the first part last time.

For the second part, it suffices to prove, by BGG Reciprocity, that $(P(\lambda) : M(w \cdot \lambda)) \leq 1$.

Now, recall that $-\rho$ is ρ -dominant $\Rightarrow M(-\rho)$ is projective. Also $w_0 \cdot \lambda$ is ρ -dominant so $w_0 \cdot \lambda + \rho \in \Lambda^+$.

Thus $L(w_0 \cdot \lambda + \rho)$ is f.dimp hence $M = M(-\rho) \otimes L(w_0 \cdot \lambda + \rho)$ is projective.

Fact: M has a standard filtration with $M(\omega \cdot \lambda)$ appearing at most once and $M(\lambda)$ appearing only at the top

Using the fact: since $M \rightarrow M(\lambda)$ we have $P(\lambda) \subseteq T$ and so $P(\lambda)$ has a std filtration with $M(\omega \cdot \lambda)$ appearing at most once, as desired.
(since T is projective)

Skip:

Proof of the fact: let v_1, \dots, v_n be a basis of weight vectors for $L(\omega_0 \cdot \lambda + \rho)$ with weights μ_1, \dots, μ_n . Reorder the basis so that $\mu_i \leq \mu_j \Rightarrow i \leq j$ and so that v_1 has weight λ (which is minimal since $\omega_0(\omega_0 \cdot \lambda + \rho) = \lambda + \rho$)

Now tensor the filtration of $U(\mathfrak{b})$ -modules

$$\mathbb{C}_{-j} \otimes_{U(\mathfrak{b})} \text{Span}_{U(\mathfrak{b})}(v_n) \subsetneq \mathbb{C}_{-j} \otimes_{U(\mathfrak{b})} \text{Span}_{U(\mathfrak{b})}(v_n, v_{n-1}) \subsetneq \dots \subsetneq \mathbb{C}_{-j} \otimes L(\omega_0 \cdot \lambda + \rho) \quad \left(\begin{array}{l} \text{with quotients } \mathbb{C}_{-j} \otimes \mathbb{C}_{\mu_i} \\ \text{Note: } \omega(\lambda + \rho) \text{ appears only once} \end{array} \right)$$

with $U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} -$ (which is exact of f.d. $U(\mathfrak{b})$ -modules). We get

$$U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} (\mathbb{C}_{-j} \otimes_{U(\mathfrak{b})} \text{Span}_{U(\mathfrak{b})}(v_n)) \subsetneq \dots \subsetneq U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} (\mathbb{C}_{-j} \otimes L(\omega_0 \cdot \lambda + \rho))$$

$$\left(U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} \mathbb{C}_{-j} \right) \otimes L(\omega_0 \cdot \lambda + \rho)$$

$$\parallel \\ M(-j) \otimes L(\omega_0 \cdot \lambda + \rho)$$

and each quotient is isomorphic to $U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} (\mathbb{C}_{-j} \otimes \mathbb{C}_{\mu_i}) = M(\mu_i - j)$, in particular $M(\mu_1 - j) = M(\lambda)$.

Remark: Result $\Leftrightarrow P_{w, \omega_0} = 1 \forall w$. Also, KL (74): $X(\gamma)$ not smooth iff $P_{w, \gamma} = 1 \forall w \leq \gamma$, so $P_{w, \omega_0} = 1 \forall w \leq \omega_0$ because

Break?

$$\forall x \quad H^i(M, M(-i, \gamma); \mathbb{Q}) = \begin{cases} \mathbb{Q} & i = 2 \dim M \\ 0 & \text{o/w} \end{cases}$$

$$X(\omega_0) = \overline{B \omega_0 B} / B = G/B \text{ smooth} \\ \text{(here not smooth)}$$

Finding the composition factors of $M(\lambda)$ (λ integral, dot-reg) for $sl_3 \mathbb{C}$

Recall that knowing $\{ \text{ch } M(w \cdot \lambda) \}_{w \in W}$ in terms of $\{ \text{ch } L(w \cdot \lambda) \}_{w \in W}$ (or vice-versa)
 \iff knowing $[M(w \cdot \lambda) : L(w \cdot \lambda)] \quad \forall w, w' \in W$.

$\mathfrak{g} = sl_3 \mathbb{C}$, λ integral, dot-regular, ρ -antidominant (e.g. $\lambda = -2\rho$)

4

So far we have $\text{ch } M(\lambda) = \text{ch } L(\lambda)$ (since λ ρ -antidom, $M(\lambda)$ simple)

Next, $[M(s_{\alpha_1} \cdot \lambda) : L(s_{\alpha_1} \cdot \lambda)] = 1$

$[M(s_{\alpha_1} \cdot \lambda) : L(\lambda)] = 1$ (by the Proposition)

and these are all the candidates, so

$$\text{ch } M(s_{\alpha_1} \cdot \lambda) = \text{ch } L(\lambda) + \text{ch } L(s_{\alpha_1} \cdot \lambda)$$

$$\text{sim: } \text{ch } M(s_{\alpha_2} \cdot \lambda) = \text{ch } L(\lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda)$$

Next, for $s_{\alpha_1} s_{\alpha_2} \cdot \lambda$ we know

$$[M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(s_{\alpha_1} s_{\alpha_2} \cdot \lambda)] = 1 = [M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(\lambda)]$$

but we need

$$[M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(s_{\alpha_1} \cdot \lambda)], [M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(s_{\alpha_2} \cdot \lambda)]$$




Fig 2: The orbit $W \cdot \lambda$. Here $\mu_1 \leftarrow \mu_2$ means $\mu_1 \uparrow \mu_2$

Jantzen's sum formula

$$\Rightarrow \sum_{i > 0} \text{ch } M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda)^i = \text{ch } M(s_{\alpha_1} \cdot \lambda) + \text{ch } M(s_{\alpha_2} \cdot \lambda)$$

$$= \text{ch } L(s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda) + 2 L(\lambda)$$

Since $[M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(\lambda)] = 1$, we deduce $M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda)^2 = L(\lambda)$ and

$$\text{ch } M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda)^2 = \text{ch } L(s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda) + \text{ch } L(\lambda)$$

$$\text{Therefore, } [M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(s_{\alpha_1} \cdot \lambda)] = 1 = [M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) : L(s_{\alpha_2} \cdot \lambda)]$$

$$\text{sim: } [M(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) : L(s_{\alpha_1} \cdot \lambda)] = 1 = [M(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) : L(s_{\alpha_2} \cdot \lambda)]$$

Finally, for $s_{\alpha_1} s_{\alpha_2} s_{\alpha_1} \cdot \lambda = w_0 \cdot \lambda$, we have

$$\sum_{i > 0} \text{ch } M(w_0 \cdot \lambda)^i = \text{ch } M(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) + \text{ch } M(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) + \text{ch } L(\lambda)$$

$$= \text{ch } L(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) + \text{ch } L(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) + 2(\text{ch } L(s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda)) + 3 \text{ch } L(\lambda)$$

Again, since $[M(w_0 \cdot \lambda) : L(\lambda)] = 1$, we deduce $M(w_0 \cdot \lambda)^3 = L(\lambda)$

Two possibilities remain:

$$\text{ch } M(w_0 \cdot \lambda)^2 = \text{ch } L(s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda) + \text{ch } L(\lambda)$$

$$\text{ch } M(w_0 \cdot \lambda)^1 = \text{ch } L(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) + \text{ch } L(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_1} \cdot \lambda) + \text{ch } L(s_{\alpha_2} \cdot \lambda) + \text{ch } L(\lambda)$$

or

$$\begin{cases} \text{ch } M(\omega_0 \cdot \lambda)^2 = \text{ch } L(\lambda) \\ \text{ch } M(\omega_0 \cdot \lambda)^1 = \text{ch } L(s_{\alpha_1} s_{\alpha_2} \cdot \lambda) + \text{ch } L(s_{\alpha_2} s_{\alpha_1} \cdot \lambda) + 2 \text{ch } L(s_{\alpha_1} \cdot \lambda) + 2 \text{ch } L(s_{\alpha_2} \cdot \lambda) + \text{ch } L(\lambda) \end{cases} \quad (*)$$

Humphreys claims this case can be dealt with using the same tools as in the previous cases, but I don't see how. Instead, here is an ad hoc argument: the weight space $M(\omega_0 \cdot \lambda)_{s_{\alpha_1} \cdot \lambda}$ has dimension 2 and

$$L(s_{\alpha_1} \cdot \lambda)_{s_{\alpha_1} \cdot \lambda} = 1, \quad L(s_{\alpha_2} s_{\alpha_1} \cdot \lambda)_{s_{\alpha_1} \cdot \lambda} = 1 \quad (\text{since } M(s_{\alpha_2} s_{\alpha_1} \cdot \lambda)_{s_{\alpha_1} \cdot \lambda} = 2 \text{ and we know } \text{ch } M(s_{\alpha_2} s_{\alpha_1} \cdot \lambda))$$

$$\text{so } [M(\omega_0 \cdot \lambda) : L(s_{\alpha_1} \cdot \lambda)] \leq 1$$

$$\text{We thus discard } (*) \text{ and conclude } [M(\omega_0 \cdot \lambda) : L(\omega_0 \cdot \lambda)] = 1 \quad \forall w \in W$$

Incidentally, this provides an example of the question we discussed last week:

$$\text{soc}(M(\mathfrak{o}) / L(-2\rho)) \supset M(\mathfrak{o})^2 / L(-2\rho) = L(s_{\alpha_1} \cdot (-2\rho)) \oplus L(s_{\alpha_2} \cdot (-2\rho)) = L(-\rho - \alpha_1) \oplus L(-\rho - \alpha_2)$$

Skip: (Proof of \oplus : let $\psi \in \text{Out}(\mathfrak{sl}_2 \mathbb{C})$ corresponding to $\begin{matrix} \xrightarrow{\alpha_1} \\ \xrightarrow{\alpha_2} \end{matrix}$. Then if $M = M(\mathfrak{o}) / L(-2\rho)$, set $M^\psi = M$, action twisted by ψ . This is isomorphic to M since $\psi(\mathfrak{o}) = \mathfrak{o}$,

However $L(-\rho - \alpha_1)^\psi = L(-\rho - \alpha_2)$ so one can't be on top of the other.)

Proof of Jantzen's Filtration Theorem

"Key Lemma"

Let A be a PID and $M = A^r$ with a nondegenerate bilinear form $(,)$ with determinant $D \neq 0$.
 Let $p \in A$ be prime. Define a filtration $M = M(0) \supset M(1) \supset \dots$ by $M(n) = \{m \in M : (m, M) \subset p^n A\}$.
 Write also $\bar{A} = A/pA$, $\bar{M} = M/pM$ etc.

Then

(a) $\nu_p(D) = \sum_{n \geq 0} \dim_{\bar{A}} \bar{M}(n)$ ($\bar{M}(n) = 0$ for n large)

(b) $(,)_n = p^{-n}(,)$ induces a nondeg form on $\bar{M}(n) / \bar{M}(n+1)$

Skip?: Proof omitted. Example instead: $A = \mathbb{C}[T]$, $M = AX \oplus AY$, bilinear form $\begin{pmatrix} 2T & 0 \\ 0 & T^3 \end{pmatrix}$

Then

$M(0) = AX \oplus AY$

$\bar{M}(0) = \mathbb{C}X \oplus \mathbb{C}Y$

$M(1) = pX \oplus AY$

$\bar{M}(1) = \mathbb{C}X \oplus \mathbb{C}Y$

$M(2) = (T)X \oplus AY$

$\bar{M}(2) = \mathbb{C}Y$

$M(3) = (T^2)X \oplus AY$

$\bar{M}(3) = \mathbb{C}Y$

$M(4) = (T^3)X \oplus (T)Y$

$\bar{M}(4) = 0$

$4 = \nu_T(T^4)$ and eg $\bar{M}(1) / \bar{M}(2)$ has form (2)

Theorem (Jantzen Filtration): let $\lambda \in \mathfrak{h}^*$ arbitrary. Then $M(\lambda)$ has a filtration $M(\lambda) = M(\lambda)^0 \supset M(\lambda)^1 \supset \dots \supset M(\lambda)^N = 0$

such that

(a) Each nonzero quotient $M(\lambda)^i / M(\lambda)^{i+1}$ has a nondegenerate contravariant form

(b) $M(\lambda)^i / M(\lambda)^{i+1} = L(\lambda)$ (i.e. $M(\lambda)^i = N(\lambda)$)

(c) $\sum_{i \geq 0} \text{ch } M(\lambda)^i = \sum_{\alpha > 0, \alpha \cdot \lambda < \lambda} \text{ch } M(\alpha - \lambda)$ ("Jantzen sum formula")

We need to define $M(\lambda)^i$. The idea is to introduce a free variable T by setting $A = \mathbb{C}[T]$, $K = \mathbb{C}(T)$ and $\mathfrak{g}_K = K \otimes_{\mathbb{C}} \mathfrak{g}$, $\mathfrak{g}_A = A \otimes_{\mathbb{C}} \mathfrak{g}$. Let $\lambda_T := \lambda + T\beta \in \mathfrak{h}_K^*$.

The theory over K is the same as over \mathbb{C} , and since $\langle \lambda_T + \beta, \alpha^\vee \rangle \notin \mathbb{Z} \forall \alpha \in \Phi^+$, $\lambda_T + \beta$ is antidominant and therefore $M(\lambda_T)$ is simple, so the contravariant form on it is nondegenerate. Also, the weight spaces are $M(\lambda_T)_{\lambda_T - \nu}$ for $\nu \in R^+$

We also have an A -form $M(\lambda_T)_A \subset M(\lambda_T)$. Write $M_{\lambda_T - \nu} = M(\lambda_T)_{\lambda_T - \nu} \cap M(\lambda_T)_A$

Finally, set $M(\lambda_T)_A^i := \sum_{\nu \in R^+} M_{\lambda_T - \nu}^i$

To get the filtration for $M(\lambda)$, set $T=0$ i.e. $M(\lambda)^i = M(\lambda_T)_A^i / T M(\lambda_T)_A^i$

The Key Lemma tells us $T^{-i}(,)$ induces a nondeg contravariant form on $M(\lambda)^i / M(\lambda)^{i+1}$. Since $M(\lambda) / M(\lambda)^1$ is also a h.w. module, it must be simple. This proves (a) and (b).

Example of this construction: $\mathfrak{sl}_3 \mathbb{C}$, $\lambda = -\lambda_1$. Then $\lambda_T = -\lambda_1 + T\beta$.

We look at a single weight space $M_{\lambda_T - \alpha_1 - \alpha_2}$. Now $M_{\lambda_T - \alpha_1 - \alpha_2}$ has two basis vectors: $a = f_{\alpha_1} f_{\alpha_2} v^+$ and $b = f_{\alpha_1 + \alpha_2} v^+$.

The contravariant form wrt this basis is $\begin{pmatrix} T^2 & -T \\ -T & 2T-1 \end{pmatrix}$

For instance, $(f_{\alpha_1} f_{\alpha_2} v^+, f_{\alpha_1} f_{\alpha_2} v^+) = (v^+, e_{\alpha_2} e_{\alpha_1} f_{\alpha_1} f_{\alpha_2} v^+)$.

$$\begin{aligned} e_{\alpha_2} e_{\alpha_1} f_{\alpha_1} f_{\alpha_2} v^+ &= e_{\alpha_2} f_{\alpha_1} e_{\alpha_1} f_{\alpha_2} v^+ + e_{\alpha_2} h_{\alpha_1} f_{\alpha_2} v^+ = e_{\alpha_2} f_{\alpha_2} h_{\alpha_1} v^+ + e_{\alpha_2} f_{\alpha_2} v^+ = (1+h_{\alpha_1}) h_{\alpha_2} v^+ \\ &= (1+h_{\alpha_1}) \underbrace{(-\lambda_1 + T\beta)}_T (\alpha_2) v^+ = T \underbrace{(1 + (-\lambda_1 + T\beta)(\alpha_1))}_{1-1+T} v^+ = T^2 v^+ \end{aligned}$$

Therefore $M_{\lambda_T - \alpha_1 - \alpha_2}(i) = \begin{cases} Aa \otimes Ab & i=0 \\ Aa \otimes (T)b & i=1 \\ Aa \otimes (T^2)b & i=2 \\ (T)a \otimes (T^2)b & i=3 \end{cases}, M(\lambda)_{\lambda - \alpha_1 - \alpha_2}^i = \begin{cases} Ca \otimes Cb & i=0 \\ Ca & i=1 \\ Ca & i=2 \\ 0 & i \geq 3 \end{cases}$

In order to verify (c) $\sum_{i \geq 0} \text{ch } M(\lambda)^i = \sum_{\alpha > 0, \delta \alpha \leq \lambda} \text{ch } M(\delta \alpha - \lambda)$, we need to compute $\sum_{i \geq 0} \dim M(\lambda)_{\lambda - \nu}^i$.

The key lemma tells us that this is $\nu_T(D_\nu(\lambda_T))$ (Here $D_\nu(\lambda_T)$ is the determinant of the form on $M_{\lambda_T - \nu} = M(\lambda_T)_\lambda \cap M(\lambda_T)_{\lambda_T - \nu}$)

Claim: (Shapovalov) $D_\nu(\lambda_T) = \prod_{\alpha > 0} \prod_{r \geq 0} (\langle \lambda_T + r\beta, \alpha^\vee \rangle - r)^{P(\nu - r\alpha)}$ (Here $P(\delta) = \#$ ways to write δ as a sum of positive roots)

Using the claim, note $\langle \lambda_T + r\beta, \alpha^\vee \rangle - r = \langle \lambda + r\beta, \alpha^\vee \rangle - r + T$ is a multiple of T iff $\langle \lambda + r\beta, \alpha^\vee \rangle = r$ iff $\delta \alpha \cdot \lambda < \lambda$. So $\nu_T(D_\nu(\lambda_T)) = P(\nu - \langle \lambda + r\beta, \alpha^\vee \rangle \alpha)$ and therefore

$$\begin{aligned} \sum_{i \geq 0} \text{ch } M(\lambda)^i &= \sum_{\nu \in \mathbb{R}^+} \sum_{\substack{\alpha > 0 \\ \delta \alpha \cdot \lambda < \lambda}} P(\nu - \langle \lambda + r\beta, \alpha^\vee \rangle \alpha) e^{\lambda - \nu} \\ &= \sum_{\substack{\alpha > 0 \\ \delta \alpha \cdot \lambda < \lambda}} \sum_{\nu \in \mathbb{R}^+} P(\nu') e^{\lambda - \langle \lambda + r\beta, \alpha^\vee \rangle \alpha - \nu'} \\ &\quad \downarrow \\ &\quad P(\nu - \langle \lambda + r\beta, \alpha^\vee \rangle \alpha) \neq 0 \\ &\quad \text{so } \nu' = \nu - \langle \lambda + r\beta, \alpha^\vee \rangle \alpha \in \mathbb{R}^+ \end{aligned}$$

On the other hand, $\text{ch } M(\delta \alpha - \lambda) = \sum_{\nu' \in \mathbb{R}^+} P(\nu') e^{\delta \alpha \cdot \lambda - \nu'}$, and (c) follows.

Skip? Shapovalov's determinantal formula

In order to get the bilinear form on $M_{\lambda-r}$ above, we computed the contravariant form by simplifying the PBW monomials as follows

$$(f_i v^+, f_j v^+) = (v^+, \tau(f_i) f_j v^+) = (v^+, t_{ij} v^+), \text{ where } t_{ij} \in U(\mathfrak{h}). \text{ Then } t_{ij} v^+ = \lambda_{\tau}(t_{ij}) v^+$$

The determinant of the form is then $\det(\lambda_{\tau}(t_{ij}))$, but we can also get this by

$$\det(t_{ij}) v^+ = \det(\lambda_{\tau}(t_{ij})) v^+$$

Ultimately, knowing $\det(t_{ij})$ will tell us $D_r(\lambda_{\tau})$ for all λ . This is Shapovalov's formula:

$$D_r = \det(t_{ij}) = \prod_{\alpha > 0} \prod_{r \geq 0} (h_{\alpha} + \langle \rho, \alpha^{\vee} \rangle - r)^{P(\nu - r\alpha)}$$

already computed, others are easy

Continuing the previous example, we would have $(t_{ij}) = \begin{pmatrix} h_{\beta}(h_{\alpha}+1) & -h_{\beta} \\ -h_{\beta} & h_{\alpha}+h_{\beta} \end{pmatrix}$,

whose determinant is $h_{\beta} h_{\alpha} (h_{\alpha} + h_{\beta} + 1)$.

Comment on the proof strategy?

Comment on what happens if we replace \mathfrak{g} by something else?

•

