

Defomation Theory Graduate Student Seminar
Fall 2021

Notes by Patrick Lei

Lectures by Various

Disclaimer

These notes were taken during the seminar using the vimtex package of the editor neovim. Any errors are mine and not the speakers'. In addition, my notes are picture-free (but will include commutative diagrams) and are a mix of my mathematical style and that of the lecturers. If you find any errors, please contact me at plei@math.columbia.edu.

Seminar Website: <https://www.math.columbia.edu/~dejong/seminar.html>

Contents

Contents • 2

- 1 Johan (Sep 24): Schlessinger's paper • 4
- 2 Ivan and Cailan (Oct 1): Deformations of schemes • 8
 - 2.1 DEFORMATIONS OF AFFINE SCHEMES • 8
 - 2.2 DEFORMATIONS OF SCHEMES • 11
- 3 Kevin (Oct 08): Deformations of coherent sheaves • 14
 - 3.1 TANGENT-OBSTRUCTION THEORY • 16
- 4 Patrick (Oct 15): Deformations of singularities • 18
 - 4.1 EXPLICIT CRITERIA FOR FLATNESS • 18
 - 4.2 HILBERT SCHEMES OF SMOOTH SURFACES • 19
 - 4.3 AN OBSTRUCTED DEFORMATION • 22
- 5 Avi (Oct 22): Local-global methods • 24
 - 5.1 CURVES WITH ISOLATED SINGULARITIES • 24
 - 5.2 SMOOTH CURVES WITH A FINITE GROUP ACTION • 26
- 6 Caleb and Morena (Oct 29): [Lemma 0E3X](#) and applications to contracting curves • 28
 - 6.1 ONE LEMMA IN THE STACKS PROJECT • 28
 - 6.2 APPLICATION TO MODULI OF CURVES • 30
 - 6.2.1 Stable curves • 31
 - 6.2.2 Contraction done wrong • 32
 - 6.2.3 Contraction done right • 32
- 7 Morena (Nov 05): Contraction morphisms between moduli stacks of curves • 33
 - 7.1 RECAP OF LAST TIME • 33
 - 7.2 CONTRACTION OF RATIONAL TAILS AND BRIDGES • 33
 - 7.3 CONTRACTION OVER ANY BASE SCHEME • 35
- 8 Baiqing (Nov 12): Deformations of group schemes • 37
 - 8.1 DEFORMATIONS OF ABELIAN SCHEMES • 37
 - 8.2 DEFORMATIONS OF SMOOTH AFFINE GROUP SCHEMES • 39

- 9 Haodong (Nov 19): Artin's axioms • 41
 - 9.1 PRESTACKS AND STACKS • 41
 - 9.1.1 Prestacks • 41
 - 9.1.2 Stacks • 43
 - 9.2 ALGEBRAIC STACKS AND ARTIN'S AXIOMS • 44
 - 9.2.1 Algebraic stacks • 44
 - 9.3 ARTIN'S AXIOMS • 44

- 10 Che (Dec 03): Stack of coherent sheaves • 47
 - 10.1 \mathcal{X} IS A STACK • 47
 - 10.2 REPRESENTABILITY OF THE DIAGONAL • 48
 - 10.3 PRESERVATION OF LIMITS • 48
 - 10.4 RIM-SCHLESSINGER • 49
 - 10.5 FINITENESS • 49
 - 10.6 FORMAL OBJECTS ARE EFFECTIVE • 49

- 11 Johan (Dec 10): Openness of versality • 51
 - 11.1 MODULI OF CURVES • 51
 - 11.2 PROPERTIES OF COTANGENT COMPLEXES • 52
 - 11.3 COHERENT SHEAVES • 53
 - 11.4 A TRICK • 53

Johan (Sep 24): Schlessinger's paper

The paper by Schlessinger is titled *Functors of Artin Rings*. Throughout this lecture, k is a field, \mathcal{C} is the category of Artinian local k -algebras A, B, C, \dots with residue field k , and $\widehat{\mathcal{C}}$ is the category of Noetherian complete local k -algebras R, S, \dots with residue field k .

Remark 1.0.1. Every $R \in \widehat{\mathcal{C}}$ is of the form $k[[x_1, \dots, x_n]]/(f_1, \dots, f_m)$ by the Cohen structure theorem. Then $R \in \mathcal{C}$ if and only if (f_1, \dots, f_m) contains $(x_1, \dots, x_n)^N$ for some N .

Remark 1.0.2. In the paper, there is a more general setup, where Λ is a complete local Noetherian ring with residue field k . Then $\mathcal{C}_\Lambda, \widehat{\mathcal{C}}$ are defined analogously, which will allow things like $\Lambda = \mathbb{Z}_p$.

The idea of deformation theory is to look at functors $F: \mathcal{C} \rightarrow \text{Set}$.

Example 1.0.3. Given $R \in \widehat{\mathcal{C}}$, we set $h_R: \mathcal{C} \rightarrow \text{Set}$ sending $A \mapsto \text{Hom}_{\widehat{\mathcal{C}}}(\widehat{R}, A)$. This is not necessarily representable because $R \notin \mathcal{C}$ in general, but it is pro-representable.

Definition 1.0.4. A functor F is *pro-representable* if $F \simeq h_R$ for some $R \in \widehat{\mathcal{C}}$.

Example 1.0.5. Let M be a variety over k and $m \in M(k)$. Then define

$$\text{Def}_{M,m}(A) = \left\{ \text{Spec } A \xrightarrow{m_A} M \mid m_A|_{\text{Spec } k} = m \right\}.$$

It is easy to see that $\text{Def}_{M,m}(A)$ is pro-representable by $\widehat{\mathcal{O}}_{M,m}$.

Observe that $h_R(k) = \{*\}$ is a singleton. Also note that $h_R(A \times_B C) = h_R(A) \times_{h_R(B)} h_R(C)$. Here, $A \times_B C$ is the fiber product of rings and not the tensor product.

Now consider the following conditions on F : let $A \rightarrow B \leftarrow C$ be a diagram in \mathcal{C} and consider the morphism

$$F(A \times_B C) \xrightarrow{(*)} F(A) \times_{F(B)} F(C).$$

- (H₁) The morphism $(*)$ is surjective if $C \rightarrow B$;
- (H₂) The morphism $(*)$ is bijective if $C = k[\varepsilon] \rightarrow k = B$;
- (H₃) $\dim_k(t_F) < \infty$ (later, we will see that we need H₂ for formulate this). Here, t_F is the tangent space to F ;
- (H₄) The morphism $(*)$ is bijective if $C \rightarrow B$.

Example 1.0.6. Fix a group G and a representation $\rho_0: G \rightarrow GL_n(k)$. Now define

$$\text{Def}_{\rho_0}^{\text{naive}}(A) = \{\rho: G \curvearrowright A^{\oplus n} \mid \rho \pmod{\mathfrak{m}_A} \cong \rho_0\} / \cong.$$

Better, we will define

$$\text{Def}_{\rho_0}(A) = \{\rho: G \curvearrowright A^{\oplus n} \mid \rho \pmod{\mathfrak{m}_A} = \rho_0\} / \ker(GL_n(A) \rightarrow GL_n(k)).$$

In general these functors fail (H_4) and $\text{Def}_{\rho_0}^{\text{naive}}$ even fails (H_2) .

Namely, if $H = \mathbb{Z}$ and ρ_0 is the trivial representation, then for $\text{Def}_{\rho_0}^{\text{naive}}$, we are looking at subsets of

$$GL_n(A \times_B C) / \text{conj} \rightarrow GL_n(A) / \text{conj} \times_{GL_n(B) / \text{conj}} GL_n(C) / \text{conj}.$$

This morphism is always surjective, but in general it is not injective.

For example, if $A = k[\varepsilon_1]$, $B = k$, $C = k[\varepsilon_2]$, we can look at elements of the form $1 + \varepsilon_1 T_1 + \varepsilon_2 T_2$ and see that on the left we can only conjugate together, while on the right we can conjugate both T_1, T_2 arbitrarily. Here $A \times_B C = k[\varepsilon_1, \varepsilon_2] = k[x_1, x_2] / (x_1^2, x_1 x_2, x_2^2)$.

Definition 1.0.7. A natural transformation $t: F \rightarrow G$ of functors on \mathcal{C} is *smooth* if for all surjections $B \rightarrow A$ the map $F(B) \rightarrow F(A) \times_{G(A)} G(B)$ is surjective.

Note that this is equivalent to the existence of a lift in the diagram below:

$$\begin{array}{ccc} \text{Spec } A & \longrightarrow & M \\ \downarrow & \nearrow & \downarrow f \\ \text{Spec } B & \longrightarrow & N. \end{array}$$

This definition is motivated by the following example: let $f: M \rightarrow N$ be a morphism of varieties over k . Let $m \in M(k)$, $n = f(m) \in N(k)$. Then the following are equivalent:

1. $\text{Def}_{M,m} \rightarrow \text{Def}_{N,n}$ is smooth.
2. f is smooth at m .

Definition 1.0.8. We say F has a *hull* if and only if $F(k) = \{*\}$ and there exists a smooth $t: h_R \rightarrow F$ for some $R \in \widehat{\mathcal{C}}$ which induces an isomorphism $t_R \cong t_F$.

Now we will say a bit about tangent spaces.

1. When $F(k) = \{*\}$, then $t_F = F(k[\varepsilon])$.
2. If F satisfies (H_2) and $F(k) = \{*\}$, then t_F has a natural k -vector space structure. Here, H_2 gives $F(k[\varepsilon_1, \varepsilon_2]) \rightarrow F(k[\varepsilon]) \times F(k[\varepsilon])$ is a bijection, and then we take $\varepsilon_1 \mapsto \varepsilon, \varepsilon_2 \mapsto \varepsilon$, which defines addition.
3. $t_R = \text{Hom}_k(\mathfrak{m}_R / \mathfrak{m}_R^2, k) = \text{Hom}_{\widehat{\mathcal{C}}}(R, k[\varepsilon]) = h_R(k[\varepsilon]) = t_{(h_R)}$.

Theorem 1.0.9 (Schlessinger). *Assume that $F(k) = \{*\}$. Then the conditions $(H_1), (H_2), (H_3)$ hold for F if and only if F has a hull. In addition, (H_3) and (H_4) hold if and only if F is pro-representable.*

Very rough idea of proof of \Rightarrow for the hull case. Let $n = \dim_k(t_F)$. Then (H_2) and $n < \infty$ imply the following: Let $S = k[[x_1, \dots, x_n]]$ and $\mathfrak{m} = \mathfrak{m}_S = (x_1, \dots, x_n)$. We can find $\xi_1 \in F(S/\mathfrak{m}^2)$ such that

$$t_S = \text{Hom}_{\hat{\mathcal{C}}}(S, k[\varepsilon]) \xrightarrow{\xi_1} t_F$$

is an isomorphism.

Next, we will choose $q \geq 2$ and consider pairs (J, ξ) where $\mathfrak{m}^{q+1} \subset J \subset \mathfrak{m}^2$ and $\xi \in F(S/J)$ such that $\xi \mapsto \xi_1 \in F(S/\mathfrak{m}^2)$. Say that $(J, \xi) \leq (J', \xi')$ if $J \subset J'$ and $\xi \mapsto \xi'$. Choose a minimal pair (J, ξ) for this ordering. We can choose J_q so that $\mathfrak{m}^{q+1} + J_{q+1} = J_q$ and ξ_{q+1} maps to ξ_q for bookkeeping purposes.

Choose $R = \lim S/J_q$, which is a quotient of S . Set $t: h_R \rightarrow F$ given by sending $\varphi: R \rightarrow A$ to the following: choose q such that φ factors as $R \rightarrow S/J_q \xrightarrow{\varphi_q} A$ and take $\xi_q \mapsto F(\varphi_q)(\xi_q) \in F(A)$.

Finally, we must show that t is smooth. Consider the diagram

$$\begin{array}{ccccc}
 S & \xrightarrow{\quad} & S/\mathfrak{m}^{q+1} & & \\
 \downarrow & & \searrow \bar{\varphi}' & \searrow \bar{\varphi} & \\
 & & & S/J_q \times_A B & \longrightarrow & B \\
 & & \swarrow \psi & \swarrow \text{pr}_1 & & \downarrow \\
 R & \longrightarrow & S/J_q & \longrightarrow & A
 \end{array}$$

(Note: A dashed arrow φ also goes from S to B .)

with $B \ni \tilde{\xi} \mapsto \xi \in A$ and $S/J_q \ni \xi_q \mapsto \xi$. First, choose $\varphi: S \rightarrow B$ making the diagram commute. We may increase q such that $\varphi(\mathfrak{m}^{q+1}) = 0$, so we now have $\bar{\varphi}: S/\mathfrak{m}^{q+1} \rightarrow B$. Now consider the fiber product $S/J_q \times_A B$ and $\text{pr}_1: S/J_q \times_A B \rightarrow S/J_q$, so we obtain $\bar{\varphi}': S/\mathfrak{m}^{q+1} \rightarrow S/J_q \times_A B$. By (H_1) , we obtain some $\tilde{\tilde{\xi}} \in F(S/J_q \times_A B)$ mapping to $\tilde{\tilde{\xi}}$ and ξ_q . We may now assume that $B \rightarrow A$ is a small extension, which means that $\dim_k \ker(B \rightarrow A) = 1$, and thus pr_1 is a small extension. Therefore, either $\bar{\varphi}'$ is surjective or its image maps isomorphically via pr_1 to S/J_q , so we have ψ which gives $R \rightarrow B$ lifting our given $r \rightarrow A$.

The tricky part is to show that $F(\psi)(\psi_q) = \tilde{\tilde{\xi}}$, and this step is deliberately omitted. \square

A generalization of this is as follows. Consider a functor $\mathcal{F}: \mathcal{C} \rightarrow \text{Grpd}$. We say that \mathcal{F} satisfies the *Rim-Schlessinger condition* (RS) if

$$\mathcal{F}(A \times_B C) \rightarrow \mathcal{F}(A) \times_{\mathcal{F}(B)} \mathcal{F}(C)$$

is an equivalence whenever $C \rightarrow B$. Let $x_0 \in \mathcal{F}(k)$ and set

$$\bar{\mathcal{F}}_{x_0}: \mathcal{C} \rightarrow \text{Set} \quad A \mapsto \{(x, \alpha) \mid x \in \mathcal{F}(A), \alpha: X_0 \rightarrow x|_k\} / \cong,$$

where $(x, \alpha) \cong (x', \alpha')$ means that $\varphi: x \rightarrow x'$ such that the diagram

$$\begin{array}{ccc}
 x|_k & \xrightarrow{\varphi} & x'|_k \\
 \alpha \uparrow & & \alpha' \uparrow \\
 x_0 & \xrightarrow{\text{id}} & x_0
 \end{array}$$

commutes.

Theorem 1.0.10. *If \mathcal{F} has (RS) then $\overline{\mathcal{F}}_{x_0}$ has (H₁) and (H₂). Therefore, if $\dim t_{\overline{\mathcal{F}}_{x_0}} < \infty$ then $\overline{\mathcal{F}}_{x_0}$ has a hull.*

In this situation, $\overline{\mathcal{F}}_{x_0}$ has (H₄) if and only if $\text{Aut}_A(x) \twoheadrightarrow \text{Aut}_B(x|_B)$ whenever $A \twoheadrightarrow B$ and $x \in \mathcal{F}_{x_0}(A)$.

Example 1.0.11. Let $\mathcal{F}(A)$ be the category of representations $G \curvearrowright A^{\oplus n}$ with morphisms being isomorphisms of representations. This has (RS).

Example 1.0.12. Let $\mathcal{F}(A)$ be the category of smooth projective families of curves of genus g over A with morphisms being isomorphisms. This has (RS).

Returning to the example of representations, it turns out that $t_{\text{Def}_{\rho_0}} = H^1(G, M_{n \times n}(k))$, where G acts on $M_{n \times n}(k)$ via ρ_0 by conjugation.

Example 1.0.13. Consider $G = \mathbb{Z} \oplus \mathbb{Z}$ and ρ_0 to be the trivial representation on $k^{\oplus 2}$. Then $t_{\text{Def}_{\rho_0}} = H^1(\mathbb{Z}^2, M_2(k)) = M_2(k) \oplus M_2(k)$. Given two matrices A, B , we have the representation

$$\begin{aligned} \mathbb{Z}^2 \rightarrow \text{GL}_2(k[\varepsilon])(1, 0) &\mapsto \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} + \varepsilon A \\ (0, 1) &\mapsto \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} + \varepsilon B. \end{aligned}$$

We get a hull R with $h_R \rightarrow \text{Def}_{\rho_0}$. We know that R is a some quotient of $k[[a_{11}, \dots, a_{22}, b_{11}, \dots, b_{22}]]$ with ρ looking like

$$(1, 0) \mapsto \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} + A \quad (0, 1) \mapsto \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} + B,$$

and of course R is the quotient of the power series ring by the ideal generated by the coefficients of $AB - BA$.

Ivan and Cailan (Oct 1): Deformations of schemes

2.1 Deformations of affine schemes

We are looking for a Cartesian diagram of schemes

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \pi \\ \text{Spec } k & \longrightarrow & S \end{array}$$

where π is flat and surjective and S is surjective. This is called a *deformation* of X over S . For the beginning of this lecture (the part given by Ivan), we are interested in $S = \text{Spec } A$, where $A \in \mathcal{C}^*$ (this category was defined in the previous lecture). This case is called a *local deformation*, and in the face where A is Artinian, it is called an *infinitesimal deformation*.

For the ring theorists, we will make the following digression. Let A be a ring and $I \subset A$ be an ideal with $I^2 = 0$. Suppose that \bar{B} is an A/I -algebra, J is an \bar{B} -module, and $h: I \rightarrow J$ is an A -module map. Then we are interested in a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & J & \longrightarrow & ? & \longrightarrow & \bar{B} \\ & & \uparrow h & & \uparrow & & \uparrow \\ 0 & \longrightarrow & I & \longrightarrow & A & \longrightarrow & A/I \longrightarrow 0, \end{array}$$

which we will call a deformation of A . Here are some interesting questions:

1. Is such a deformation unique?
2. If \bar{B} is flat over A/I , does that mean that B is flat over A ?

Returning to the case of schemes, we will say that two deformations $\mathcal{X}, \mathcal{X}'$ of X over S are isomorphic if there exists an S -isomorphism $\phi: \mathcal{X}' \rightarrow \mathcal{X}$ commuting with the inclusions of the central fibers $X \rightarrow \mathcal{X}, \mathcal{X}'$.

Example 2.1.1. The most basic example of a family is the trivial deformation

$$\begin{array}{ccc} X & \longrightarrow & X \times_k S \\ \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & S. \end{array}$$

Definition 2.1.2. A scheme X is *rigid* if all deformations of X are isomorphic to the trivial deformation.

Theorem 2.1.3. If X is a smooth affine k -scheme and $S = \text{Spec } A$ for some local Artinian ring, then X is rigid.

Definition 2.1.4. A closed immersion $i: S_0 \hookrightarrow S$ of schemes is called a *first (resp. n th) order thickening* if the ideal sheaf $\mathcal{I} = \ker(i^b: \mathcal{O}_S \rightarrow \mathcal{O}_{S_0})$ satisfies $\mathcal{I}^2 = 0$ (resp. $\mathcal{I}^{n+1} = 0$).

Definition 2.1.5. A morphism $f: X \rightarrow S$ is called *formally smooth* (resp. *unramified*, resp. *étale*) if for all first order thickenings $i: T_0 \rightarrow T$ of affine schemes and diagrams

$$\begin{array}{ccc} T_0 & \xrightarrow{u_0} & X \\ \downarrow i & \nearrow \tilde{u}_0 & \downarrow f \\ T & \longrightarrow & S \end{array}$$

there exists a lift \tilde{u}_0 (resp. there is at most one such \tilde{u}_0 , resp. there exists a unique \tilde{u}_0).

Example 2.1.6.

1. Open immersions are formally étale. This is clear because T_0, T have the same underlying topological space.
2. Closed immersions are formally unramified. This is clear because $X \rightarrow S$ induces an injection on T -points.
3. $\mathbb{A}_S^n \rightarrow S$ is formally smooth. To see this, assume $S = \text{Spec } R$ is affine and then consider the corresponding lifting problem in commutative algebra.

Proposition 2.1.7. The classes of formally smooth (resp. étale, resp. unramified) morphisms are closed under base change, composition, and products and local on both source and target.

Definition 2.1.8. A $f: X \rightarrow S$ is *smooth* if it is formally smooth and locally of finite presentation.

We will now consider differentials. Let $X = \text{Spec } A$ be an affine scheme over k and choose a k -point and consider the diagram

$$\begin{array}{ccc} \text{Spec } k & \longrightarrow & X \\ \downarrow & & \downarrow \\ \text{Spec } k[\varepsilon] & \longrightarrow & \text{Spec } k. \end{array}$$

If X is smooth, then there exists a lift $\text{Spec } k[\varepsilon] \rightarrow X$. But this is given by a morphism

$$\tilde{\phi}: A \rightarrow k[\varepsilon]/\varepsilon^2 \quad a \mapsto \phi(a) + d(a)\varepsilon.$$

This motivates the following definition:

Definition 2.1.9. Let $R \rightarrow A$ be a morphism of rings and M be an A -module. A *derivation* $d: A \rightarrow M$ is an A -linear map satisfying the Leibniz rule.

Proposition 2.1.10. *There exists an A -module $\Omega_{A/k}^1$ equipped with a derivation $d: \Omega_{A/k}^1$ that is universal among derivations from A . This means that all derivations $\tilde{d}: A \rightarrow M$ factor through d , and formally, we have an identity*

$$\text{Der}_R(A, M) \simeq \text{Hom}_A(\Omega_{A/k}^1, M).$$

Definition 2.1.11. For an A -module M with derivation $d: A \rightarrow M$, define the ring $A[M]$ as the module $A \oplus M$ with the multiplication

$$(a, m) \cdot (a', m') = (aa', am' + a'm).$$

There is a sequence $\phi: A \rightarrow A[M] \rightarrow A$.

Proposition 2.1.12. *Let $S \leftarrow R \rightarrow A \rightarrow B$ be a diagram of rings. Then*

1. $\Omega_{A \otimes_R S/S}^1 \simeq \Omega_{A/R}^1 \otimes_R S$;
2. The sequence $\Omega_{A/R}^1 \otimes_A B \rightarrow \Omega_{B/R}^1 \rightarrow \Omega_{B/A}^1 \rightarrow 0$ is exact.
3. If $B = A/I$ for some ideal I , we have an exact sequence

$$I/I^2 \rightarrow \Omega_{A/R}^1 \otimes_A B \rightarrow \Omega_{B/R}^1 \rightarrow 0.$$

4. For all $f \in A$, we have $\Omega_{A[f^{-1}]/R}^1 \simeq \Omega_{A/R}^1 \otimes_A A[f^{-1}]$.

Remark 2.1.13. If $J = \ker(A \otimes_R A \rightarrow A)$, then $\Omega_{A/R}^1 = J/J^2$.

Theorem 2.1.14. *Let $f: X \rightarrow S$ be locally of finite presentation. The following are equivalent:*

1. f is smooth;
2. f is flat with smooth fibers;
3. f is flat and has smooth geometric fibers.

We will finally return to deformation theory.

Lemma 2.1.15. *Let Z_0 be a closed subscheme of Z determined by a nilpotent ideal sheaf \mathcal{N} . If Z_0 is affine, then so is Z .*

Proof of this result can be found in EGA, Chapter I.5.9.

Proof of Theorem 2.1.3. Recall that we have a diagram of the form

$$\begin{array}{ccc} B & \longrightarrow & B_0 \\ \uparrow & & \uparrow \\ A & \longrightarrow & k, \end{array}$$

where $A \rightarrow B$ is flat and $B_0 \simeq B \otimes_A k$ is a smooth k -algebra. We need to prove that $B_0 \simeq B \otimes_A k$. The first step is to prove this result for first-order deformations. Suppose that $A = k[\varepsilon]$ is a square-zero extension.

Lemma 2.1.16. *For a ring R with M, N flat over R , nilpotent ideal $I \subset R$, and $f: M \rightarrow N$, then if $f \otimes_R R/I$ is an isomorphism, then so is f .*

To prove the lemma, note that the cokernel of f is preserved by I , so it must vanish. Returning to our case, we know that B is a smooth $k[\varepsilon]$ -algebra. Now we obtain a square-zero extension $B_0[\varepsilon]$ of B_0 and a diagram

$$\begin{array}{ccc} B & \longrightarrow & B_0 \\ & & \uparrow \\ k[\varepsilon] & \xrightarrow{f} & B_0[\varepsilon] \end{array}$$

with a lift $B \rightarrow B_0[\varepsilon]$. But now by the lemma, we have $B \otimes_{k[\varepsilon]} k = B_0[\varepsilon] \otimes_{k[\varepsilon]} k$. The rest of the proof follows using an inductive argument that was verbalized but now written down. \square

2.2 Deformations of schemes

The main theorem of this section is

Theorem 2.2.1. *Assume X is a smooth R -scheme. Then there is a bijection*

$$\text{Def}_X^{\text{sm}}(k[I]) \simeq H^1(X, T_{X/k} \otimes I).$$

Proof. Let X' be a smooth deformation over $k[I]$. Then the diagram

$$\begin{array}{ccc} X & \longrightarrow & X' \\ \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \text{Spec } k[I] \end{array}$$

is cartesian. Then if $U_k = \text{Spec } B_k$ is an affine cover of X and $U'_k = \text{Spec } D_k$ is an affine cover of X' , we have a $k[I]$ -linear ring isomorphism

$$\varphi_k: k[I] \otimes_k B_k \rightarrow D_k \quad (k, i) \otimes b \mapsto s(b) + i.$$

Modulo I , φ_k is the identity on B_k . Without loss of generality, we may assume that $U_{kj} = U_k \cap U_j$ is a distinguished open for both U_k and U_j , so let $U_{kj} = \text{Spec } B_{kj}$ and $U'_{kj} = \text{Spec } D_{kj}$. Now note that both

$$\varphi_k, \varphi_j: k[I] \otimes_k B_{kj} \rightarrow D_{kj}$$

induce the identity on B_{kj} modulo I . Now we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \longrightarrow & D_{kj} & \longrightarrow & B_{kj} \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \varphi_j^{-1} \varphi_k & & \downarrow \text{id} \\ 0 & \longrightarrow & I & \longrightarrow & D_{kj} & \longrightarrow & B_{kj} \longrightarrow 0. \end{array}$$

Lemma 2.2.2. *The morphism $g = \varphi_j^{-1} \varphi_k$ must be of the form*

$$g(i + b) = i + b + \delta(b),$$

where $\delta: B_{kj} \rightarrow I$ is a derivation.

In particular, this means that $\varphi_j^{-1} \circ \varphi_k(b, b') = (b, \alpha_{kj}(b) + b')$, where $\alpha_{kj}: B_{kj} \rightarrow I \otimes_k B_{kj}$ is a derivation.

By definition, we have

$$\begin{aligned} (T_{X/k} \otimes_k I)(B_{kj}) &= \text{Hom}_{B_{kj}}(\Omega_{B_{kj}/k}^1, B_{kj}) \otimes_k I \\ &= \text{Hom}_{B_{kj}}(\Omega_{B_{kj}/k}^1, B_{kj} \otimes_k I) \\ &= \text{Der}_k(B_{kj}, B_{kj} \otimes_k I). \end{aligned}$$

Therefore, $\alpha_k \in H^0(B_{kj}, T_X \otimes_k I)$. Note that

$$\varphi_\ell^{-1} \circ \varphi_j \circ \varphi_j^{-1} \circ \varphi_k^{-1} = \varphi_\ell^{-1} \circ \varphi_k,$$

which implies that

$$(b, \alpha_{j\ell}(b) + \alpha_{kj}(b) + b') = (b, \alpha_{k\ell}(b) + b')$$

and thus $\{\alpha_{kj}\} \in Z^1(\{U_k\}, T_X \otimes_k I)$.

If two deformations are the same, note that φ_k is defined using a ring section $s_k: B_k \rightarrow D_k$ of the canonical map $\pi_k: D_k \rightarrow B_k$. If φ'_k is defined using another section s'_k , then define $\theta_k = s'_k - s_k \in \text{Der}(B_k, I \otimes_k B_k)$. We now compute

$$((\varphi'_j)^{-1} \circ \varphi'_k - \varphi_j^{-1} \circ \varphi_k)(b, b') = (0, \theta_k(b) - \theta_j(b)),$$

and thus the two differ by the desired coboundaries. \square

We will now consider some obstructions. We are looking for a diagram of the form

$$\begin{array}{ccc} X' & \longrightarrow & X'' \\ \downarrow f & & \downarrow \\ \text{Spec } A' & \longrightarrow & \text{Spec } A'' \end{array}$$

for each pair (j, k) , we have a isomorphism $\psi_{jk}: V'_j \rightarrow V'_k$ and a cocycle

$$c_{jkl} = \psi_{kl} \circ \psi_{jk} \circ \psi_{jl}^{-1}.$$

This induces $B_{jkl} \in \text{Der}_A(D_{jkl}, J \otimes_A D_{kl}) = Z^2(U, T_{X'/A} \otimes_A J)$.

Now we will discuss some examples.

Theorem 2.2.3. *Let C be a smooth projective curve, $T = T_C$, and $K = \Omega_C^1$. We have the following table:*

Table 2.1: Cohomology

	degree	h^0	h^1	h^2	
K	$2g-2$	g	1	0	where $\varepsilon = 0$ where $g \geq 2$, $\varepsilon = 1$ if $g = 1$, and $\varepsilon = 3$ if $g = 0$.
T	$2-2g$	ε	$\varepsilon + 3g - 3$	0	

For $g \geq 2$, $\deg T < 0$, and by Riemann-Roch and Serre duality, we have $h^1(C, T_C) = 3g - 3$.

Theorem 2.2.4. \mathbb{P}^n has no infinitesimal deformations.

Proof. Consider the Euler sequence

$$0 \rightarrow \mathcal{O} \rightarrow \mathcal{O}(1)^{\oplus n+1} \rightarrow T_{\mathbb{P}^n} \rightarrow 0$$

and use the long exact sequence in cohomology. Because positive degree line bundles have no higher cohomology, we have $H^1(T_{\mathbb{P}^n}) = 0$. \square

Kevin (Oct 08): Deformations of coherent sheaves

There will be no mixed characteristic funny business during this lecture. Let X be a projective k -scheme (proper might be fine, but this makes certain facts more true) and \mathcal{F} be a coherent sheaf on X . Consider the deformation functor

$$D_{\mathcal{F}}: \text{Art}_k \rightarrow \text{Set} \quad A \mapsto \{\mathcal{F}_A \in \text{Coh}(X_A) \mid \mathcal{F}_A|_X \cong \mathcal{F}, \mathcal{F}_A \text{ flat over } A\}.$$

We want to study the properties of this functor, which means we will check Schlessinger's conditions:¹ Let $A \rightarrow B \leftarrow C$ be a diagram in \mathcal{C} and consider the morphism

$$D(B \times_A C) \xrightarrow{r} D(B) \times_{D(A)} D(C).$$

- (H₁) The morphism r is surjective if $C \rightarrow A$;
- (H₂) The morphism r is bijective if $C = k[\varepsilon] \rightarrow k = A$;
- (H₃) $\dim_k(t_D) < \infty$ (later, we will see that we need H₂ for formulate this). Here, t_D is the tangent space to D ;
- (H₄) The morphism r is bijective if $C \rightarrow A$.

Recall from Johan's lecture that (H₁), (H₂), (H₃) are equivalent to the existence of a hull and (H₃), (H₄) are equivalent to D being pro-representable.

We only need to check (H₁) for small extensions, which are extensions by a k -vector space

$$0 \rightarrow I \rightarrow C \rightarrow A \rightarrow 0,$$

where I is killed by the maximal ideal of C .

Theorem 3.0.1. *The functor $D_{\mathcal{F}}$ admits a hull.*

Lemma 3.0.2. *Let (A, \mathfrak{m}) be a local Artinian ring.*

1. *If $\mathfrak{m}M = M$, then $M \cong 0$.*
2. *If $M \rightarrow N$ induces an isomorphism $M/\mathfrak{m}M \cong N/\mathfrak{m}N$ and N is flat over A , then $M \cong N$.*
3. *If M is flat, then M is free.*

¹Neither Kevin nor Johan knows why these conditions are called H

Proof. We know that $m^d = 0$, so $m^d M = 0$, and thus $M = mM = m^2 M = \dots = m^d M = 0$. Next, suppose $M \rightarrow N$ induces an isomorphism after killing m . Then we know that the kernel and cokernel vanish because they are killed by m , so $M \rightarrow N$ must be an isomorphism. The last part is left as an exercise. \square

Proof of theorem. We will simply prove $(H_1), (H_2), (H_3)$:

1. Suppose that $C \rightarrow A$ is a small extension and consider a pair $(\mathcal{F}_B, \mathcal{F}_C) \in D(B) \times_{D(A)} D(C)$. We know that we have isomorphisms $\mathcal{F}_B|_{X_A} \cong \mathcal{F}_A, \mathcal{F}_C|_{X_A} \cong \mathcal{F}_A$, and so we take the fiber product

$$\mathcal{F}_{B \times_A C} := \mathcal{F}_B \times_{\mathcal{F}_A} \mathcal{F}_C.$$

We only need to show that our sheaf is flat over $B \times_A C$ because it clearly restricts to \mathcal{F}_B and \mathcal{F}_C . We can consider each sheaf as a module M , and so we know M_B is free over B by the lemma. Choose a basis $\{e_i\}$. Also consider the diagram

$$\begin{array}{ccc} M_B \times_{M_A} M_C & \longrightarrow & M_C \\ \downarrow & & \downarrow v \\ M_B & \xrightarrow{u} & M_A. \end{array}$$

Then M_A has A -basis $u(e_i)$. Because M_C surjects onto M_A , we can lift the $u(e_i)$ to $f_i \in C$, and these form a C -basis for M_C . This all implies that $M_B \times_{M_A} M_C$ is free with basis (e_i, f_i) .

2. It suffices to prove injectivity. Suppose $\mathcal{G} \in D(B \times_k k[\varepsilon])$ maps to $(\mathcal{F}_B, \mathcal{F}_{k[\varepsilon]}) \in D(B) \times D(k[\varepsilon])$, and so we have morphisms

$$\begin{array}{ccc} \mathcal{G} & \longrightarrow & \mathcal{F}_{k[\varepsilon]} \\ \downarrow & & \downarrow \\ \mathcal{F}_B & \longrightarrow & \mathcal{F}. \end{array}$$

We will prove that this diagram is Cartesian. By the lemma, the morphism $\mathcal{G} \rightarrow \mathcal{F}_B \times_{\mathcal{F}} \mathcal{F}_{k[\varepsilon]}$ is an isomorphism.

3. We will prove that $T_D = \text{Ext}_X^1(\mathcal{F}, \mathcal{F})$. We will only prove this in the case where \mathcal{F} is a vector bundle \mathcal{E} of rank r . In this case, we have $\text{Ext}_X^1(\mathcal{E}, \mathcal{E}) = H^1(X, \text{End}(\mathcal{E}))$. Now we will associate cocycles to deformations. To each $\mathcal{E}_{k[\varepsilon]}$, we will associate an open cover (U_j) and

$$h_{ij} \in \text{Aut}(\mathcal{O}_{X_{k[\varepsilon]}}^{\oplus r})(U_{ij}),$$

and we write $g_{ij} + \varepsilon f_{ij}$, where $g_{ij} \in \text{Aut}_{\mathcal{O}_X^{\oplus r}}(U_{ij})$ and $f_{ij} \in \text{End}(\mathcal{O}_X^{\oplus r})(U_{ij})$. The cocycle condition is that

$$g_{ik} + \varepsilon f_{ik} = (g_{ij} + \varepsilon f_{ij})(g_{jk} + \varepsilon f_{jk}),$$

which is the same as

$$f_{ik} = g_{ij} f_{jk} + f_{ij} g_{jk},$$

which is exactly the Čech 1-cocycle condition. Proving that equivalent cocycles give the same deformation is easy. \square

Theorem 3.0.3. *The condition (H_4) holds when \mathcal{F} is simple, which means that $k \simeq \text{End}_X(\mathcal{F})$.*

3.1 Tangent-obstruction theory

Suppose D is a deformation functor. Then a *tangent-obstruction theory* for D is given by finite-dimensional k -vector spaces (T^1, T^2) . Suppose we have a small extension

$$0 \rightarrow I \rightarrow B \rightarrow A \rightarrow 0.$$

Then we have another exact sequence

$$T^1 \otimes_k I \rightarrow D(B) \rightarrow D(A) \xrightarrow{\text{ob}} T^2 \otimes_k I,$$

which means that

1. $\xi_A \in D(A)$ lifts to $D(B)$ if and only if $\text{ob}(\xi_A) = 0$;
2. $T^1 \otimes I$ acts transitively on the fibers of $D(B) \rightarrow D(A)$;
3. If $A = k$, then the action of $T^1 \otimes I$ acts simply transitively on $D(B)$.

Note that because T^1 acts simply transitively on $D(k[\varepsilon])$, we must have $T^1 = D(k[\varepsilon])$. On the other hand, T^2 is not canonical.

Theorem 3.1.1. *The deformations $D_{\mathcal{F}}$ admits a tangent-obstruction theory with $T^1 = \text{Ext}_X^1(\mathcal{F}, \mathcal{F})$ and $T^2 = \text{Ext}_X^2(\mathcal{F}, \mathcal{F})$.*

Proof. We claim that if D satisfies (H_1) and (H_2) , then $D(k[\varepsilon]) \otimes I$ naturally acts on $D(B)$ for small extensions $0 \rightarrow I \rightarrow B \rightarrow A \rightarrow 0$. To see this, note that $D(k[\varepsilon]) \otimes_k I = D(k[I])$. We also note that by (H_2) , $D(k[I]) \times D(B) = D(k[I] \times_k B)$. Now define $\alpha: k[I] \times_k B \rightarrow B$ by $\alpha(1+i, b) = 1+b$, and this gives us an action of $D(k[I]) \times D(B) = D(k[I] \times_k B) \xrightarrow{\alpha_*} D(B)$. To prove transitivity, apply (H_1) to the diagram

$$\begin{array}{ccc} k[I] \times_k B & \xrightarrow{\alpha} & B \\ \downarrow \pi_B & & \downarrow \\ B & \longrightarrow & A. \end{array}$$

Now we will consider obstructions. We will assume again that \mathcal{F} is a rank r vector bundle, which we will call \mathcal{E} . Let

$$0 \rightarrow I \rightarrow B \rightarrow A \rightarrow 0$$

be a small extension, so we will consider $H^2(X, \text{End}(\mathcal{E}))$. Consider an open cover (U_i) and $g_{ij} \in \text{Aut}(\mathcal{O}_X^{\oplus r} \otimes_k A)(U_{ij})$. We want to lift these to $h_{ij} \in \text{Aut}(\mathcal{O}_X^{\oplus r} \otimes_k B)(U_{ij})$. If this is possible, we have a cocycle

$$h_{ij}^{-1} h_{ij} h_{jk} \in 1 + \text{End}(\mathcal{O}_X^{\oplus r} \otimes_k I)(U_{ij}),$$

and the cocycle condition is satisfied when $h_{ij}^{-1} h_{ij} h_{ik} = 1$. If any other $h'_{ij} = h_{ij} + s_{ij}$, then we note that

$$(h'_{ij})^{-1} h'_{ij} h'_{jk} = h_{ik}^{-1} h_{ij} h_{jk} + (-s_{ik} g_{ij} g_{jk} + g_{ik}^{-1} s_{ij} g_{ij} + g_{ik}^{-1} g_{ij} s_{jk}),$$

and this gives us a class in $H^2(X, \text{End}(\mathcal{E})) \otimes I$. □

Remark 3.1.2. Let R be the hull of D , which means we have a morphism $h_R \rightarrow D$. Then we know $R = k[[t_1, \dots, t_{d_1}]]/(f_1, \dots, f_{d_2})$. We also know that $d_1 - d_2 \leq \dim R \leq d_1$.

Example 3.1.3 (Good example). Let X be a smooth projective curve and \mathcal{E} be a rank r vector bundle. Then we know that

$$T^1 = H^1(X, \text{End}(\mathcal{E})), \quad T^2 = H^2(X, \text{End}(\mathcal{E})) = 0,$$

so deformations of \mathcal{E} are unobstructed. Now assume that \mathcal{E} is simple. Then $H^0(X, \text{End}(\mathcal{E})) = k$ by definition, and we also know that D_ε is pro-represented by some ring R with

$$\dim R = h^1(X, \text{End}(\mathcal{E})) = r^2(g-1) + 1$$

by Riemann-Roch.

Example 3.1.4 (Bad example). Let X be a smooth projective variety and \mathcal{E} be a rank r vector bundle on X . Let $\varepsilon_1, \varepsilon_2 \in D_\varepsilon(k[\varepsilon])$. Then $(\varepsilon_1, \varepsilon_2) \in D_\varepsilon(k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_1\varepsilon_2, \varepsilon_2^2))$, and we would like to lift to $D_\varepsilon(k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_2^2))$.

We will compute the obstruction explicitly. We know $\varepsilon_1, \varepsilon_2$ give us classes $u_1, u_2 \in H^1(X, \text{End}(\mathcal{E}))$, and after some magical computation, the obstruction to lifting is given by

$$u_1 \smile u_2 + u_2 \smile u_1,$$

where the cup product comes from the algebra structure on $\text{End}(\mathcal{E})$.

Now let $X = C_1 \times C_2$ be a product of curves. Then $H^1(X, \mathcal{O}_X) = H^1(C_1, \mathcal{O}_{C_1}) \oplus H^1(C_2, \mathcal{O}_{C_2})$ and $H^2(X, \mathcal{O}_X) = H^1(C_1, \mathcal{O}_{C_1}) \otimes H^1(C_2, \mathcal{O}_{C_2})$. Suppose that $\alpha_1 \in H^1(C_1, \mathcal{O}_{C_1})$ and $\alpha_2 \in H^1(C_2, \mathcal{O}_{C_2})$ with nonzero cup product. Then we simply set $\mathcal{E} = \mathcal{O}_X \oplus \mathcal{O}_X$ and

$$u_1 = \begin{pmatrix} 0 & \alpha_1 \\ 0 & 0 \end{pmatrix} \quad u_2 = \begin{pmatrix} 0 & 0 \\ \alpha_2 & 0 \end{pmatrix} \quad u_1 \smile u_2 + u_2 \smile u_1 = \begin{pmatrix} \alpha_1 \smile \alpha_2 & \\ & \alpha_2 \smile \alpha_1 \end{pmatrix}.$$

this gives us our obstructed deformation.

Patrick (Oct 15): Deformations of singularities

We begin by fixing some notation. Let k be a field and $R = P/I$, where $P = k[x_1, \dots, x_n]$ and $I = (f_1, \dots, f_r)$ is an ideal. Throughout this lecture, we will denote local Artinian rings with residue field k by A, B, C, \dots and rings by R, S, T, \dots . Finally, denote $Z = \text{Spec } R$.

4.1 Explicit criteria for flatness

We will study (embedded) deformations of singular affine schemes embedded in \mathbb{A}^n . The first thing we want to understand is to explicitly understand flatness of some R_A over A , where $R_A \otimes_A k = R$. We will write $R_A = P_A/I_A$, where $P_A = A[x_1, \dots, x_n] = A \otimes_k P$. Recall that over a Noetherian local ring S with residue field k , a module M is flat if and only if it is free, and this is equivalent to $\text{Tor}_1^S(M, k) = 0$ by standard results in commutative algebra.

Now consider the exact sequence

$$0 \rightarrow I_A \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

After tensoring with k , we have

$$0 \rightarrow \text{Tor}_1(R_A, k) \rightarrow I_A \otimes_A k \rightarrow P \rightarrow R \rightarrow 0.$$

Therefore, we know that R_A is flat over A if and only if $I_A \otimes_A k = I$. We would like to understand this statement.

Consider a presentation

$$P_A^s \rightarrow P_A^r \rightarrow I_A \rightarrow 0$$

of I_A . Then we know R_A is flat over A if and only if after tensoring with k , we obtain an exact sequence

$$P^s \rightarrow P^r \rightarrow I \rightarrow 0.$$

Note that to give this presentation $P^s \rightarrow P^r \rightarrow I \rightarrow 0$ is the same as giving a complete set of relations among the generators of I .

Proposition 4.1.1. *Suppose that*

$$(4.1) \quad P^s \rightarrow P^r \rightarrow P \rightarrow R \rightarrow 0$$

is exact and

$$(4.2) \quad P_A^s \rightarrow P_A^r \rightarrow P_A \rightarrow R_A \rightarrow 0$$

is a complex such that $P_A^r \rightarrow R_A \rightarrow R_A \rightarrow 0$ is exact and tensoring (2) with k gives (1). Then R_A is flat over A .

Proof. Note that the hypotheses are equivalent to the fact that all relations in I can be lifted to I_A . Now given $g'_1, \dots, g'_r \in P_A$ such that

$$\sum_{i=1}^r g'_i f'_i = 0,$$

this clearly descends to a relation in I by killing the maximal ideal of A . But now if we choose a complete set of relations for I_A , this descends to a complete set of relations in I , so we may in fact assume that (2) is exact.

In this case, there exists some L_A such that the sequence splits as

$$P_A^s \rightarrow L_A \rightarrow 0 \quad 0 \rightarrow L_A \rightarrow P_A^r \rightarrow I_A \rightarrow 0 \quad 0 \rightarrow I_A \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

By right exactness of the tensor product, we know $P_A^s \otimes k \rightarrow L_A \otimes k \rightarrow 0$ is exact. We also know that

$$L_A \otimes k \rightarrow P_A^r \otimes k \rightarrow I_A \otimes k \rightarrow 0$$

is exact, again by right exactness. But this means that $I_A \otimes k$ is the cokernel of $P^s \rightarrow P^r$, and therefore $I_A \otimes k = I$. This means that R_A is flat. \square

Corollary 4.1.2. *Let $R = P/I$ and $R_A = P_A/I_A$, where $I = (f_1, \dots, f_r)$ and $I_A = (f'_1, \dots, f'_r)$ such that f'_i is a lift of f_i . Then R_A is flat over A if and only if every relation among the f_i lifts to a relation among the f'_i .*

Remark 4.1.3. This result essentially gives us that first-order embedded deformations of $\text{Spec } R \subset \mathbb{A}^n$ are given by $\text{Hom}(I, R)$. The first-order (not embedded) deformations of Z are given by the cokernel of

$$0 \rightarrow T_X \rightarrow T_{\mathbb{A}^n}|_X \rightarrow N_{X/\mathbb{A}^n},$$

which arises from the exact sequence

$$I/I^2 \rightarrow \Omega_{\mathbb{A}^n}^1|_X \rightarrow \Omega_X^1 \rightarrow 0,$$

and this is supported on the singular points of X , so when X has isolated singularities, this is finite-dimensional.

Note that if $\text{Spec } R \subset \mathbb{A}^n$ is a complete intersection, then I is generated by a regular sequence, so in particular the Koszul complex is a free resolution of R and therefore there are only trivial relations among the f_i (this means the relations are generated by $f_i f_j - f_j f_i = 0$). Clearly, because we are only considering commutative rings (after all, this is normal algebraic geometry), this means that all deformations of $\text{Spec } R$ are unobstructed.

4.2 Hilbert schemes of smooth surfaces

We will prove that deformations of finite length closed subschemes of \mathbb{A}^2 are unobstructed. In particular, this will imply that the Hilbert scheme $\text{Hilb}(\mathbb{A}^2, n)$ is smooth.

Let $Z \subset \mathbb{A}^2$ be a closed subscheme of dimension 0. Then because $P = k[x, y]$ has dimension 2, there exists a free resolution

$$0 \rightarrow P^s \xrightarrow{(g_{ij})} P^r \rightarrow P \rightarrow R \rightarrow 0$$

of R . In this case it is possible to understand the matrix (g_{ij}) , and in fact this is the special case of a more general result. First, when we study the local behavior, we have the following result.

Theorem 4.2.1 (Hilbert, Burch). *Let P be a regular local ring of dimension n and $R = P/I$ be a Cohen-Macaulay quotient of codimension 2. Then there exists an $(r-1) \times r$ matrix $G = (g_{ij})$ whose maximal minors f_1, \dots, f_r minimally generate I , and there is a free resolution*

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0.$$

Proof. Note that the fact that the free resolution has this length is a corollary of the Auslander-Buchsbaum formula, which says that for a ring R and module M , we have

$$\text{depth } M + \text{proj. dim } M = \text{depth } R$$

and the fact that depth equals dimension for Cohen-Macaulay things. Thus we have a free resolution

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(a_i)} P \rightarrow R \rightarrow 0,$$

where a_1, \dots, a_r are a minimal set of generators for I . Let f_i is $(-1)^i$ times the determinant of the i -th minor of g_{ij} . We will prove that the map (f_i) is the same as the map (a_i) ; clearly

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0.$$

is a resolution. This is because at the generic point of P , we know (g_{ij}) is injective, so at least one f_i is nonzero. But then we know $\text{coker}(g_{ij})$ is torsion-free (because I is torsion-free), and so it in fact must vanish by rank reasons. Thus (a_1, \dots, a_r) and (f_1, \dots, f_r) are isomorphic as P -modules.

At a codimension 1 point in $\text{Spec } P$, note that $0 \rightarrow P^{r-1} \rightarrow P^r \xrightarrow{(a_i)} P \rightarrow B \rightarrow 0$ is split exact (because I has codimension 2). This implies that at least one f_i is a unit, and thus (f_1, \dots, f_r) has codimension at least 2. But then the isomorphism $I \cong (f_1, \dots, f_r)$ is given by multiplication by some nonzero element of P which is a unit away from codimension 2. But this means it is a unit everywhere. \square

Considering the global picture in \mathbb{A}^n , we obtain the following result.

Theorem 4.2.2 (Hilbert, Schaps). *Let $Z = \text{Spec } R \subset \mathbb{A}^n$ be a Cohen-Macaulay closed subscheme of codimension 2. Then $R = P/I$ has a free resolution of the form*

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0$$

where the f_i are the maximal minors of the matrix (g_{ij}) .

This result in fact holds over any Artinian local ring A , which we will use later.

Next, we want to understand what happens if we choose some Artinian local ring with residue field k and lift the g_{ij} to g'_{ij} , where $g'_{ij} \in P_A$.

Theorem 4.2.3 (Schaps). *If A is a square zero extension of k , then the sequence*

$$0 \rightarrow P_A^{r-1} \xrightarrow{(g'_{ij})} P_A^r \xrightarrow{(f'_i)} P_A \rightarrow R_A \rightarrow 0$$

is exact. Moreover, any lifting of R over A arises by lifting the matrix (g_{ij}) .

Proof. We know that

$$L_A^\bullet := P_A^{r-1} \rightarrow P_A^r \rightarrow P_A$$

is a complex. This is because composing the two maps amounts to evaluating determinants with a repeated column. Because P_A is free (and therefore flat), we can tensor with the exact sequence

$$0 \rightarrow \mathfrak{m}_A \rightarrow A \rightarrow k \rightarrow 0$$

to obtain an exact sequence of complexes

$$0 \rightarrow L_A^\bullet \otimes_A \mathfrak{m}_A \rightarrow L_A^\bullet \rightarrow L_A^\bullet \otimes_A k \rightarrow 0.$$

Note that

$$L_A^\bullet \otimes_A k = P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P = L^\bullet.$$

In particular, this term is exact by Hilbert-Schaps. In addition, clearly $L_A^\bullet \otimes_A \mathfrak{m}_A = L^\bullet \otimes_k \mathfrak{m}_A$ because $A \rightarrow k$ is a square zero extension, so the complex $L_A^\bullet \otimes_A \mathfrak{m}_A$ is exact. By the long exact sequence in homology, we know that L_A^\bullet is exact. Note that L^\bullet extends to an exact sequence

$$0 \rightarrow P^{r-1} \rightarrow P^r \rightarrow P \rightarrow R \rightarrow 0,$$

and L_A^\bullet extends to an exact sequence

$$0 \rightarrow P_A^{r-1} \rightarrow P_A^r \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

However, by the homology long exact sequence, we have an exact sequence

$$0 \rightarrow R \otimes_k \mathfrak{m}_A \rightarrow R_A \rightarrow R \rightarrow 0.$$

But this implies that $R_A \otimes_A k = R$. Finally, by the local criterion for flatness, we see that R_A is flat over A .

Let $R_A = P_A/I_A$ be a lifting of R over A . Lift $f_i \in I$ to $h_i \in I_A$. By Nakayama, these generate I_A , so we obtain a free resolution

$$0 \rightarrow P_A^{r-1} \xrightarrow{(g'_{ij})} P_A^r \xrightarrow{(h_i)} P_A \rightarrow R_A \rightarrow 0,$$

where g'_{ij} lift the g_{ij} . However, we already have a lift

$$0 \rightarrow P_A^{r-1} \xrightarrow{(g'_{ij})} P_A^r \xrightarrow{(f'_i)} P_A \rightarrow R'_A \rightarrow 0,$$

and so we must show $R_A = R'_A$. But we know that the ideals $I_A = (h_1, \dots, h_r)$ and $I'_A = (f'_1, \dots, f'_r)$ are isomorphic as P_A -modules. But then if we restrict this isomorphism to $\mathbb{A}_A^n \setminus \text{supp } B$, we obtain a unit in $H^0(\mathbb{A}_A^n \setminus \text{supp } B, \mathcal{O}_{\mathbb{A}_A^n})$. Because functions extend over codimension 2, we have $H^0(\mathbb{A}_A^n \setminus \text{supp } B, \mathcal{O}_{\mathbb{A}_A^n}) = P_A$, so this is a global unit. This gives the desired result. \square

This result holds if we replace $A \rightarrow k$ with any square-zero extension of Artinian local rings $B \rightarrow A$ and P, P_A with flat things, and so we see that (embedded) deformations of codimension 2 Cohen-Macaulay subschemes of \mathbb{A}^n are unobstructed. In particular, any dimension 0 closed subscheme $Z \subset \mathbb{A}^2$ is automatically Cohen-Macaulay (because it is dimension 0), so its embedded deformations are unobstructed. By some cohomological argument, the tangent space to $\text{Hilb}(\mathbb{A}^2, n)$ is isomorphic to $\text{Hom}(R, R)$ and has dimension $2n$, so

4.3 An obstructed deformation

Let $R = k[x, y, z]/(z^2, xy, xz, yz)$. Note that this scheme has an embedded point at the origin, so in particular it is **not** Cohen-Macaulay.




Figure 4.1: Drawing of $\text{Spec } R$

We will study embedded deformations of $\text{Spec } R$ and see that they are obstructed. In particular, we will choose two deformations of R over $k[\varepsilon]$ that cannot be simultaneously lifted. We claim that a complete set of relations (using the ordering (xy, xz, yz, z^2) for the generators of I) is given by the matrix

$$G = \begin{pmatrix} z & -y & 0 & 0 \\ z & 0 & -x & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y \end{pmatrix}.$$

Now a first-order deformation of $\text{Spec } R$ is given by lifting (xy, xz, yz, z^2) over $k[\varepsilon]$, and the first candidate is to consider $I_{\varepsilon_1} = (xy + \varepsilon_1 y, xz, yz, z^2)$. Then we note that

$$G \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz \\ z^2 \end{pmatrix} = \varepsilon_1 \begin{pmatrix} yz \\ yz \\ 0 \\ 0 \end{pmatrix},$$

and we can lift G to kill this vector with the matrix

$$G_{\varepsilon_1} = \begin{pmatrix} z & -y & -\varepsilon_1 & 0 \\ z & 0 & -x - \varepsilon_1 & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y \end{pmatrix} = G + \begin{pmatrix} 0 & 0 & -\varepsilon_1 & 0 \\ 0 & 0 & -\varepsilon_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} =: G + G_1.$$

Next consider the deformation given by $I_{\varepsilon_2} = (xy, xz, yz + \varepsilon_2 z, z^2)$. We note that

$$G \begin{pmatrix} xy \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} = \varepsilon_2 \begin{pmatrix} 0 \\ -xz \\ 0 \\ z^2 \end{pmatrix},$$

and we can lift G to kill this vector with the matrix

$$G_{\varepsilon_2} = \begin{pmatrix} z & -y & 0 & 0 \\ z & \varepsilon_2 & -x & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y - \varepsilon_2 \end{pmatrix} = G + \begin{pmatrix} 0 & \varepsilon_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\varepsilon_2 \\ 0 & 0 & 0 & 0 \end{pmatrix} =: G + G_2.$$

Now we consider $I_{\varepsilon_1^2, \varepsilon_2^2, \varepsilon_1 \varepsilon_2} = (xy + \varepsilon_1 y, xz, yz + \varepsilon_2 z, z^2)$ and attempt to lift this deformation to $k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_2^2)$. Note that

$$\begin{aligned} (G + G_1 + G_2) \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} &= \begin{pmatrix} z & -y & -\varepsilon_1 & 0 \\ z & \varepsilon_2 & -x - \varepsilon_1 & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y - \varepsilon_2 \end{pmatrix} \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} \\ &= \varepsilon_1 \varepsilon_2 \begin{pmatrix} -z \\ -z \\ 0 \\ 0 \end{pmatrix}, \end{aligned}$$

and clearly $z \notin I$, so in fact we cannot lift this deformation to $k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_2^2)$. This proves obstructedness.

Avi (Oct 22): Local-global methods

5.1 Curves with isolated singularities

Here, we will consider the same lifting problems that we have considered before. Let X be a curve over k and consider $p: \text{Def}_X \rightarrow \mathcal{C}$ be the deformations of X as a category cofibered over \mathcal{C} .

Definition 5.1.1. A functor $F: D_1 \rightarrow D_2$ is *smooth* if given $\varphi: A' \rightarrow A$ and $(Z, A') \rightarrow F(Y, Z)$ over φ and $(Y', A') \rightarrow (Y, A)$ over φ in D_1 , then there exists some $F(Y', A') \rightarrow (Z, A')$ over the identity in A' .

There is an absolute notion, for D_1 to be smooth, where we set $D_2 = \text{Def}_k$ to be the category of trivial deformations. In this case, we also called D_1 *unobstructed*, and this corresponds to being unobstructed in the tangent-obstruction theory.

Theorem 5.1.2 (Local-to-global). *Let X/k be separated of dimension at most 1 and smooth away from finitely many points. At each singularity p_1, \dots, p_n , consider the inclusions*

$$\mathcal{O}_{X,p} \rightarrow \mathcal{O}_{X,p}^h \rightarrow \widehat{\mathcal{O}}_{X,p}$$

into the henselizations and completions, respectively. Then the functors

$$\text{Def}_X \rightarrow \prod_i \text{Def}_{\mathcal{O}_{X,p_i}} \xrightarrow{(2)} \prod \text{Def}_{\mathcal{O}_{X,p}^h} \xrightarrow{(3)} \prod \text{Def}_{\widehat{\mathcal{O}}_{X,p}}$$

are all smooth and (2), (3) induce isomorphisms on tangent spaces.

This means that we only have to check unobstructedness at the completions of the local rings at each singular point.

Proof. First, we can reduce to the affine case.¹ If X is a curve, then $X = U_1 \cup U_2$ can be covered by two affine opens such that $U_1, U_1 \cap U_2$ are smooth. Because deformations of smooth affine schemes are unobstructed, we can essentially ignore U_1 . To do this, we will prove that $\text{Def}_X \rightarrow \text{Def}_{U_2}$ is smooth.

Because smoothness (roughly) respects products, the functor

$$\text{Def}_X \rightarrow \text{Def}_{U_1} \times_{\text{Def}_{U_1 \cap U_2}} \text{Def}_{U_2}$$

¹Avi says that Johan sketched this proof and then said not to give it.

is smooth by formal reasons. Because $U_1, U_1 \cap U_2$ are smooth, we can ignore $\text{Def}_{U_1}, \text{Def}_{U_1 \cap U_2}$ (in fact, the arrow is an equivalence) and project down to Def_{U_2} .

Now $U_2 = \text{Spec } P$ is affine. Then each p_i gives a maximal ideal \mathfrak{m}_i , and so we set $J = \bigcap_i \mathfrak{m}_i$. If we consider the completion \widehat{P}_J with respect to J , this decomposes as

$$\widehat{P}_J = \prod_i \widehat{P}_i.$$

Applying [Lemma 91.12.5](#) in the Stacks Project, the functor

$$\text{Def}_P \rightarrow \text{Def}_{\widehat{P}_J} \simeq \prod_i \text{Def}_{\widehat{P}_i} = \prod_i \text{Def}_{\widehat{O}_{X, p_i}}$$

is smooth and an isomorphism on tangent spaces. The henselization step is similar. \square

Example 5.1.3. Let $X = \text{Spec } k[x, y]/(xy)$. Then $\widehat{O}_{X, 0} = k[[x, y]]/(xy)$. We will show that all deformations look like $k[x, y]/(xy - t)$ in some sense. Really, we will show that Def_X has a hull, which is given by $k[[t]]$ with universal deformation

$$X_t := \text{Spec } k[[t]][x, y]/(xy - t) \rightarrow \text{Spec } k[[t]].$$

We want T such that $\text{Hom}(T, -) \rightarrow \text{Def}_X$ is smooth and induces an isomorphism on tangent spaces. Given a morphism

$$f: k[[t]] \rightarrow A \ni f(t),$$

we obtain a deformation

$$A[[x, y]]/(xy - f(t))$$

of $k[[x, y]]/(xy)$. We can check that this is smooth, and so we need to check that we have an isomorphism of tangent spaces.

We want to compute $\text{Def}_X(k[\varepsilon])$. If $S \in \text{Def}_X(k[\varepsilon])$, set $R = k[[x, y]]/(x, y)$, and we have a diagram

$$\begin{array}{ccc} k[\varepsilon] & \longrightarrow & k \\ \downarrow & & \downarrow \\ S & \longrightarrow & R. \end{array}$$

If we consider the exact sequence

$$0 \rightarrow k \rightarrow k[\varepsilon] \rightarrow k \rightarrow 0$$

and tensor with S , by flatness we have an exact sequence

$$0 \rightarrow k \otimes_{k[\varepsilon]} S \rightarrow S \rightarrow k \otimes_{k[\varepsilon]} S \rightarrow 0.$$

But because $k \otimes_{k[\varepsilon]} S = R$, we have a decomposition $S = R \oplus \varepsilon R$. If we choose lifts \tilde{x}, \tilde{y} of x, y , the product $\tilde{x}\tilde{y}$ lifts $xy = 0$, so $\tilde{x}\tilde{y} = (f(x) + g(y))\varepsilon$. If we choose different lifts $\tilde{x}' = \tilde{x} + \varepsilon h, \tilde{y}' = \tilde{y} + \varepsilon h$, we see that

$$\tilde{x}'\tilde{y}' = \tilde{x}\tilde{y} + (xj + yh)\varepsilon.$$

But now $xj + yh \in (x, y)$, and so if we set $t = f(0) + g(0)$, then we can write $\tilde{x}\tilde{y} \in t\varepsilon + (x, y)\varepsilon$, and so there is a canonical choice of the element of $(x, y)\varepsilon$, namely 0. Thus we can set

$$S_t = k[\varepsilon][[\tilde{x}, \tilde{y}]]/(\tilde{x}\tilde{y} - t\varepsilon),$$

and now we have a diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & R & \longrightarrow & S_t & \longrightarrow & R & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & R & \longrightarrow & S & \longrightarrow & R & \longrightarrow & 0. \end{array}$$

By the five lemma, $S_t \simeq S$, and so we are done.

5.2 Smooth curves with a finite group action

In this section, we will consider lifting from characteristic $p > 0$ to characteristic 0. This means we will have to redefine what lifting. Here, we will suppose $k = \bar{k}$ with characteristic p . Let Λ be a Noetherian complete local ring with residue field k . We define \mathcal{C}_Λ be the category of local Artinian Λ -algebras with residue field k .²

Forgetting about the requirement that k is algebraically closed, if $k = \mathbb{F}_p$, then we can set $\Lambda = \mathbb{Z}_p$. Let $W = W(k)$ be the ring of Witt vectors in k . This is apparently some universal Noetherian local ring lifting k , and for example, $W(\mathbb{F}_p) = \mathbb{Z}_p$, and $W(\bar{\mathbb{F}}_p) = \mathcal{O}_{\bar{\mathbb{Q}}_p}$. We will consider some extension \mathcal{O}/W of rings and write K for the fraction field of \mathcal{O} .

We are interested in deforming smooth proper curves X/k with the action of some finite group G , conveniently denoted as a pair (X, G) .

Definition 5.2.1. The pair (X, G) *lifts to characteristic 0* if there exists \mathcal{O} and \mathcal{X}/\mathcal{O} a projective smooth curve with an action of G such that $(\mathcal{X} \times_{\mathcal{O}} k, G) \simeq (X, G)$.

Given the action of G on X , we can construct the quotient X/G . To remember the data of the action, we remember the ramified Galois cover $\pi: X \rightarrow X/G$.³

Consider the action of $\mathrm{PGL}_2(\mathbb{F}_q)$ on \mathbb{P}_k^1 . This includes in $\mathbb{P}_{\mathcal{O}}^1$, which has an action of $\mathrm{PGL}_2(\mathcal{O}) \subset \mathrm{PGL}_2(K)$. We want an embedding $\mathrm{PGL}_2(\mathbb{F}_q) \subseteq \mathrm{PGL}_2(K)$, but generally this is not possible. For example, set $q = 9$ and consider the matrices

$$\begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \quad \begin{pmatrix} a+1 & a \\ -a & 1-a \end{pmatrix},$$

which generate a $(\mathbb{Z}/3)^3$. But then all finite subgroups of $\mathrm{PGL}_2(K)$ are cyclic, dihedral, A_4 , A_5 , or S_4 , so this action clearly cannot lift.

Proposition 5.2.2. *Suppose (X, G) is such that for all $x \in X$, the stabilizer G_x has order $|G_x|$ prime to p . Then (X, G) lifts to characteristic 0.*

Note that this is some local statement, so we need some kind of local-to-global method. Recall that $\widehat{\mathcal{O}}_{X,x} = k[[t]]$, and this has an action of G_x . Thus we define a local action as some action of a finite group G on $k[[t]]$, and this lifts if there exists \mathcal{O} with an action of G on $\mathcal{O}[[t]]$ specializing to the original action on $k[[t]]$.

Theorem 5.2.3. *For (X, G) , suppose that the local action of G_x on $\widehat{\mathcal{O}}_{X,x}$ lifts to characteristic 0. Then (X, G) lifts to characteristic 0.⁴*

²Apparently this is interesting in infinite-combinatorics number theory.

³This is useful if you want to look at some of the omitted proofs.

⁴Avi was lured in to a paper with French title and French body but English abstract, and in the end did not read this paper.

Caleb and Morena (Oct 29): [Lemma 0E3X](#) and applications to contracting curves

6.1 One lemma in the Stacks Project

Lemma 6.1.1. In [Example 0DY7](#) let $f: X \rightarrow Y$ be a morphism of schemes over k . If $f_*\mathcal{O}_X = \mathcal{O}_Y$ and $R^1f_*\mathcal{O}_X = 0$, then the morphism of deformation categories

$$\mathrm{Def}_{X \rightarrow Y} \rightarrow \mathrm{Def}_X$$

is an equivalence.

Let A be an Artinian local ring with residue field k . Remember that $\mathrm{Def}_X(A)$ is the set of isomorphism classes of diagrams

$$(6.1) \quad \begin{array}{ccc} X & \longrightarrow & X_A \\ \downarrow & & \downarrow \alpha \\ \mathrm{Spec} k & \longrightarrow & \mathrm{Spec} A \end{array}$$

with α flat. Also, $\mathrm{Def}_{X \rightarrow Y}$ consists of diagrams of the form

$$\begin{array}{ccccc} & & Y & \longrightarrow & Y_A \\ & \nearrow f & & & \nearrow f_A \\ X & \longrightarrow & X_A & & \\ \downarrow & & \downarrow \alpha & & \downarrow \beta \\ \mathrm{Spec} k & \longrightarrow & \mathrm{Spec} A & & \end{array}$$

with α, β flat.

Lemma 6.1.2 ([Lemma 063Y](#)). Let $(f, f'): (X, X') \rightarrow (S, S')$ be a morphism of first order thickenings such that f is flat. Then the following are equivalent:

1. f' is flat and $X = S \times_{S'} X'$;
2. The canonical map $f^*C_{S/S'} \rightarrow C_{X/X'}$ is an isomorphism, where C is the conormal sheaf.

Proof. In the affine case, write $X = \text{Spec } B, X' = \text{Spec } B', S = \text{Spec } A, S' = \text{Spec } A'$. Then we are looking for a diagram of the form

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \longrightarrow & A' & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow & & \downarrow f' & & \downarrow f & & \\ 0 & \longrightarrow & J & \longrightarrow & B' & \longrightarrow & B & \longrightarrow & 0. \end{array}$$

The two conditions become

1. f is flat and $B = B' \otimes_{A'} A$;
2. $I/I^2 \otimes_A B = J/J^2$ and $I \otimes_A B = J$.

To begin, note that $B = B' \otimes_{A'} A$ is equivalent to $B'/J = B' \otimes_{A'} A'/I$, and this implies that $J = IB'$ and thus that $I \otimes_A B' \rightarrow J$ is surjective. To prove injectivity, by flatness of B' , the map $0 \rightarrow I \rightarrow A'$ remains injective after tensoring with B' , so $I \otimes_{A'} B' \rightarrow B'$ is injective.

In the other direction, we may cite [Lemma 051C](#). Alternatively, we give the following argument. Assuming that $I \otimes_A B \rightarrow J$ is an isomorphism, we know $J = IB'$, and thus $B = B' \otimes_{A'} A$. To prove that B' is flat over A' , we know that B'/IB' is flat over A because B/A is flat and $J = IB'$. We will prove that if $\mathfrak{a} \subset A'$ is an ideal, then $\mathfrak{a} \otimes_{A'} B' \rightarrow B'$ is injective.

By some inexplicable brilliancy, we simply need to fill in the diagram

$$\begin{array}{ccccccccc} ? & \longrightarrow & \mathfrak{a} \otimes_{A'} B' & \longrightarrow & ? & \longrightarrow & 0 & & \\ & & \downarrow & & \downarrow ? & & \downarrow & & \\ 0 & \longrightarrow & IB' & \longrightarrow & B' & \longrightarrow & B'/IB' & \longrightarrow & 0. \end{array}$$

By diagram chasing reasons, we will have exactness. Consider the exact sequence

$$0 \rightarrow I \cap \mathfrak{a} \rightarrow \mathfrak{a} \rightarrow (I + \mathfrak{a})/I \rightarrow 0.$$

After tensoring with B' , we obtain a right exact sequence

$$(I \cap \mathfrak{a}) \otimes_{A'} B' \rightarrow \mathfrak{a} \otimes_{A'} B' \rightarrow (I + \mathfrak{a})/I \otimes_{A'} B' \rightarrow 0.$$

This gives us the desired items in the question marks.

Now we want to prove that $(I \cap \mathfrak{a}) \otimes_{A'} B' \rightarrow IB'$ is injective. If we consider $0 \rightarrow I \cap \mathfrak{a} \rightarrow I$ and tensor with B'/IB' , we obtain

$$0 \rightarrow (I \cap \mathfrak{a}) \otimes_A B'/IB' \rightarrow I \otimes_A B'/IB',$$

but this is clearly actually

$$(I \cap \mathfrak{a}) \otimes_{A'} B' \hookrightarrow I \otimes_A B'.$$

For the other part, we simply take

$$0 \rightarrow (I + \mathfrak{a})/I \rightarrow A$$

and tensor with B'/IB' . □

Proof of Lemma 0E3X. We need to prove that $\beta: (Y, f_*\mathcal{O}_{X_A}) \rightarrow \text{Spec } A$ is flat. We can compose any thickening as a sequence

$$\begin{array}{ccccccc} X & \longrightarrow & X_2 & \longrightarrow & \cdots & \longrightarrow & X_{n-1} & \longrightarrow & X_A \\ \downarrow \alpha_1 & & \downarrow \alpha_2 & & & & \downarrow \alpha_{n-1} & & \downarrow \alpha_n \\ \text{Spec } k & \longrightarrow & \text{Spec } A/\mathfrak{m}_A^2 & \longrightarrow & \cdots & \longrightarrow & \text{Spec } \mathfrak{m}_A^{n-1} & \longrightarrow & \text{Spec } A. \end{array}$$

Now we apply Lemma 063Y to each square and obtain

$$\mathcal{O}_X \otimes_k \mathfrak{m}_A^i/\mathfrak{m}_A^{i+1} = \alpha_i^*(\mathfrak{m}_A^i/\mathfrak{m}_A^{i+1}) = \mathfrak{m}_A^i \mathcal{O}_{X_A}/\mathfrak{m}_A^{i+1} \mathcal{O}_{X_A}.$$

Now if we consider the exact sequence

$$0 \rightarrow \mathfrak{m}_A^i \mathcal{O}_{X_A}/\mathfrak{m}_A^{i+1} \mathcal{O}_{X_A} \rightarrow \mathcal{O}_{X_A}/\mathfrak{m}_A^{i+1} \mathcal{O}_{X_A} \rightarrow \mathcal{O}_{X_A}/\mathfrak{m}_A^i \mathcal{O}_{X_A} \rightarrow 0.$$

Applying f_* and the assumption that $\mathbf{R}^1 f_* \mathcal{O}_X = 0$, we obtain an exact sequence

$$0 \rightarrow f_* \mathcal{O}_X \otimes_k \mathfrak{m}_A^i/\mathfrak{m}_A^{i+1} \rightarrow f_*(\mathcal{O}_{X_A}/\mathfrak{m}_A^{i+1} \mathcal{O}_{X_A}) \rightarrow f_*(\mathcal{O}_{X_A}/\mathfrak{m}_A^i \mathcal{O}_{X_A}) \rightarrow 0.$$

Now if we consider the diagram

$$\begin{array}{ccccccc} Y & \longrightarrow & Y_2 & \longrightarrow & \cdots & \longrightarrow & Y_{n-1} & \longrightarrow & Y_A \\ \downarrow \beta & & \downarrow \beta & & & & \downarrow \beta_{n-1} & & \downarrow \beta \\ \text{Spec } k & \longrightarrow & \text{Spec } A/\mathfrak{m}_A^2 & \longrightarrow & \cdots & \longrightarrow & \text{Spec } \mathfrak{m}_A^{n-1} & \longrightarrow & \text{Spec } A, \end{array}$$

we want to prove that β is flat starting with β_1 being flat. But here, we know

$$\mathcal{O}_{Y_A} = f_* \mathcal{O}_{X_A},$$

and therefore we have

$$\begin{aligned} \beta_i^*(\mathfrak{m}_A^i/\mathfrak{m}_A^{i+1}) &= \mathcal{O}_{Y_i} \otimes_{A/\mathfrak{m}_A^i} \mathfrak{m}_A^i/\mathfrak{m}_A^{i+1} \\ &= \mathcal{O}_Y \otimes_k \mathfrak{m}_A^i/\mathfrak{m}_A^{i+1} \\ &= \mathfrak{m}_A \mathcal{O}_{Y_A}/\mathfrak{m}_A^{i+1} \mathcal{O}_{Y_A}. \end{aligned}$$

Now we may apply Lemma 063Y repeatedly to obtain flatness of β . □

6.2 Application to moduli of curves

In this part of the lecture, we wish to define a contraction map $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ that deletes a marked point. This was used by Knudsen to prove that $\overline{\mathcal{M}}_{g,n}$ is a smooth and proper stack over $\text{Spec } \mathbb{Z}$. Other applications include relations between $\overline{\mathcal{M}}_G$ and $\overline{\mathcal{M}}_{g+1}$ and their Chow groups. The roadmap for this section is:

1. We will discuss stable curves over an algebraically closed field k and define $\overline{\mathcal{M}}_{g,n}$.
2. We will define $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ in the wrong way.
3. We will define contraction of rational tails and bridges over an algebraically closed field correctly.
4. Finally, we will use Lemma 0E3X to define contraction over any scheme.

6.2.1 Stable curves We will define n -marked genus g stable curves C over $k = \bar{k}$. Here, we will take C to be a connected 1-dimensional scheme of finite type over k . We will work only with nodal curves, which are curves where for every $x \in C(k)$ either x is smooth and $\mathcal{O}_{C,x}$ is a regular local ring and thus a UFD by Auslander-Buchsbaum or x is a node and

$$\widehat{\mathcal{O}}_{C,x} \simeq k[[x,y]]/(xy).$$

Remark 6.2.1. In fact, $\widehat{\mathcal{O}}_{C,x}$ is reduced and thus $\mathcal{O}_{C,x}$ is reduced, so every nodal curve is reduced.

Now let

$$\bigcup_i \tilde{C}_i = \tilde{C} \xrightarrow{\nu} C = \bigcup_i C_i$$

be the normalization. We know that $\nu^{-1}(\text{node}) = \text{Spec } k \sqcup \text{Spec } k$. To check this, recall that the normalization of a reduced scheme is constructed by gluing the local maps

$$\text{Spec}(\overline{A_{\text{red}}})^{Q(A_{\text{red}})} \rightarrow \text{Spec } A_{\text{red}}.$$

But now we know that

$$Q(A_{\text{red}}) = \prod_{\mathfrak{p}_i \text{ minimal}} Q(A_{\text{red}}/\mathfrak{p}_i).$$

However, we know that

$$\overline{k[[x,y]]/(xy)}^{Q(k[[x,y]]/(xy))=k((x)) \times k((y))} = k[[x]] \times k[[y]].$$

This implies that $\overline{\mathcal{O}_{C,p}}^{Q(\mathcal{O}_{C,p})}$ is not local because $k[[x]] \times k[[y]]$ is not local.

Proposition 6.2.2. *The arithmetic genus is given by*

$$g = \sum_i g(\tilde{C}_i) + \#(\text{nodes}) - \#(\text{components}) + 1.$$

Proof. Recall that $\nu^{-1}(\text{node}) = \text{Spec } k \sqcup \text{Spec } k$. Then we have the exact sequence

$$0 \rightarrow \mathcal{O}_C \hookrightarrow \nu_* \mathcal{O}_{\tilde{C}} \rightarrow \bigoplus_{\mathfrak{p} \text{ node}} K_{\mathfrak{p}} \rightarrow 0.$$

Taking the long exact sequence in cohomology, we obtain

$$0 \rightarrow H^0(C, \mathcal{O}_C) \rightarrow H^0(C, \nu_* \mathcal{O}_{\tilde{C}}) \rightarrow k^{\#(\text{nodes})} \rightarrow H^1(C, \mathcal{O}_C) \rightarrow H^1(C, \nu_* \mathcal{O}_{\tilde{C}}).$$

Because ν is finite, the Leray spectral sequence computing $H^1(\tilde{C}, \mathcal{O}_{\tilde{C}})$ degenerates at the E_2 -page, and thus

$$H^p(C, \nu_* \mathcal{O}_{\tilde{C}}) = H^p(\tilde{C}, \mathcal{O}_{\tilde{C}}).$$

But then we know $H^1(C, \nu_* \mathcal{O}_{\tilde{C}}) = \bigoplus H^1(\mathcal{O}_{\tilde{C}_i}) = \sum g(\tilde{C}_i)$. By connectedness of C , we know that $H^0(C, \mathcal{O}_C) = k$ and $H^0(C, \nu_* \mathcal{O}_{\tilde{C}}) = k^{\#(\text{components})}$. \square

We now need to add marked points.

Definition 6.2.3. A n -marked, genus g nodal curve C over k is a nodal curve of genus g with n smooth points $x_1, \dots, x_n \in C(k)$.

This definition allows too many curves, so we want to define a notion of stability.

Definition 6.2.4. A genus g nodal curve C with n marked points is *stable* if for all irreducible components \tilde{C}_i , we have

$$2g - 2 + \#(\text{special points}) > 0,$$

where a special point is a node or a marked point.

Example 6.2.5. Over C , consider a genus 2 curve with two marked points x, y , an elliptic curve with marked point z , and a nodal cubic with no marked points all intersecting in a triangle. Here, we have three marked points and four nodes.¹ By observation, this curve is stable.

Remark 6.2.6. All of the \tilde{C}_i are stable if and only if $g(\tilde{C}_i) \geq 2$, $g(\tilde{C}_i) = 1$ and \tilde{C}_i has at least one special point, or if $g(\tilde{C}_i) = 0$ and there are at least three special points.

6.2.2 Contraction done wrong We will denote by $\overline{\mathcal{M}}_{g,n}$ be the fibered category whose objects over a scheme S are given by the following data:

- A map $f: \mathcal{C} \rightarrow S$ proper and flat with sections $\sigma_1, \dots, \sigma_n: S \rightarrow \mathcal{C}$ and for any geometric point $s \in S$, the map $f_s: \mathcal{C}_s \rightarrow k(s)$ and sections $\sigma_{1,s}, \dots, \sigma_{n,s}$ form a stable curve of genus g .
- Morphisms are given by cartesian diagrams

$$\begin{array}{ccc} \mathcal{C}' & \longrightarrow & \mathcal{C} \\ \sigma'_i \uparrow \downarrow & & \downarrow \uparrow \sigma_i \\ S' & \longrightarrow & S \end{array}$$

that respect the sections.

Clearly the forgetful functor simply forgets the last section. Unfortunately, this does not actually respect the stability condition. The problem is when the component containing $\sigma_{n+1}(s)$ satisfies $2g_i - 2 + \#(\text{special}) = 1$, because when we delete σ_{n+1} we lose stability. This only happens when $g(\tilde{C}_i) = 1$ and there is exactly one marked point (which is actually smooth and integral, so is an elliptic curve) or when $g(\tilde{C}_i) = 0$ and there are exactly three special points. In the second case, we may have \mathbb{P}^1 with three marked points, two marked points and a node (rational tail), or one node and two marked points (rational bridge). Also, we may have a nodal cubic with one marked point.

6.2.3 Contraction done right In this part, we construct from a prestable curve $\mathcal{C}_{k(s)}$ a stable curve $\tilde{\mathcal{C}}_{k(s)}$. In the rational tail case, we are contracting the entire \mathbb{P}^1 to the node and in the rational bridge case we identify the two nodes and collapse the \mathbb{P}^1 to that point, leaving the two other components intersecting at a node (locally; globally this can be either one or two components). Of course, we need to check that $f_* \mathcal{O}_{\mathcal{C}_{k(s)}} = \mathcal{O}_{\tilde{\mathcal{C}}_{k(x)}}$ and $R^1 f_* \mathcal{O}_{\mathcal{C}_{k(s)}} = 0$. Next, we need to check that contraction can be extended to a neighborhood in a canonical way when $S = \text{Spec } \mathcal{O}_{S,s}^h$.

¹Anna commented that Morena's drawing had the curves tangent to each other, but we can just pretend that they are nodes.

Morena (Nov 05): Contraction morphisms between moduli stacks of curves

7.1 Recap of last time

Recall that for $f: X \rightarrow Y$, if $f_*\mathcal{O}_X = \mathcal{O}_Y$ and $R^1f_*\mathcal{O}_X = 0$, then we have an equivalence of categories

$$\mathrm{Def}_{X \rightarrow Y} \simeq \mathrm{Def}_X.$$

We also attempted to construct a morphism $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ using the forgetful functor, but this does not work. Remember the bad cases were called the rational tail and rational bridge. Also recall that the stability condition for every component was $2g - 2 + \#(\text{special points}) > 0$, where special points are nodes and marked points.

7.2 Contraction of rational tails and bridges

In the rational tail case, our stable curve should be the scheme-theoretic closure (so just topological closure) $\overline{C/C_i}$, where C_i is the rational tail. Because C is reduced, we see that C is the pushout

$$\begin{array}{ccc} \mathrm{Spec} k & \hookrightarrow & C_i \\ \downarrow & & \downarrow \\ \overline{C \setminus C_i} & \longrightarrow & C. \end{array}$$

Of course, this gives us a map $C \rightarrow \overline{C \setminus C_i}$ by contracting C_i . Now we need to check that $\overline{C \setminus C_i}$ is a prestable curve of genus g which is actually stable and that if $c: C \rightarrow \overline{C \setminus C_i}$, then $c_*\mathcal{O}_C \simeq \mathcal{O}_{\overline{C \setminus C_i}}$ and $R^1c_*\mathcal{O}_C = 0$.

To check the sheafy conditions, we have an exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow j^*\mathcal{O}_{\overline{C \setminus C_i}} \oplus j_{i*}\mathcal{O}_{C_i} \rightarrow i_{x*}k \rightarrow 0.$$

This gives us a longer exact sequence

$$0 \rightarrow c_*\mathcal{O}_C \rightarrow c_*j_*\mathcal{O}_{\overline{C \setminus C_i}} \oplus c_*j_{i*}\mathcal{O}_{C_i} \rightarrow c_*i_{x*}k \rightarrow R^1c_*\mathcal{O}_C \rightarrow R^1c_*j_*\mathcal{O}_{\overline{C \setminus C_i}} \oplus R^1c_*j_{i*}\mathcal{O}_{C_i}.$$

To see that $R^1 c_* \mathcal{O}_C = 0$, we only need to check that $R^1 c_* j_* \mathcal{O}_{\overline{C \setminus C_i}} = 0$ because $R^1 c_* j_{i*} \mathcal{O}_{C_i} = H^1(C_i, \mathcal{O}_{C_i}) = 0$. But here we have $c_* j_* \mathcal{O}_{\overline{C \setminus C_i}} = 0$ because $c_* j_* = 0$, and so we need to prove surjectivity of the direct sum onto the skyscraper sheaf. This is clear.

Considering

$$\begin{array}{ccc} \text{Spec } k & \longrightarrow & \overline{C \setminus C_i} \\ \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \overline{C \setminus C_i}, \end{array}$$

we see that $c_* \mathcal{O}_C \simeq \mathcal{O}_{\tilde{C}}$ and

$$\mathcal{O}_{\overline{C \setminus C_i}} = c_* j_* \mathcal{O}_{\overline{C \setminus C_i}} \times_{i_{x*} k} c_* j_{i*} \mathcal{O}_{C_i}.$$

To check stability of $\overline{C \setminus C_i}$, we simply note that stability condition of the component C_j attached to C_i is unchanged, and everything else was untouched, so we have a stable curve.

We now check the rational bridge case. Here we consider the pushout

$$\begin{array}{ccc} \{x_1\} \sqcup \{x_2\} & \longrightarrow & \mathcal{E} \setminus \overline{C_i} \\ \downarrow & & \downarrow \\ \{y\} & \longrightarrow & \overline{C}. \end{array}$$

This exists and has nice properties. What this really means is that we contract C_i and introduce a new self-intersection at $x_1 = x_2$. Now the contraction morphism is given by considering the total pushout of

$$\begin{array}{ccc} \{x_1\} \sqcup \{x_2\} & \longrightarrow & \mathcal{E} \setminus \overline{C_i} \\ \downarrow & & \downarrow \\ C_i & \longrightarrow & C \\ \downarrow & & \downarrow \text{---} \\ \{y\} & \longrightarrow & \overline{C}. \end{array}$$

Restricting to an affine piece, where $\overline{C} = \text{Spec } A$ and $C \setminus \overline{C_i} = \text{Spec } A'$, we obtain a fiber product diagram of rings

$$\begin{array}{ccc} A & \longrightarrow & k \\ \downarrow & & \downarrow \\ A' & \longrightarrow & k \times k. \end{array}$$

We can now tensor with A_y , and we need to show that that $\widehat{A}_y = k[[x, y]]/(xy)$. Because completions are exact, we can consider the fiber product diagram

$$\begin{array}{ccc} \widehat{A}_y & \longrightarrow & k \\ \downarrow & & \downarrow \\ A' \otimes_A \widehat{A}_y & \longrightarrow & k \times k. \end{array}$$

Because $A' \otimes_{\Lambda} \widehat{A}_y = \widehat{A}'_{x_1} \times \widehat{A}'_{x_2} = k[[t_1]] \times k[[t_2]]$, we see that

$$\widehat{A}_y = \{\ell + t_1 p(t_1) + t_2 p(t_2)\} = k[[t_1, t_2]] / (t_1 t_2),$$

and thus y is a node.

7.3 Contraction over any base scheme

Theorem 7.3.1. *Let $(S, f: \mathcal{C} \rightarrow S, \sigma_1, \dots, \sigma_{n+1}) \in \overline{\mathcal{M}}_{g, n+1}(S)$. Then there exists a contraction such that*

$$f = \mathcal{C} \xrightarrow{c} \overline{\mathcal{C}} \xrightarrow{g} S$$

and the following conditions hold:

1. $(S, g: \overline{\mathcal{C}} \rightarrow S, c \circ \sigma_1, \dots, c \circ \sigma_n) \in \overline{\mathcal{M}}_{g, n}(S)$.
2. $c_* \mathcal{O}_{\mathcal{C}} \simeq \mathcal{O}_{\overline{\mathcal{C}}}$ and $R^1 c_* \mathcal{O}_{\mathcal{C}} = 0$, and this is stable under base change. In addition, for all geometric points $\overline{s} \rightarrow S$, $c_{\overline{s}}$ is either an isomorphism or a contraction of a rational tail or rational bridge.

Moreover, $c: \mathcal{C} \rightarrow \overline{\mathcal{C}}$ is unique up to unique isomorphism.

Corollary 7.3.2. *The morphism $\overline{\mathcal{M}}_{g, n+1} \rightarrow \overline{\mathcal{M}}_{g, n}$ is defined over \mathbb{Z} .*

Proof. We know what happens to objects. We know that morphisms are cartesian diagrams

$$\begin{array}{ccc} S' & \longrightarrow & S \\ \downarrow \sigma'_i & & \downarrow \sigma_i \\ \mathcal{C}' & \xrightarrow{b} & \mathcal{C} \\ \downarrow & & \downarrow \\ S' & \xrightarrow{a} & S. \end{array}$$

But now we have two candidate contractions of \mathcal{C}' , namely $\overline{\mathcal{C}'}$ and $\overline{\mathcal{C}} \times_S S'$. But these have a unique isomorphism. These fit into the diagram

$$\begin{array}{ccccc} & \mathcal{C} \times_S S' & \longrightarrow & \mathcal{C} & \\ & \swarrow & & \searrow & \\ \overline{\mathcal{C}'} & \xrightarrow{\quad} & \overline{\mathcal{C}} \times_S S' & \longrightarrow & \overline{\mathcal{C}} \\ & \swarrow & & \searrow & \\ & S' & \longrightarrow & S & \end{array}$$

This gives us compatibility with morphisms. □

To prove the theorem, we want to work étale locally over S , with a cover $(S_i \rightarrow S)$. Then there exists a factorization $\mathcal{C}_i \xrightarrow{a} \overline{\mathcal{C}}_i \rightarrow S_i$, and then we need to check that the cocycle condition works on the overlaps. The existence of a global factorization means that our data is effective, and this is the same as $\overline{\mathcal{M}}_{g, n}$ is a stack.

To prove existence of a factorization étale locally, over $\overline{s} \in S_i$, we have a contraction $c_s: \mathcal{C}_s \rightarrow \overline{\mathcal{C}}_s$. We want to prove that there exists an étale neighborhood $(U, u) \rightarrow (S, s)$ such that there exists a factorization $\mathcal{C}_U \rightarrow \overline{\mathcal{C}}_U$ extending c_s . In order to do this, we need to do some reduction. If $S = \text{Spec } A$ is affine is of finite type over \mathbb{Z} , we want $A_i \subseteq A$ such that $A = \varinjlim A_i$, and thus $S = \text{Spec } A = \varprojlim \text{Spec } A_i$.

Theorem 7.3.3 (0E6U, 0E6V, 0DSS, 0CMV). *The fibered category $\text{Curves}_g^{\text{prestable}}$ is limit-preserving.*

Lemma 7.3.4. *The category of schemes of finite presentation over S is the colimit of the categories of schemes of finite presentation over S_i .*

Now if $S = \text{Spec } A$, then $\mathcal{O}_{S,s}^h = \text{colim}_{(U,u) \rightarrow (S,s)} \mathcal{O}_U(U)$. Thus we may consider $S = \text{Spec}(\mathcal{O}_{S,s}^h)$. Write Λ for this henselization. Then if we consider $\Lambda \rightarrow \Lambda/\mathfrak{m}^n\Lambda$, we have fiber product diagrams

$$\begin{array}{ccccc} C_s & \longrightarrow & C_u & \longrightarrow & C \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \text{Spec } \Lambda/\mathfrak{m}^n\Lambda & \longrightarrow & S. \end{array}$$

This gives us a formal object of $\text{Def}_{C_s}(\widehat{\Lambda})$. But now we know that

$$\text{Def}_{C_s \rightarrow \overline{C}_s} \simeq \text{Def}_{C_s},$$

and thus we actually have deformations of $C_s \rightarrow \overline{C}_s$ over $\Lambda/\mathfrak{m}^n\Lambda$. But this gives us a formal element of

$$\text{Def}_{C_s \rightarrow \overline{C}_s}(\widehat{\Lambda}).$$

Now using 01W0, we have $\overline{C}_n \in \text{Curve}_g$.

Theorem 7.3.5. *The data Curve_g is effective.*

For a sketch of this, note that the first stability condition is an open condition. If p_1, \dots, p_n are the marked points, then write $D = p_1 + \dots + p_n$. Stability is equivalent to $\omega_C(D)$ being ample, and so over C_s we have an ample line bundle $\omega_{C_s}(D)$. We can lift this to $(\overline{C}_n, \mathcal{L}_n)$ because obstructions live in

$$H^2(\overline{C}_{n-1}, (1+m)^n \mathcal{O}_{\overline{C}_n^*}) = 0.$$

Using the Grothendieck algebraization theorem, we see that if $X_i \rightarrow S_i$ is proper and \mathcal{L}_i is ample, there exists a proper morphism $X \rightarrow S$ and \mathcal{L} an ample line bundle such that base change to S_n recovers (X_n, \mathcal{L}_n) .

Here, we know that $\overline{C} \rightarrow \text{Spec } \widehat{\Lambda}$ is finite type and separated while $\mathcal{C} \rightarrow \text{Spec } \widehat{\Lambda}$ is proper, and so there exists a unique $\mathcal{C} \rightarrow \overline{C}$. We now need to return back to $\text{Spec } \Lambda$. But now we know that $\widehat{\Lambda}$ is the direct limit of its finitely-generated (over Λ) subalgebras. This gives us $\Lambda \subseteq \Lambda^1 \subseteq \widehat{\Lambda}$. We know that

$$\Lambda = \mathcal{O}_{S,s}^h \rightarrow \widehat{\mathcal{O}}_{S,s}$$

is regular, it is flat, and thus $\widehat{\Lambda}/\mathfrak{m}\widehat{\Lambda}$ is noetherian and geometrically regular over $k(\mathfrak{m})$ by Popescu. We have the diagram

$$\begin{array}{ccc} \Lambda_{\text{sm}} & \longrightarrow & \Lambda_{\text{ét}} \\ \uparrow u & \searrow & \downarrow \\ \Lambda & \longrightarrow & \Lambda/\mathfrak{m}\Lambda. \end{array}$$

Because Λ is henselian, we have a section $\Lambda_{\text{ét}} \rightarrow \Lambda$, and thus we have a base change to Λ .

Baiqing (Nov 12): Deformations of group schemes

8.1 Deformations of abelian schemes

Definition 8.1.1. An *abelian scheme* $p: X \rightarrow S$ is a proper smooth group scheme over a connected scheme S with geometrically connected fibers.

If $S = \text{Spec } k$ is a field, then X is projective. However, this is not true in general.

Definition 8.1.2. The deformation functor $\text{Def}_X^{\text{AS}}: \Lambda_W \rightarrow \text{Set}$ is defined for an abelian variety X over k and Λ_W the category of local Artinian W -algebras with residue field k . It is given by

$$A \mapsto \{(X', \varphi) \mid X' \rightarrow \text{Spec } A \text{ abelian scheme, } \varphi: X'_k \simeq X\}/$$

where $(X'_1, \varphi'_1) \sim (X'_2, \varphi'_2)$ if there exists an isomorphism $\psi: X'_1 \rightarrow X'_2$ of abelian schemes such that $\varphi'_2 \circ \psi_k = \varphi'_1$.

Theorem 8.1.3. Let X be an abelian variety of dimension g over k . Then the deformation functor Def_X^{AS} is pro-representable by $W(k)[[t_1, t_2, \dots, t_{g^2}]]$.

Recall Schlessinger's criterion from Johan's lecture.¹ Note that if F is formally smooth and $F(R') \rightarrow F(R)$ is surjective for all $R' \twoheadrightarrow R$, then F is pro-representable by a power series ring $W[[t_1, \dots, t_d]]$, where $d = \dim_k F(k[\varepsilon])$.

We will prove the conditions (H_3) , (H_4) and then prove that the deformation functor is formally smooth. Before this, we will consider the geometry of abelian schemes.

Lemma 8.1.4 (Rigidity lemma). *Given a diagram*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p & \swarrow q \\ & & S, \end{array}$$

suppose that S is connected, p is flat and proper, and $H^0(X_s, \mathcal{O}_{X_s}) \simeq k(s)$ for all $s \in S$. If for some point $s \in S$, $f(X_s)$ is set-theoretically a single point, then there exists a section $\eta: S \rightarrow Y$ such that $f = \eta \circ p$.

Corollary 8.1.5. Let X, Y be abelian schemes over S and $f: X \rightarrow Y$. If $f \circ \varepsilon_X = \varepsilon_Y$, then f is a homomorphism. Here, ε_X refers to the identity section.

¹Baiqing wrote them all down on the board, but I am too lazy to type them yet again.

This result holds in general if we replace Y by an arbitrary group scheme.

Sketch of proof. Consider the diagram

$$\begin{array}{ccc} X \times_S X & \longrightarrow & G_X \\ & \searrow & \swarrow \\ & X & \end{array}$$

Consider the morphisms $X \times_S X \rightarrow G \times_S X$ given by $\psi_1(x_1, x_2) = (f(x_1 x_2), x_2)$ and $\psi_2(x_1, x_2) = (f(x_1), x_2)$. Now we argue that $f(g_1 g_2) = f(g_1) f(g_2)$. \square

Also, it is clear that abelian schemes are commutative because inversion takes the identity to itself and thus is a homomorphism. If $\phi_1, \phi_2: X \rightarrow Y$ are two homomorphisms and $(\phi_1)_s = (\phi_2)_s$ for some $s \in S$, then $\phi_1 = \phi_2$. Next, if $f: X \rightarrow Y$ is a morphism of schemes between abelian schemes, then $f - f(0) = f - f \circ \varepsilon \circ p$ is a homomorphism.

We will now prove (H_4) for Def_X^{AS} . For a small extension $R'' \rightarrow R$, we show that

$$\text{Def}_X^{\text{AS}}(R'' \times_R R') \rightarrow \text{Def}_X^{\text{AS}}(R'') \times_{\text{Def}_X^{\text{AS}}(R)} \text{Def}_X^{\text{AS}}(R')$$

is a bijection. This is equivalent to proving that in the diagram

$$\begin{array}{ccc} \text{Def}_X^{\text{AS}}(R'' \times_R R') & \longrightarrow & \text{Def}_X^{\text{AS}}(R'') \\ \downarrow \pi' & & \downarrow \pi \\ \text{Def}_X^{\text{AS}}(R') & \longrightarrow & \text{Def}_X^{\text{AS}}(R), \end{array}$$

if $(X', \varphi') \mapsto (X, \varphi)$ along the bottom arrow, then $\pi'^{-1}((X', \varphi')) = \pi^{-1}((X, \varphi))$.

Proposition 8.1.6. $\pi^{-1}((X, \varphi))$ is the set of isomorphism classes of (X'', φ''_R) such that X'' is flat over R'' and $X''_R \rightarrow X$ is an isomorphism as schemes over R .

Choose $(X'', \varphi'') \in \pi^{-1}((X, \varphi))$. Then there exists an isomorphism $\phi: X''_R \rightarrow X$ of abelian schemes. Now we will send $(X'', \varphi'') \mapsto (X'', \phi)$. Checking uniqueness is easy with the diagram

$$\begin{array}{ccc} X''_k & \xrightarrow{\phi_k} & X_k \\ & \searrow \varphi'' & \swarrow \varphi \\ & X & \end{array}$$

Now we check that if $(X''_1, \varphi''_1) \sim (X''_2, \varphi''_2)$, then they are sent to the same thing. This was erased from the board by Baiqing before I could process it.

Now we prove injectivity. If $(X''_1, \varphi''_1), (X''_2, \varphi''_2)$ map to equivalent $(X''_1, \phi_1), (X''_2, \phi_2)$, then there exists $\psi: X''_1 \simeq X''_2$ such that $\phi_2 \circ \psi_R = \phi_1$. This implies that $\varphi''_2 \circ \psi_k = \varphi''_1$. Unfortunately, ψ is not an isomorphism of abelian schemes, so we can replace it by $\psi = \psi - \psi(0)$.

Finally, we prove surjectivity, which is the difficult part of the argument.

Proposition 8.1.7. Let $S = \text{Spec } A$, where A is an Artinian local ring, $\mathfrak{m} \subset A$ be the maximal ideal, and $I \subset A$ such that $\mathfrak{m} \cdot I = 0$. Let $\pi: X \rightarrow S$ be proper and smooth, $\varepsilon: S \rightarrow X$ be a section of π , $S_0 = \text{Spec } A/I$, and $X_0 = X \times_S S_0$. Then if X_0 is an abelian scheme with identity $\varepsilon|_{S_0}$, then X is an abelian scheme with identity ε .

Theorem 8.1.8. *Let $X, Y/R$ are smooth schemes. Suppose that $\pi: R' \rightarrow R$ is a small extension. Let $X' \in \text{Def}_X(R')$ and $Y' \in \text{Def}_Y(R')$. For any $f: X \rightarrow Y$ over R , there is a canonically associated class $o(f) \in H^1(X_k, f_k^* T_{Y_k/k}) \otimes_k \ker \pi$. If $o(f) = 0$, then $H^0(X_k, f_k^* T_{Y_k/k}) \otimes_k \ker \pi$.*

Proof of proposition. Given (m, ε, ι) , we want (μ, ε) , where $\mu(x, y) = x - y$. We have μ on X_0 , and so we want to deform this to μ' on X . We know that

$$o(\mu) \in H^1(X_k \times_k X_k, \mu_k^* T_{X_k/k}) \otimes_k \ker \pi,$$

but because X_k is an abelian variety, we have

$$T_{X_k/k} = H^0(X_k, \Omega_{X_k/k})^* \otimes_k \mathcal{O}_{X_k}.$$

We want to show that $o(\mu) = 0$. Consider $g_1 = \Delta: X_0 \rightarrow X_0 \times_{S_0} X_0$ and $g_2 = (\text{id}, \varepsilon p): X_0 \rightarrow X_0 \times_{S_0} X_0$. Then $\mu \circ g_1 = \varepsilon_0 \circ \pi$, and thus $o(\mu \circ g_1) = 0$. Also, $\mu \circ g_2 = 1_{X_0}$ and $o(\mu \circ g_2) = 0$. We know that

$$o(\mu \circ g_1) = (g_1)_k^* o(\mu) = 0, \quad o(\mu \circ g_2) = (g_2)_k^* o(\mu) = 0.$$

Consider the morphisms $(g_1)_k^*: H^1(X_k \times X_k, \mu_k^* T_{X_k/k}) \rightarrow H^1(X_k, (\varepsilon \circ \pi)^* T_{X_k/k})$ and $(g_2)_k^*: H^2(X_k \times X_k, \mu_k^* (T_{X_k/k})) \rightarrow H^1(X_k, (\varepsilon \circ \pi)^* T_{X_k/k})$. But now we see that they are given by $(g_1)_k^*: (x, y) \otimes v \mapsto (x + y) \otimes v$ and $(x, y) \otimes v \mapsto x \otimes v$, and so $o(\mu) = 0$.

Now consider $H^0(X_k \times_k X_k, \mu_k^* T_{X_k/k}) \otimes_k \ker \pi \simeq \mathfrak{t} \otimes_k I$ and let μ' be a deformation of μ . Then

$$S \xrightarrow{(\varepsilon, \varepsilon)} X \times_S X \xrightarrow{\mu'} X$$

is a deformation of ε_0 . Under the identification

$$H^0(\text{Spec } k, (\varepsilon_0)_k^* T_{X_k/k}) \otimes_k \ker \pi \simeq \mathfrak{t} \otimes_k I$$

and using what we have done previously, we have $\mu' \circ (\varepsilon, \varepsilon) = \varepsilon$ and therefore an abelian scheme structure. \square

Finally we prove that $\text{Def}_X^{\text{AS}}(R') \rightarrow \text{Def}_X^{\text{AS}}(R)$ is surjective for $R' \rightarrow R$. Let $(\mathcal{X}, \varphi) \in \text{Def}_X^{\text{AS}}(R)$. Then $o(\mathcal{X}) \in H^2(X, T_{X/k}) \otimes_k \ker \pi = \mathfrak{t} \otimes_k (\mathfrak{t}^* \wedge \mathfrak{t}^*)$. We want to prove that $\pi^{-1}((\mathcal{X}, \varphi))$ is nonempty. We know that ι^* induces -1 on $\mathfrak{t}, \mathfrak{t}^*$, and so $o(\mathcal{X}) = -o(\mathcal{X})$, and if we are working not in characteristic 2, $o(\mathcal{X}) = 0$.

To compute the dimension of the tangent space, we want to compute $\text{Def}_X^{\text{AS}}(k[\varepsilon])$, but this is $H^1(X, T_{X/k}) \simeq \mathfrak{t} \otimes \mathfrak{t}^*$, which has dimension g^2 .

8.2 Deformations of smooth affine group schemes

We will now discuss deformations of G_m and G_a . Let G/k be an affine smooth algebraic group scheme. We will consider the deformation problem

$$\text{Def}_G(R) = \left\{ (G', \phi) \mid G'/R \text{ group scheme, } G_s \xrightarrow{\phi} \sim G \right\} / \text{iso}.$$

For example, for G_m , we know that $\text{Spec } k[\varepsilon][t, t^{-1}]$ is a deformation of G_m . We want to deform the multiplication, and so we have $m': T \rightarrow T_1 T_2 (1 + \varepsilon \Delta(T_1, T_2))$. To check associativity, we obtain

$$\Delta(T_1, T_2) + \Delta(T_1 T_2, T_2) = \Delta(T_1, T_2 T_3) + \Delta(T_2, T_3).$$

We will also consider Δ, Δ' equivalent if there exists f such that $\Delta'(T_1, T_2) = \Delta(T_1, T_2) + f(T_1 T_2) - f(T_1) - f(T_2)$. Therefore we have

$$\text{Def}_{G_m}(k[\varepsilon]) \cong \frac{\{\Delta \mid \text{associative}\}}{\{f(T_1 T_2) - f(T_1) - f(T_2)\}}.$$

Proposition 8.2.1. *In fact, we have $\text{Def}_{G_m}(k[\varepsilon]) = 0$.*

Proof. let $f(T) = \sum a_i T^i$. Then $f(T_1 T_2) - f(T_1) - f(T_2) = \sum a_i (T_1^i T_2^i - T_1^i - T_2^i)$. If we set

$$\Delta(T_1, T_2) = \lambda_{ij} T_1^i T_2^j,$$

adjust Δ by f such that $\lambda_{i0} = 0$. Now we have the equation

$$\sum \lambda_{ij} T_1^i T_2^j = \sum_{i,j} T_1^i T_2^i T_3^j - \left(\sum \lambda_{ij} T_1^i T_2^j T_3^j + \sum \lambda_{ij} T_2^i T_3^j \right),$$

and thus $\lambda_{ij} = 0$. □

For the additive group, our deformed multiplication is given by $T \rightarrow T_1 + T_2 + \varepsilon \Delta(T_1, T_2)$. Associativity is equivalent to the condition

$$\Delta(T_1, T_2) + \Delta(T_1 + T_2, T_3) = \Delta(T_2, T_3) + \Delta(T_1, T_2 + T_3).$$

The trivial deformations are given by $f(T_1 + T_2) - f(T_1) - f(T_2)$, and thus we have

$$\text{Def}_{G_a}(k[\varepsilon]) \simeq \frac{\Delta \mid \text{associative}}{\{f(T_1 + T_2) - f(T_1) - f(T_2)\}}.$$

This deformation space vanishes in characteristic 0 and is infinite-dimensional in positive characteristic. If we apply $\frac{\partial}{\partial T_3}$ to the associativity relation, we obtain

$$\Delta_2(T_1 + T_2, T_3) = \Delta_2(T_2, T_3) + \Delta_2(T_1, T_2 + T_3),$$

and if we apply $\frac{\partial}{\partial T_1}$, we obtain

$$\Delta_{12}(T_1 + T_2, T_3) = \Delta_{12}(T_1, T_2 + T_3),$$

and therefore $\Delta_{12}(T_1, T_2) = f(T_1 + T_2)$. In characteristic 0, f has a primitive, and so taking the primitive twice we obtain a desired \tilde{F} . Primitives do not exist in positive characteristic, so this proof does not work. However, it is possible to work out the following:

Proposition 8.2.2. *In positive characteristic p , Δ is a linear combination of $B_p(T_1, T_2)^{p^n}$ for $n \geq 0$ and $T_1^{p^n} T_2^{p^m}$ for $m \geq n + 1$, where $B_p(T_1, T_2) = \frac{(T_1 + T_2)^p - T_1^p - T_2^p}{p} \in \mathbb{Z}[T_1, T_2]$.*

Haodong (Nov 19): Artin's axioms

9.1 Prestacks and stacks

First, we will generalize the Zariski topology a little bit.

Definition 9.1.1. Let \mathcal{S} be a category. Then a *Grothendieck topology* on \mathcal{S} is given by a set $\text{Cov}(X)$ for each $x \in X$, where each element in $\text{Cov}(X)$ is a collection of morphisms $\{x_i \rightarrow x\}_I$ in \mathcal{S} . We require that

1. All isomorphisms $x' \xrightarrow{\sim} x$ are in $\text{Cov}(x)$.
2. If $\{x_i \rightarrow x\}_I \in \text{Cov}(X)$ and $y \rightarrow x$, then $x_i \times_x y$ exist and $\{x_i \times_x y \rightarrow y\}_I \in \text{Cov}(y)$.
3. If $\{x_i \rightarrow x\}_I$ is a covering of X and $\{x_{ij} \rightarrow x_i\}_{J_i}$ is a covering of x_i for all i , then $\{x_{ij} \rightarrow x\}$ is a covering of X .

A *site* is a category with a Grothendieck topology.

Example 9.1.2. Let S be a scheme and let $\mathcal{C} = \text{Sch}/S$. For every $T \rightarrow S$, we say that $\{f_i: T_i \rightarrow T\}$ is a covering if all f_i are open immersions and the f_i are jointly surjective. This gives the big Zariski site of S . If we replace open immersion with étale morphism, then we obtain the big étale site.

9.1.1 Prestacks

Definition 9.1.3. Let \mathcal{S} be a category and $p: \mathcal{X} \rightarrow \mathcal{S}$ be a functor. We will denote objects of \mathcal{X} by a, b, \dots and objects in \mathcal{S} by S, T, \dots . We will denote morphisms in \mathcal{X}, \mathcal{S} by α, f respectively. We say that p is a *prestack* if the following conditions are satisfied:

1. If $p(b) = T$ and $S \rightarrow T$ is a morphism, there exists a and a morphism $a \rightarrow b$ such that $p(a) = S$ filling the diagram

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \downarrow \\ S & \xrightarrow{f} & T. \end{array}$$

2. If we have

$$\begin{array}{ccccc}
 & & \curvearrowright & & \\
 a & & & b & \longrightarrow & c \\
 \downarrow & & & \downarrow & & \downarrow \\
 R & \longrightarrow & S & \longrightarrow & T,
 \end{array}$$

there exists a unique morphism $a \rightarrow b$ filling in the above arrow.

Exercise 9.1.4. For every $s \in \mathcal{S}$, the category $\mathcal{X}(s)$ over s has morphisms those lying over the identity and is thus a groupoid.

Example 9.1.5. Let $F: \mathcal{S} \rightarrow \text{Set}$ be a functor. We will define \mathcal{X}_F to have objects (a, s) , where $s \in \mathcal{S}$ and $a \in F(s)$. Then morphisms $(a, s) \rightarrow (a', s')$ are morphisms $f: s \rightarrow s'$ such that $F(f)(a) = a'$. We will define $p: (a, s) \mapsto s$, and this defines a prestack, which we will simply call F .

Example 9.1.6. Let S be a scheme and $\mathcal{C} = \text{Sch}/S$. If $T \rightarrow S$ is an S -scheme, then $\text{Hom}(-, T)$ is a presheaf on \mathcal{C} and by the previous example defines a prestack over \mathcal{C} , which we will call T .

Example 9.1.7. Consider the functor $\mathcal{M}_g \rightarrow \text{Sch}/\mathcal{C}$ where the objects are morphisms $\mathcal{C} \rightarrow S$, where S is a scheme over \mathcal{C} and $\mathcal{C} \rightarrow S$ is smooth and proper with all geometric fibers connected curves of genus g . Morphisms are simply pairs $\alpha: \mathcal{C} \rightarrow \mathcal{C}', f: S \rightarrow S'$ such that $\mathcal{C} = \mathcal{C}' \otimes_S S$.

Definitions 9.1.8. Let \mathcal{S} be a site.

1. A *morphism of prestacks* $\mathcal{X} \rightarrow \mathcal{Y}$ is a functor f such that for all $a \in \mathcal{X}$, $p_{\mathcal{X}}(a) = p_{\mathcal{Y}}(f(a))$.
2. If $f, g: \mathcal{X} \rightarrow \mathcal{Y}$ are morphisms of prestacks, then a *2-morphism* $\alpha: f \rightarrow g$ is a natural transformation such that for all $a \in \mathcal{X}$, $\alpha_a: f(a) \rightarrow g(a)$ lies over the identity in \mathcal{S} . In particular, α is an isomorphism.
3. A diagram

$$\begin{array}{ccc}
 \mathcal{X} & \xrightarrow{f'} & \mathcal{Y} \\
 \downarrow g' & & \downarrow g \\
 \mathcal{Y}' & \xrightarrow{f} & \mathcal{Z}
 \end{array}$$

is a *2-commutative diagram* if there exists a 2-morphism $\alpha: fg' \rightarrow gf'$.

4. $f: \mathcal{X} \rightarrow \mathcal{Y}$ is called an *equivalence* if there exists $g: \mathcal{Y} \rightarrow \mathcal{X}$ such that $f \circ g \xrightarrow{\sim} \text{id}$ and $g \circ f \xrightarrow{\sim} \text{id}$.

Lemma 9.1.9 (2-Yoneda lemma). *Let $\mathcal{X} \rightarrow \mathcal{S}$ be a prestack and $s \in \mathcal{S}$. Then $\text{Hom}_{\mathcal{S}}(-, s)$ is a prestack on \mathcal{S} and the functor $\text{Hom}(s, \mathcal{X}) \rightarrow \mathcal{X}(s)$ given by $f \mapsto f_s(\text{id}_s)$ is an equivalence of categories.*

Now we want a fiber product of prestacks for $\mathcal{X} \rightarrow \mathcal{Y}, \mathcal{Y}' \rightarrow \mathcal{Y}$, which is simply the final object in all 2-commutative diagrams

$$\begin{array}{ccc}
 \mathcal{Z} & \longrightarrow & \mathcal{X} \\
 \downarrow & & \downarrow \\
 \mathcal{Y}' & \longrightarrow & \mathcal{Y}
 \end{array}$$

Fortunately, these do exist.

Example 9.1.10. The product $\mathcal{X} \times \mathcal{X}$ exists and is a prestack. We also have a diagonal $\Delta: \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$.

Example 9.1.11. Let \mathcal{X} be a prestack and $a: \mathcal{Y} \rightarrow \mathcal{X}, b: \mathcal{Y}' \rightarrow \mathcal{X}$ be morphisms. Then there exists a 2-cartesian diagram

$$\begin{array}{ccc} \mathcal{Y} \times_{\mathcal{X}} \mathcal{Y}' & \longrightarrow & \mathcal{Y} \times \mathcal{Y}' \\ \downarrow & & \downarrow (a,b) \\ \mathcal{X} & \xrightarrow{\Delta} & \mathcal{X} \times \mathcal{X}. \end{array}$$

9.1.2 Stacks

Definition 9.1.12. A prestack \mathcal{X} over a site \mathcal{S} is called a *stack* if for every $\{U_i \rightarrow U\} \in \text{Cov}(U)$, we have

1. (morphisms glue) There exists a unique $a \rightarrow b$ filling in the diagram

$$\begin{array}{ccccc} & & a|_{U_i} & & \\ & \nearrow & \searrow & \curvearrowright & \\ a|_{U_{ij}} & & & & b \\ & \searrow & \nearrow & \dashrightarrow & \\ & & a|_{U_j} & & \end{array}$$

lying over

$$\begin{array}{ccc} & U_i & \\ & \nearrow & \searrow \\ U_{ij} & & U \\ & \searrow & \nearrow \\ & U_j & \end{array}$$

Precisely, this means that given $a, b \in \mathcal{X}(U)$ and $\phi_i: a|_{U_i} \rightarrow b$ such that $\phi_i|_{U_{ij}} = \phi_j|_{U_{ij}}$, then there exists a unique $\phi: a \rightarrow b$ such that $\phi|_{U_i} = \phi_i$ for all i .

2. (objects glue) Given a_i, a_j and isomorphisms $\alpha_{ij}: a_i|_{U_{ij}} \rightarrow a_j|_{U_{ij}}$ satisfying the cocycle condition on U_{ij} , there exists $a \in \mathcal{X}(U)$ and isomorphisms $\phi_i: a|_{U_i} \rightarrow a_i$ such that $\alpha_{ij} \circ \phi_i|_{U_{ij}} = \phi_j|_{U_{ij}}$.

Example 9.1.13. Consider the prestack Sheaves over Sch with objects (S, F) , where F is a sheaf on S . Then $(S, F) \rightarrow (S', F')$ is a pair $f: S \rightarrow S'$ and $\alpha: F' \rightarrow f_*F$ such that the adjoint of α is an isomorphism $F \simeq f^{-1}F'$. We know that sheaves and their morphisms can be glued in the Zariski topology, so the prestack Sheaves is a stack over Sch_{Zar} .

Example 9.1.14. Consider the prestack Schemes over Sch with objects $(T \rightarrow S)$ and morphisms $(T \rightarrow S) \rightarrow (T' \rightarrow S')$ is a pair $f: T \rightarrow T'$ and $g: S \rightarrow S'$ such that the two compositions $T \rightarrow S'$ agree. Of course schemes can be glued in the Zariski topology, so Sch is a stack in Sch_{Zar} .

Proposition 9.1.15. \mathcal{M}_g is a stack over $\text{Sch}_{\text{ét}}$ for $g \geq 2$.

Remark 9.1.16. The stackiness conditions mainly come from descent theory.

9.2 Algebraic stacks and Artin's axioms

From now on, we will work in the category Sch/S of schemes over S .

9.2.1 Algebraic stacks

Definition 9.2.1. A morphism $\mathcal{X} \rightarrow \mathcal{Y}$ of prestacks is *representable by schemes* if for all schemes $T \rightarrow \mathcal{Y}$, the fiber product $\mathcal{X} \times_{\mathcal{Y}} T$ is also a scheme. A representable $\mathcal{X} \rightarrow \mathcal{Y}$ is surjective, smooth, etc if for all schemes $T \rightarrow \mathcal{Y}$, the morphism $\mathcal{X} \times_{\mathcal{Y}} T \rightarrow T$ is surjective, smooth, etc.

Definition 9.2.2. An *algebraic space* is a sheaf F on $(\text{Sch}/S)_{\text{ét}}$ such that there exists a scheme U and a surjective étale $U \rightarrow F$ which is representable by schemes.

Definition 9.2.3. A morphism $\mathcal{X} \rightarrow \mathcal{Y}$ is called *representable* if for all schemes $T \rightarrow \mathcal{Y}$, $\mathcal{X} \times_{\mathcal{Y}} T$ is an algebraic space. Moreover, a representable $f: \mathcal{X} \rightarrow \mathcal{Y}$ is called surjective, smooth, etc if $U \rightarrow \mathcal{X} \times_{\mathcal{Y}} T \rightarrow T$ is surjective, smooth, etc.

Definition 9.2.4. A *algebraic stack* over $(\text{Sch}/S)_{\text{ét}}$ is a stack such that there exists a scheme U with a morphism $U \rightarrow \mathcal{X}$ that is smooth, surjective, and representable.

This is equivalent to the following conditions (taken together):

1. The diagonal $\mathcal{X} \xrightarrow{\Delta} \mathcal{X} \times \mathcal{X}$ is representable.
2. There exists a scheme U and a smooth surjective morphism $U \rightarrow \mathcal{X}$.

A useful fact that the product and fiber products of algebraic stacks exist (in the category of algebraic stacks), and this is the same as their fiber products as prestacks.

Definition 9.2.5. We say that $f: \mathcal{X} \rightarrow \mathcal{Y}$ of algebraic stacks is *locally of finite type* if for all (for some) smooth presentations $V \rightarrow \mathcal{Y}$ and $U \rightarrow \mathcal{X} \times_{\mathcal{Y}} V$, the composition

$$U \rightarrow \mathcal{X} \times_{\mathcal{Y}} V \rightarrow V$$

is locally of finite type.

9.3 Artin's Axioms

Definition 9.3.1. Let \mathcal{X} be a stack over $(\text{Sch}/S)_{\text{ét}}$. Then \mathcal{X} is *limit-preserving* if

$$\varinjlim \mathcal{X}(\text{Spec } B_i) \rightarrow \mathcal{X}(\text{Spec}(\varinjlim B_i))$$

is an equivalence of categories. Explicitly, this means:

1. Every object on the right hand side comes from $a_i|_{\text{Spec } B}$ for some i and some $a_i \in \mathcal{X}(\text{Spec } B_i)$.
2. For $a, b \in \mathcal{X}(\text{Spec } B_i)$, we have

$$\text{Hom}_{\text{RHS}}(a|_{\text{Spec } B}, b|_{\text{Spec } B}) = \varinjlim_{i' \geq i} \text{Hom}_{\mathcal{X}(\text{Spec } B_{i'})}(a|_{B_{i'}}, b|_{i'}).$$

This should be viewed as a finiteness condition because if $f: X \rightarrow S$ is a scheme, then $\text{Hom}(-, X)$ is limit-preserving if and only if f is locally of finite presentation.

Definitions 9.3.2. Let \mathcal{X} be a prestack over Sch/k . A *formal object* of \mathcal{X} is $(R, \{\xi_n\}, \{f_n\})$, where R is a complete local Noetherian k -algebra, $\xi_n: \text{Spec } R/\mathfrak{m}_R^n \rightarrow \mathcal{X}$, and $f_n: \xi_n \rightarrow \xi_{n+1}$ are morphisms in \mathcal{X} lying over $\text{Spec } R/\mathfrak{m}_R^n \hookrightarrow \text{Spec } R/\mathfrak{m}_R^{n+1}$.

A *morphism of formal objects* $(R, \{\xi_n\}, \{f_n\}) \rightarrow (T, \{\eta_n\}, \{g_n\})$ is a collection of morphisms $\alpha_n: \xi_n \rightarrow \eta_n$ such that

$$\begin{array}{ccc} \xi_n & \xrightarrow{\alpha_n} & \eta_n \\ \downarrow f_n & & \downarrow g_n \\ \xi_{n+1} & \xrightarrow{\alpha_{n+1}} & \eta_{n+1} \end{array}$$

commutes. These define morphisms $\text{Spec } R/\mathfrak{m}_R^n \rightarrow \text{Spec } T/\mathfrak{m}_T^n$ that are compatible, and hence a morphism $\text{Spec } R \rightarrow \text{Spec } T$.

There is a functor from $\mathcal{X}(\text{Spec } R)$ to the category of formal objects (R, \dots) . We say that a formal object is *effective* if it is in the essential image of this functor.

Definition 9.3.3. Let R be a complete local Noetherian k -algebra and $\xi \in \mathcal{X}(\text{Spec } R)$. We say that ξ is *versal* if for any diagram

$$\begin{array}{ccccc} \text{Spec } k & \longrightarrow & \text{Spec } B & \longrightarrow & \text{Spec } C \\ & & \downarrow & & \downarrow \xi \\ & & \text{Spec } B' & \xrightarrow{\eta'} & \mathcal{X} \end{array}$$

such that $B' \twoheadrightarrow B$ is a surjection of local Artinian k -algebras and $\alpha: \xi|_{\text{Spec } B} \rightarrow \eta'|_{\text{Spec } B}$ is an isomorphism, then there exists $\text{Spec } B' \rightarrow \text{Spec } R$ such that $\alpha' = \xi|_{\text{Spec } B'} \simeq \eta'$ extending α .

A formal object $(R, \{\xi_n\}, \{f_n\})$ is *versal* if in the same diagram, if we replace ξ with ξ_n and R with R/\mathfrak{m}_R^n , we can find a lift to $\text{Spec } R/\mathfrak{m}_R^m$ for some $m \geq n$.

Theorem 9.3.4 (Artin's axioms). *Let \mathcal{X} be a stack over $(\text{Sch}/S)_{\text{ét}}$. Then \mathcal{X} is an algebraic stack locally of finite type over k if and only if*

0. \mathcal{X} is limit-preserving.
1. The diagonal $\Delta: \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is representable.
2. (formal deformations) for every $x: \text{Spec } k \rightarrow \mathcal{X}$, there exists a complete local Noetherian k -algebra (R, \mathfrak{m}) and a versal formal object $(R, \{\xi_n\}, \{f_n\})$ such that $\xi_1 = x$.
3. Every formal object is effective.
4. (openness of versality) Let $\xi_{\mathcal{U}}: \mathcal{U} \rightarrow \mathcal{X}$, where \mathcal{U} is a scheme of finite type over k , and $u \in \mathcal{U}$ be a k -point such that $\xi_{\mathcal{U}}|_{\text{Spec } \widehat{\mathcal{O}}_{\mathcal{U}, u}}$ is versal. Then $\xi_{\mathcal{U}}$ is versal at all k -points in an open neighborhood of u .

Remark 9.3.5. Suppose we want to prove that Artin's axioms imply that \mathcal{X} is an algebraic stack locally of finite type over k . Given a formal object $(R, \{\xi_n\}, \{f_n\})$, we have an actual object $\xi \in \mathcal{X}(\text{Spec } R)$. By approximation and algebraization, we have $\mathcal{U} \rightarrow \mathcal{X}$ finite type and versal at $X = \mathcal{U}$. Now $\mathcal{U} \rightarrow \mathcal{X}$ is smooth, so we have $\bigsqcup \mathcal{U} \rightarrow \mathcal{X}$ smooth, surjective, and representable.

Remark 9.3.6. In many modular problems, the condition about formal deformations follows from the Rim-Schlessinger condition that for a diagram

$$\begin{array}{ccc} A \times_B C & \longrightarrow & C \\ \downarrow & & \downarrow \\ A & \longrightarrow & B \end{array}$$

of Artinian local k -algebras, the map

$$\mathcal{X}(\mathrm{Spec} A \times_B C) \rightarrow \mathcal{X}(A) \times_{\mathcal{X}(B)} \mathcal{X}(C)$$

is an equivalence of categories. Here, if $x_0 \in \mathcal{X}(\mathrm{Spec} k)$, define F_{C, x_0} by $A \mapsto x \in \mathcal{X}(\mathrm{Spec} A)$ such that $x_0 \rightarrow x$ lies over $\mathrm{Spec} k \rightarrow \mathrm{Spec} A$. Then $TF_{C, x_0} = F_{C, x_0}(k[\varepsilon])$ is a k -vector space. If $\dim_k TF_{C, x_0} < \infty$, then we have the condition on formal deformations. Then smoothness of $\mathcal{U} \rightarrow F_{C, x_0}$ gives versality.

Remark 9.3.7. The third condition follows from Grothendieck's existence theorem, and a deformation-obstruction theory gives openness of versality.

Che (Dec 03): Stack of coherent sheaves

Let k be an algebraically closed field of characteristic 0, $(\text{Sch}/k)_{\text{ét}}$ be the étale site of schemes over k , and X be a projective scheme over k .

Definition 10.0.1. Let Coh_X be the category defined as follows:

- Objects are tuples (T, \mathcal{F}) of a scheme T and \mathcal{F} is a quasicoherent sheaf on $X \times T$ of finite presentation flat over T .
- Morphisms from (T, \mathcal{F}) to (T', \mathcal{F}') are pairs (h, φ) , where $h: T \rightarrow T'$ is a morphism of schemes and $\varphi: (h')^* \mathcal{F}' \rightarrow \mathcal{F}$ is an isomorphism of \mathcal{O}_{X_T} -modules. Here, $h': X_T \rightarrow X_{T'}$ is the morphism induced from h .

There is a functor $p: \text{Coh}_X \rightarrow (\text{Sch}/k)_{\text{ét}}$ given by $(T, \mathcal{F}) \mapsto T$, and we want to prove that $\mathcal{X} := \text{Coh}_X$ is an algebraic stack. Also, we will abuse notation and write h for h' . We will prove that \mathcal{X} satisfies Artin's axioms. Recall that these are

- (0) \mathcal{X} is a stack. This means that \mathcal{X} is a prestack and objects and morphisms glue.
- (1) $\Delta: \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is representable by algebraic spaces.
- (2) \mathcal{X} is limit-preserving.
- (3) \mathcal{X} satisfies the Rim-Schlessinger condition.
- (4) The tangent spaces $T\mathcal{F}_{\mathcal{X}, X_0}$ and $\text{Inf}(\mathcal{F}_{\mathcal{X}, X_0})$ are finite-dimensional.
- (5) Every formal object is effective.
- (6) \mathcal{X} satisfies openness of versality.

These will imply that there is a smooth surjective covering of \mathcal{X} by a scheme.

10.1 \mathcal{X} is a stack

First, we will prove that \mathcal{X} is a prestack. We note that $\mathcal{X}(T)$ has objects finitely presented \mathcal{F} on X_T flat over T and morphisms (id_T, φ) , where $\varphi: \mathcal{F}' \rightarrow \mathcal{F}$ is an isomorphism. Thus $\mathcal{X}(T)$ is a groupoid. We now want to prove that pullbacks exist, which is clear because given $h: T \rightarrow T'$ and an object (T', \mathcal{F}') , our pullback is simply $(T, h^* \mathcal{F}')$. To show that pullbacks are universal, consider

$h_1: T_1 \xrightarrow{h} T_2 \xrightarrow{h_2} T$. Now suppose that we have morphisms $(T_1, \mathcal{F}_1) \rightarrow (T, \mathcal{F})$ and $(T_2, \mathcal{F}_2) \rightarrow (T, \mathcal{F})$. Now we want to prove that there exists an isomorphism $\mathcal{F}_1 \simeq h^*\mathcal{F}_2$. But we know that

$$h^*\mathcal{F}_2 \simeq h^*h_2^*\mathcal{F} \simeq h_1^*\mathcal{F} \simeq \mathcal{F}_1,$$

and so we are done.

Now we want to check that objects glue. Given T and a covering $\{a_i: T_i \rightarrow T\}$, suppose we have \mathcal{F}_i on X_{T_i} flat over T_i . Suppose that the \mathcal{F}_i are isomorphic on intersections and satisfy the cocycle condition. We want to construct a \mathcal{F} on X_T of finite presentation, flat over T . This follows from étale descent, so we are done.

We now want to prove that morphisms glue. To do this, we need to introduce some new notions.

Definition 10.1.1. Given $X = (T, \mathcal{F})$ and $Y = (T, \mathcal{G})$, define the *Isom presheaf* $\text{Isom}_{\mathcal{X}}(X, Y)$ sending a scheme $S \rightarrow T$ to the set $\text{Hom}(\mathcal{F}|_S, \mathcal{G}|_S)$ in $\mathcal{X}(S)$.

It is easy to see that morphisms glue if and only if this is a sheaf, and we will omit the proof.

10.2 Representability of the diagonal

We want to prove that for all schemes S over k , the stack \mathcal{Y} given by pullback in the diagram

$$\begin{array}{ccc} \mathcal{Y} & \longrightarrow & S \\ \downarrow & & \downarrow \\ \mathcal{X} & \xrightarrow{\Delta} & \mathcal{X} \times \mathcal{X} \end{array}$$

is an algebraic space. By the 2-Yoneda lemma, a map $S \rightarrow \mathcal{X} \times \mathcal{X}$ is given by $\xi = (S, \mathcal{F}), \eta = (S, \mathcal{G})$ in $\mathcal{X}(S)$. If we compute the fiber product $S \times_{\mathcal{X} \times \mathcal{X}} \mathcal{X}$, we actually obtain the Isom presheaf $\text{Isom}_{\mathcal{X}}(\xi, \eta)$. We will use without proof the fact that $\text{Isom}_{\mathcal{X}}(\xi, \eta)$ is a closed subfunctor of $\underline{\text{Hom}}(\mathcal{F}, \mathcal{G})$ (defined by $T \mapsto \text{Hom}(\mathcal{F}_T, \mathcal{G}_T)$).

We will only prove that $\underline{\text{Hom}}(\mathcal{F}, \mathcal{G})$ is representable by an algebraic space when \mathcal{F}, \mathcal{G} are locally free (and X is a point apparently).¹ In this case,

$$\underline{\text{Hom}}(\mathcal{F}, \mathcal{G})(T) = \text{Hom}(\mathcal{F}_T, \mathcal{G}_T) = H^0((\mathcal{F}^\vee \otimes \mathcal{G})_T).$$

But now this is clearly represented by the total space $\text{Spec Sym}(\mathcal{F} \otimes \mathcal{G}^\vee)$ because²

$$\begin{aligned} \text{Hom}_S(T, \text{Spec Sym}(\mathcal{F} \otimes \mathcal{G}^\vee)) &= \text{Hom}_{\mathcal{O}_S}(\text{Sym}(\mathcal{F} \otimes \mathcal{G}^\vee), f_*\mathcal{O}_T) \\ &= \text{Hom}_{\mathcal{O}_S}(\mathcal{F} \otimes \mathcal{G}^\vee, f_*\mathcal{O}_T) \\ &= \text{Hom}_{\mathcal{O}_T}(f^*(\mathcal{F} \otimes \mathcal{G}^\vee), \mathcal{O}_T) \\ &= H^0((\mathcal{F}^\vee \otimes \mathcal{G})_T). \end{aligned}$$

10.3 Preservation of limits

Given $T = \lim T_i$, we want to prove that $\mathcal{X}(T)$ is the colimit of the $\mathcal{X}(T_i)$. Given \mathcal{F} on $X \times T$ of finite presentation and flat over T , we want to show that there exist T_i, \mathcal{F}_i such that $\mathcal{F} = (X \rightarrow X_i)^*\mathcal{F}_i$.

¹Here, Johan intervened and came to the board to talk about cohomology and base change things.

²Apparently the following is wrong, but is preserved here for recordkeeping.

Proposition 10.3.1. *Let R, R_i be rings such that R is the colimit of the R_i . Let M be an R -module of finite presentation. Then there exists i and a finitely-presented R_i -module M_i such that $M = M_i \otimes_R R$.*

Proof. We know that M is finitely presented, so we have an exact sequence

$$R^{\oplus m} \xrightarrow{(a_{jk})} R^{\oplus n} \rightarrow M \rightarrow 0.$$

Because R is the colimit of the R_i , there exists i such that a_{jk} lift to R_i . If we define M_i by

$$R_i^{\oplus m} \xrightarrow{(a_{jk})} R_i^{\oplus n} \rightarrow M_i \rightarrow 0,$$

this is clearly the desired M_i . □

Checking flatness is too hard, so we will not do it.

10.4 Rim-Schlessinger

We will not prove this, but we will see that this is a natural condition to satisfy. Given a pushout diagram

$$\begin{array}{ccc} U & \longrightarrow & U' \\ \downarrow & & \downarrow \\ V & \longrightarrow & V' \end{array}$$

where U, U', V, V' are spectra of local Artinian rings of finite type over k and $U \rightarrow U'$ is a closed embedding, we want the functor

$$\mathcal{X}(V') \rightarrow \mathcal{X}(V) \times_{\mathcal{X}(U)} \mathcal{X}(U')$$

to be an equivalence of categories.

10.5 Finiteness

We want to prove that the tangent space $T\mathcal{F}_{\mathcal{X}, X_0}$ and the infinitesimal automorphisms $\text{Inf}(\mathcal{F}_{\mathcal{X}, X_0})$ are finite-dimensional. Given $X_0: \text{Spec } k \rightarrow \mathcal{X}$ (which is just a finitely presented sheaf \mathcal{F} on X), the tangent space is

$$T\mathcal{F}_{\mathcal{X}, X_0} := \{ \mathcal{F}'/X \times k[\varepsilon] \text{ finitely-presented, flat over } k[\varepsilon], \mathcal{F}'|_X \cong \mathcal{F} \} / \sim = \text{Ext}^1(\mathcal{F}, \mathcal{F}).$$

In addition, we know that

$$\text{Inf}(\mathcal{F}_{\mathcal{X}, X_0}) = \ker(\text{Aut}(\mathcal{F} \otimes k[\varepsilon]/X \times k[\varepsilon]) \rightarrow \text{Aut}(\mathcal{F}/X)) = \text{Ext}^0(\mathcal{F}, \mathcal{F}).$$

Because X is projective, these Ext groups are finite-dimensional.

10.6 Formal objects are effective

We will now prove that formal objects are effective. Let $R \in \widehat{\mathcal{C}}$ be a complete Noetherian local ring. Recall that a formal object is $\{\xi_n\}$, where $\xi_n \in \mathcal{X}(\text{Spec } R/\mathfrak{m}_R^n)$ along with $f_n: \xi_n \rightarrow \xi_{n+1}$ living over the natural inclusions. A formal object is effective if it comes from an actual object over R .

In our case, a formal object is given by \mathcal{F}_n on $X \times \text{Spec } R/\mathfrak{m}_R^n$ flat and finitely presented and $f_n: i_n^* \mathcal{F}_{n+1} \cong \mathcal{F}_n$. We want to show that there exists \mathcal{F} over $\text{Spec } R$ restricting to each \mathcal{F}_n .

Theorem 10.6.1 (Grothendieck existence theorem). *Let A be a Noetherian ring which is complete with respect to some ideal I . Let $f: X \rightarrow \text{Spec } A$ be a proper morphism. Let $\mathcal{J} = I\mathcal{O}_X$. Then the functor*

$$\text{Coh}(X) \rightarrow \left\{ \mathcal{F}_\infty \leftarrow \mathcal{F}_\epsilon \leftarrow \cdots \mid \mathcal{F}_\epsilon \text{ coherent, annihilated by } \mathcal{J}^\epsilon, \mathcal{F}_{\epsilon+1}/\mathcal{J}^\epsilon \mathcal{F}_{\epsilon+1} \simeq \mathcal{F}_\epsilon \right\}$$

is an equivalence.

In our case, take $A = \mathbb{R}$ and $I = \mathfrak{m}$, and now we are done.

Johan (Dec 10): Openness of versality

Let k be an algebraically closed field. In this lecture, all schemes live over k for simplicity.

11.1 Moduli of curves

Recall that $\overline{\mathcal{M}}_g$ is a stack such that a morphism $U \rightarrow \overline{\mathcal{M}}_g$ is a family $C \rightarrow U$ of stable curves of genus g . Assume that U is of finite type over k and let $u_0 \in U(k)$.

Definition 11.1.1. We say that $U \rightarrow \overline{\mathcal{M}}_g$ is *versal* at u_0 if $\widehat{\mathcal{O}}_{U, u_0}$ and the map

$$h_{\widehat{\mathcal{O}}_{U, u_0}} \rightarrow \text{Def}_{C_{u_0}}$$

given by $C|_{\text{Spec } \widehat{\mathcal{O}}_{U, u_0}}$ are a hull.

Lemma 11.1.2. $U \rightarrow \overline{\mathcal{M}}_g$ is versal if and only if U is smooth at u_0 and $T_{u_0}U \rightarrow T\text{Def}_{C_{u_0}}$ is surjective.

Proof. Earlier, we discussed that deformations of C_{u_0} are unobstructed. Therefore any hull is a power series ring over k . Thus U must be smooth at u_0 . If U is smooth at u_0 , then look at $\widehat{\mathcal{O}}_{U, u_0} \leftarrow R$, where R is the deformation ring of C_{u_0} . But now this is a map of power series rings, and therefore defines a smooth transformation of functors if and only if the map on tangent spaces is surjective. \square

Lemma 11.1.3. We have openness of versality for $\overline{\mathcal{M}}_g$ and \mathcal{M}_g .

Proof for \mathcal{M}_g . By the previous lemma, we may assume that U is smooth. Call $f: C \rightarrow U$ and consider the exact sequence

$$0 \rightarrow T_{C/U} \rightarrow T_C \rightarrow f^*T_U \rightarrow 0.$$

This gives $T_U = f_*f^*T_U \rightarrow R^1f_*T_{C/U}$. Taking fibers at u_0 , we obtain

$$T_{u_0}U = T_U \otimes \kappa(u_0) \rightarrow R^1f_*T_{C/U} \otimes \kappa(u_0) = H^1(C_{u_0}, T_{C_{u_0}}) = T_{u_0}\text{Def}_{C_{u_0}}.$$

Also, $R^1f_*T_{C/U}$ is a vector bundle of rank $3g - 3$ over U . By the lemma and the assumption of versality at u_0 , we see this is surjective. Thus this is surjective in an open neighborhood. \square

Proof for $\overline{\mathcal{M}}_g^1$. Consider the exact sequence

$$0 \rightarrow f^* \Omega_{U/k} \rightarrow \Omega_{C/k} \rightarrow \Omega_{C/U} \rightarrow 0.$$

We think of this as $\Omega_{C/U} \rightarrow f^* \Omega_{U/k}[1]$ in the derived category, and now if we tensor with the relative dualizing sheaf, we have

$$\Omega_{C/U} \otimes \omega_{C/U} \rightarrow f^* \Omega_{U/k} \otimes \omega_{C/U}[1] = f^!(\Omega_{U/k}),$$

where $f^!$ is the right adjoint to Rf_* in this case. This gives

$$Rf_*(\Omega_{C/U} \otimes \omega_C) \rightarrow \Omega_{U/k}.$$

By positivity properties, $Rf_*(\Omega_{C/U} \otimes \omega_{C/U}) = f_*(\Omega_{C/U} \otimes \omega_{C/U})$. Taking the fiber at u_0 , we have

$$H^0(\Omega_{C_{u_0}/k} \otimes \omega_{C_{u_0}})^\vee = \text{Ext}^1(\Omega_{C_{u_0}/k} \otimes \omega_{C_{u_0}}, \omega_{C_{u_0}/k}) = \text{Ext}_{C_{u_0}}^1(\Omega_{C_{u_0}}, \mathcal{O}_{C_{u_0}}) \cong T_{u_0} \text{Def}_{C_{u_0}}.$$

By the same argument as before, we are done. \square

11.2 Properties of cotangent complexes

Suppose that U is of finite type over k . Then there is a complex $L_{U/k} \in D_{\text{coh}}^{\leq 0}(\mathcal{O}_U)$ such that $H^0(L_{U/k}) = \Omega_{U/k}$ called the *cotangent complex*. Suppose we write

$$\widehat{\mathcal{O}}_{U, u_0} \cong k[[x_1, \dots, x_n]] / (f_1, \dots, f_m) = k[[x]] / I$$

with n, m minimal. This implies that $f_1, \dots, f_m \in (x_1, \dots, x_n)^2$. Then $H^0(L_{U/k} \otimes \kappa(u_0))$ is the cotangent space of U at u_0 and thus has dimension n . Next we note that

$$H^{-1}L_{U/k} \otimes \kappa(u_0) = I/mI,$$

and this has dimension m .

Now let $g: U \rightarrow V$ be a morphism of schemes of finite type over k . Write $u_0 \mapsto v_0$. Then if we consider the distinguished triangle

$$Lg^*L_{V/k} \rightarrow L_{U/k} \rightarrow L_{U/V},$$

we see that g is smooth at u_0 if and only if $H^0(L_{U/k} \otimes \kappa(u_0)) \leftarrow H^0(L_{V/k} \otimes \kappa(v_0))$ is injective and $H^{-1}(L_{U/k} \otimes \kappa(u_0)) \leftarrow H^{-1}(L_{V/k} \otimes \kappa(v_0))$ is surjective.

Remark 11.2.1. Suppose that $f: X \rightarrow U$ is a proper flat morphism corresponding to $U \rightarrow \mathcal{X}$, where \mathcal{X} is the prestack parameterizing families of flat proper schemes. Then we have $L_{X/U} \rightarrow Lf^*L_{U/k}[1]$, and tensoring with the dualizing complex, we have

$$L_{X/U} \otimes \omega_{X/U}^\bullet \rightarrow Lf^*L_{U/k} \otimes \omega_{X/U}^\bullet[1] = f^!(L_{U/k})[1].$$

We should consider instead

$$\text{can}_{X/U}: Rf_*(L_{X/U} \otimes \omega_{X/U}^\bullet)[-1] \rightarrow L_{U/k}.$$

Versality is then related to properties of $H^i(\text{can}_{X/U} \otimes^{\mathbb{L}} \kappa(u_0))$ for $i = 0, -1$.

11.3 Coherent sheaves

Let $X \rightarrow \text{Spec } k$ be proper. Then $\text{Coh}_{X/k}$ is a stack such that for all U/k of finite type, a morphism $U \rightarrow \text{Coh}_{X/k}$ is a coherent sheaf \mathcal{F} on $X \times U$ which is flat over U .

Lemma 11.3.1. *We have openness of versality for $U \rightarrow \text{Coh}_{X/k}$.*

Easy case. We will assume that \mathcal{F}_{u_0} is a vector bundle and $\text{Ext}_X^2(\mathcal{F}_{u_0}, \mathcal{F}_{u_0}) = 0$. As before, we may assume that U is smooth and \mathcal{F} is a vector bundle. We then have the Atiyah extension

$$0 \rightarrow \mathcal{F} \otimes \Omega_{X \times U}^1 \rightarrow \mathcal{P}(\mathcal{F}) \rightarrow \mathcal{F} \rightarrow 0$$

and a map $\mathcal{F} \otimes \Omega_{X \times U/k}^1 \rightarrow \mathcal{F} \otimes p^* \Omega_{U/k}$. Then we have the Atiyah class

$$\mathcal{F} \rightarrow (\mathcal{F} \otimes \otimes_{X \times U/k}^\infty)[\infty].$$

Dually, we have $f^* T_{U/k} \rightarrow \text{Hom}(\mathcal{F}, \mathcal{F})[1]$,

$$T_{U/k} \rightarrow \text{Rf}_* f^* T_{U/k} \rightarrow \text{Rf}_*(\text{Hom}(\mathcal{F}, \mathcal{F}))[1],$$

etc. By similar arguments as before, we obtain the desired result. \square

General case (terrible). We have an Atiyah class

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\xi} & \mathcal{F} \otimes L_{X \times U/k}[1] \\ & \searrow \xi' & \\ & & \mathcal{F} \otimes L_{p^* L_{U/k}}[1]. \end{array}$$

This yields

$$\xi'' : \mathcal{F} \otimes \text{RHom}(\mathcal{F}, q^* \omega_{X/k}^\bullet) \rightarrow p^* L_{U/k} \otimes q^* \omega_{X/k}^\bullet[1] = p^!(L_{U/k}).$$

The adjunction gives us

$$\text{Rp}_*(\mathcal{F} \otimes \text{RHom}(\mathcal{F}, q^* \omega_{X/k}^\bullet))[-1] \rightarrow L_{U/k}.$$

We need to show that formation of the left hand side commutes with base change and that

$$H^i(X, \mathcal{F}_{u_0} \otimes \mathcal{F}_{u_0}, \omega_{X/k}^\bullet) = \text{Ext}_X^{-i}(\mathcal{F}_{u_0}, \mathcal{F}_{u_0}),$$

and then we can use cohomology and base change. \square

11.4 A trick

Sometimes we can get openness of versality for a prestack \mathcal{X} . Here, openness of versality holds for the prestack \mathcal{X} if

1. $\mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is representable by algebraic spaces.
2. We have the condition (RS_*) , which is a version of RS where the rings do not need to be Artinian.
3. \mathcal{X} is limit-preserving.

4. The following effectiveness holds: If $A = \lim A_n$ where

$$A \rightarrow \cdots \rightarrow A_3 \rightarrow A_2 \rightarrow A_1$$

where each $A_n \rightarrow A_{n-1}$ is surjective with square zero kernel, then $\mathcal{X}(A) = \lim \mathcal{X}(A_n)$.

In the example on $\text{Coh}_{X/k}$, we have \mathcal{F}_n on $X \otimes A_n$, and then we can attempt to take the limit \mathcal{F} of the \mathcal{F}_n . This fails because the limit is not quasicohherent, but we can fix it.