

RANK 2 LOCAL SYSTEMS AND ABELIAN VARIETIES

RAJU KRISHNAMOOTHY AND AMBRUS PÁL

ABSTRACT. Let X/\mathbb{F}_q be a smooth geometrically connected variety. Inspired by work of Corlette-Simpson over \mathbb{C} , we formulate a conjecture that absolutely irreducible rank 2 local systems with infinite monodromy on X come from families of abelian varieties. When X is a projective variety, we prove a Lefschetz-style theorem for abelian schemes of GL_2 -type on X , modeled after a theorem of Simpson. If one assumes a strong form of Deligne's *(p-adic) companions conjecture* from Weil II, this implies that the our conjecture for projective varieties also reduces to the case of projective curves. We also answer affirmatively a question of Grothendieck on extending abelian schemes via their p -divisible groups.

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Notation and Conventions | 5 |
| 3. Coefficient Objects | 5 |
| 4. Compatible Systems and Companions | 8 |
| 5. Barsotti-Tate Groups and Dieudonné Crystals | 11 |
| 6. A question of Grothendieck | 13 |
| 7. Lefschetz Theorems I | 17 |
| 8. Lefschetz Theorems II | 20 |
| References | 25 |

1. INTRODUCTION

The goal of this article is to prove the following Lefschetz-style theorem. If X/k is a smooth variety over a perfect field, $\mathbf{F}\text{-Isoc}(X)_{\overline{\mathbb{Q}}_p}$ denotes the category of F -isocrystals on X with coefficients in $\overline{\mathbb{Q}}_p$. On a smooth *proper* variety X/\mathbb{F}_q , an F -isocrystal is a p -adic analog of a lisse l -adic sheaf.

Theorem (8.11). *Let X/\mathbb{F}_q be a smooth projective variety. Then there exists an open subset $U \subset X$, whose complement has codimension at least 2, such that the following holds.*

Let \mathcal{E} be a semi-simple object of $\mathbf{F}\text{-Isoc}(X)_{\overline{\mathbb{Q}}_p}$ such that

- *the irreducible constituents of \mathcal{E} have rank 2 and determinant $\overline{\mathbb{Q}}_p(-1)$, and*
- *there exists a smooth projective curve $C \subset U$, the complete intersection of smooth ample divisors in X , and an abelian scheme $A_C \rightarrow C$ of dimension g such that*

$$\mathbb{D}(A_C[p^\infty]) \otimes \overline{\mathbb{Q}}_p \cong \mathcal{E}_C$$

Then there exists an abelian scheme $B_U \rightarrow U$ of dimension g such that

$$\mathbb{D}(B_U[p^\infty]) \otimes \overline{\mathbb{Q}}_p \cong \mathcal{E}_U.$$

In particular, $A_C \rightarrow C$ extends to an abelian scheme over U after an isogeny.

Here, if $G \rightarrow S$ is a p -divisible group in characteristic p , $\mathbb{D}(G)$ denotes the contravariant Dieudonné crystal attached to G . This theorem is modeled on the following, which easily follows from a very special case of a corollary of Simpson.

Theorem 1.1. [Sim92, Corollary 4.3] *Let X/\mathbb{C} be a smooth projective variety and $C \subset X$ a smooth curve that is the complete intersection of smooth ample divisors. Let L_X be a \mathbb{C} -local system on X and let L_C denote the restriction to C . Suppose there exists an abelian scheme $\pi_C: A_C \rightarrow C$ such that $R^1(\pi_C^{an})_*\mathbb{C} \cong L_C$. Then there is a unique abelian scheme $\pi_X: A_X \rightarrow X$ extending $A_C \rightarrow C$, and*

$$R^1(\pi_X^{an})_*\mathbb{C} \cong L_X.$$

Therefore, Theorem 8.11 is an analog of Theorem 1.1 over \mathbb{F}_q for abelian schemes of GL_2 -type. In contrast to Theorem 1.1, Theorem 8.11 has an intervening $U \subset X$ and also an isogeny.

The authors conjecture the following. For the definition of l -adic companions, see Remark 4.2.

Conjecture 1.2. (Conjecture R2) *Let X/\mathbb{F}_q be a smooth, geometrically connected, quasi-projective variety, let $l \neq p$ be a prime, and let L be a lisse $\overline{\mathbb{Q}}_l$ -sheaf of rank 2 such that*

- L has determinant $\overline{\mathbb{Q}}_l(-1)$ and
- L is irreducible with infinite geometric monodromy.

Then L comes from a family of abelian varieties: there exists a non-empty open $U \subset X$ together with an abelian scheme

$$\pi: A_U \rightarrow U$$

such that

$$R^1\pi_*\overline{\mathbb{Q}}_l \cong \bigoplus (\sigma L|_U)^m$$

where σL runs over the l -adic companions of L and $m \in \mathbb{N}$.

In particular, we conjecture such L come from H^1 of an abelian scheme on an open $U \subset X$. The main evidence for Conjecture 1.2 comes from Drinfeld's first work on the Langlands correspondence.

Theorem 1.3. (Drinfeld) *Let C/\mathbb{F}_q be a smooth affine curve and let L be as in Conjecture 1.2. Suppose L has infinite (geometric) monodromy around some point at $\infty \in \overline{C} \setminus C$. Then L comes from a family of abelian varieties in the following sense: let E be the field generated by the Frobenius traces of L and suppose $[E : \mathbb{Q}] = g$. Then there exists an abelian scheme*

$$\pi: A_C \rightarrow C$$

of dimension g and an isomorphism $E \cong \text{End}_C(A) \otimes \mathbb{Q}$, realizing A_C as a GL_2 -type abelian scheme, such that L occurs as a summand of $R^1\pi_\overline{\mathbb{Q}}_l$. Moreover, $A_C \rightarrow C$ is totally degenerate around ∞ .*

See [ST18, Proof of Proposition 19, Remark 20] for how to recover this result from Drinfeld's work.¹ For general smooth X/\mathbb{F}_q , an *overconvergent F -isocrystal* is a good p -adic analog of a lisse l -adic sheaf. Conjecture 1.2 can then be formulated in the $l = p$ case, replacing L with $\mathcal{E} \in \mathbf{F}\text{-Isoc}^\dagger(X)_{\overline{\mathbb{Q}}_p}$, an overconvergent F -isocrystal with coefficients in $\overline{\mathbb{Q}}_p$. If X is projective, then any F -isocrystal is automatically overconvergent. Combined with a refined form of Deligne's p -adic companions conjecture, we obtain the following application to Conjecture 1.2.

Corollary (8.13). *Let X/\mathbb{F}_q be a smooth projective variety with $\dim(X) \geq 2$ and let L be a rank 2 lisse $\overline{\mathbb{Q}}_l$ sheaf with cyclotomic determinant and infinite geometric monodromy. Suppose there exists a complete set of p -adic companions (\mathcal{E}_v) for L (as predicted by the p -adic companions conjecture). Then there exists an open subset $U \subset X$, whose complement has codimension at least 2, such that*

- if $C \subset U$ is a smooth proper curve that is the complete intersection of smooth ample divisors and
- if L_C comes from an abelian scheme on $A_C \rightarrow C$, in the sense of Conjecture 1.2,

then L_U comes from an abelian scheme $B_U \rightarrow U$, i.e., Conjecture 1.2 is true for (X, L) .

¹This amounts to combining [Dri83, Main Theorem, Remark 5] with [Dri77, Theorem 1].

In other words, if one believes the p -adic companions conjecture, then Conjecture 1.2 for a projective variety reduces to the case of a single sufficiently generic curve $C \subset X$. (See Definition 4.5 for the definition of a *complete set of p -adic companions*.) In general, the existence of a complete set of p -adic companions is a strong form of Deligne’s *petits camarades cristallin* conjecture: see Conjectures 4.8 and 4.9. When we place certain conditions on the splitting of p in E , the field of traces of \mathcal{E} , the existence of a single p -adic companion guarantees the existence of all of them by Corollary 4.16.

We briefly describe the strategy. First, we construct a (non-canonical) p -divisible group on $U \subset X$. Then, we use Serre-Tate theory to construct a formal (polarizable) abelian scheme over the formal scheme X/C . Here the positivity of $C \subset X$ is used. Finally, using work of Grothendieck, Hartshorne, and Hironaka and the positivity of $C \subset X$, we globalize the family. As a key step, we record an affirmative answer to a question of Grothendieck [Gro66, 4.9]:

Theorem. (Corollary 6.8) *Let X be a locally noetherian normal scheme and $U \subset X$ be an open dense subset whose complement has characteristic p . Let $A_U \rightarrow U$ be an abelian scheme. Then A_U extends to an abelian scheme over X if and only if $A_U[p^\infty]$ extends to a p -divisible group over X .*

Combined with algebraization techniques, there is the following useful consequence, which is a p -adic analog of Simpson’s Theorem 1.1.

Corollary. (Corollary 8.6) *Let X/\mathbb{F}_q be a smooth projective variety and let $C \subset X$ be a smooth curve that is the complete intersection of smooth ample divisors. Let $A_C \rightarrow C$ be an abelian scheme. Suppose there exists a Zariski open neighborhood $U \supset C$ of X such that $A_C[p^\infty]$ extends to a Barsotti-Tate group \mathcal{G}_U on U . Then there exists a unique abelian scheme $A_U \rightarrow U$, extending A_C , such that $A_U[p^\infty] \cong \mathcal{G}_U$.*

Remark 8.10 shows that hypothesis on the dimension is necessary. The proof of Corollary 8.6 makes use of p -to- l companions. See Remark 8.8 and Corollary 8.9 for what we know over general fields k of characteristic p . We now make some general remarks on Conjecture 1.2.

Remark 1.4. Our motivation to formulate Conjecture 1.2 partly comes from the following celebrated theorem of Corlette-Simpson [CS08, Theorem 11.2]:

Theorem. (Corlette-Simpson) *Let X/\mathbb{C} be a smooth, connected, quasi-projective variety and let L be a rank 2 \mathbb{C} -local system on X such that*

- L has trivial determinant,
- L has quasi-unipotent monodromy along the divisor at ∞ ,
- L has Zariski-dense monodromy inside of $SL_2(\mathbb{C})$, and
- L is rigid.

Then L comes from a family of abelian varieties: there exists an abelian scheme $\pi: A_X \rightarrow X$ such that

$$R^1\pi_*\mathbb{C} \cong \bigoplus_{\sigma \in \Sigma} (\sigma L)^m$$

where $\Sigma \subset \text{Aut}(\mathbb{C})$ is a finite subset of automorphisms of \mathbb{C} containing the identity, σL is the local system obtained by applying σ to the matrices in the associated $SL_2(\mathbb{C})$ representation, and $m \in \mathbb{N}$.

When X is projective, L being rigid means it yields an isolated (though not necessarily reduced) point in the character variety associated to $\pi_1(X)$. For general quasi-projective X , the notion of rigidity involves a character variety that remembers Jordan blocks of the various monodromies around ∞ , see [CS08, Section 6] or [EG17, Section 2] for a precise definition. Corlette-Simpson’s theorem verifies the rank 2 case of a conjecture of Simpson that roughly states: rigid semi-simple \mathbb{C} -local systems on smooth complex varieties are motivic. ²

Remark 1.5. It is conjectured that if X/\mathbb{F}_q is an irreducible, smooth variety and L is a lisse $\overline{\mathbb{Q}}_l$ -sheaves L with trivial determinant, then L is “of geometric origin” up to a Tate twist (Esnault-Kerz attribute this to Deligne [EK11, Conjecture 2.3], see also [Dri12, Question 1.4]). More precisely, given such an L , it is conjectured that there exists an open dense subset $U \subset X$, a smooth projective morphism

²When X is proper, this is [Sim91, Conjecture 4].

$\pi : Y_U \rightarrow U$, an integer i , and a rational number j such that $L|_U$ is a sub-quotient (or even a summand) of $R^i\pi_*\overline{\mathbb{Q}}_l(j)$.

Remark 1.6. Compared to Corlette-Simpson’s theorem, there are two fewer hypotheses in Conjecture 1.2: there is no “quasi-unipotent monodromy at ∞ ” condition and there is no rigidity condition. The former is automatic by Grothendieck’s quasi-unipotent monodromy theorem [ST68, Appendix].

As for the latter: the local systems showing up in Simpson’s work are representations of a *geometric fundamental group*, while the local systems occurring in Conjecture 1.2 are representations of an *arithmetic fundamental group*. Let X/\mathbb{F}_q be a smooth variety. Then it follows from [Del12] that there are only finitely many isomorphism classes of lisse $\overline{\mathbb{Q}}_l$ -sheaves that have finite order determinant, bounded rank, and bounded ramification [EK12, Theorem 5.1]. For this reasons, the authors view such local systems as rigid.³

Remark 1.7. A further difference between Conjecture 1.2 and Corlette-Simpson’s theorem is the intervention of an open set $U \subset X$ in the positive characteristic case. If $U \subset X$ is a Zariski open subset of a smooth and irreducible \mathbb{C} -variety with complement of codimension at least 2, then any map $U \rightarrow \mathcal{A}_g \otimes \mathbb{C}$ extends (uniquely) to a map $Y \rightarrow \mathcal{A}_g \otimes \mathbb{C}$ [Gro66, Corollaire 4.5]. This extension property is not true in characteristic p ; one may construct counterexamples using a family of supersingular abelian surfaces over $\mathbb{A}^2 \setminus \{(0,0)\}$ [Gro66, Remarques 4.6]. On the other hand, we do not know of a single example that requires the intervention of an open subset $U \subsetneq X$ in Conjecture 1.2.

Remark 1.8. There is recent work of Snowden-Tsimerman that uses Drinfeld’s results to prove something similar to Conjecture 1.2 for curves over a number field and rank 2 local systems with Frobenius traces in \mathbb{Q} [ST18]. This work was very inspiring for us, but the techniques used there are rather different from those used here. In particular, they use Drinfeld’s result “mod p ” for infinitely many p , together with a Hilbert scheme argument based on the boundedness of maps between projective hyperbolic curves. Our techniques instead use deformation theory and algebraization.

We briefly summarize the sections.

- Section 3 sets out definitions and conventions for lisse l -adic sheaves and (overconvergent) F -isocrystals. Following Kedlaya, if X/\mathbb{F}_q is a smooth variety, we define a *coefficient object* to be either a lisse l -adic sheaf or an overconvergent F -isocrystal.
- Section 4 briefly describes the companions conjecture.
- Section 5 reviews what we need from the theory of Barsotti-Tate (a.k.a. p -divisible) groups.
- Section 6 proves that an abelian scheme on a noetherian, normal scheme extends across a closed subset of characteristic p if and only if its p -divisible group extends.
- Section 7 recalls a result of Hartshorne that allows us to globalize formal maps.
- Section 8 puts all of the ingredients together and contains a proof of Theorem 8.11.

The authors hope this work helps further reveal the geometric content of p -adic coefficient objects; more specifically that they are analogous to variations of Hodge structures. Deligne’s Conjecture 4.8 was presumably formulated with the hope that such local systems are of geometric origin; here, we deduce “geometric origin” conclusions from the existence of p -adic companions.

Acknowledgement. This work was born at CIRM (in Luminy) at “ p -adic Analytic Geometry and Differential Equations”; the authors thank the organizers. R.K. warmly thanks Tomoyuki Abe, Ching-Li Chai, Marco d’Addezio, Philip Engel, Hélène Esnault, and especially Johan de Jong, with whom he had stimulating discussions on the topic of this article. R.K. also thanks Daniel Litt, who patiently and thoroughly explained several algebraization techniques; in particular, Litt explained the proof of Lemma 8.5. R.K. thanks the arithmetic geometry group at FU Berlin for a lively and intellectually rich atmosphere. R.K. thanks Dino Lorenzini for comments on an earlier draft of this article. R.K. gratefully acknowledges financial support from the NSF under Grants No. DMS-1605825 and No. DMS-1344994 (RTG in Algebra, Algebraic Geometry and Number Theory at the University of Georgia).

³See also [Lit18, Theorem 3.2.1] for another indication of rigidity.

2. NOTATION AND CONVENTIONS

- The field with p elements is denoted by \mathbb{F}_p and \mathbb{F} denotes a fixed algebraic closure.
- A variety X/k is a geometrically integral scheme of finite type.
- If X is a scheme, then X° is the set of closed points.
- If k is a field, l denotes a prime different than $\text{char}(k)$.
- If E is a number field, then λ denotes an arbitrary prime of E .
- An λ -adic local field is a *finite* extension of \mathbb{Q}_λ .
- If L/K is a finite extension of fields and \mathcal{C} is a K -linear abelian category, then \mathcal{C}_L is the base-changed category. If M/K is an algebraic extension, then \mathcal{C}_M is the 2-colimit of the categories \mathcal{C}_L as M ranges through the finite extensions of K contained in M : $K \subset L \subset M$.
- If $\mathcal{G} \rightarrow S$ is a p -divisible (a.k.a. Barsotti-Tate) group, then $\mathbb{D}(\mathcal{G})$ denotes the contravariant Dieudonné crystal.
- If X/k is a smooth scheme of finite type over a perfect field k , then $\mathbf{F}\text{-Isoc}^\dagger(X)$ is the category of overconvergent F -isocrystals on X .
- If X is a noetherian scheme and $Z \subsetneq X$ is a non-empty closed subscheme, then X/Z is the formal completion of X along Z .

3. COEFFICIENT OBJECTS

For a more comprehensive introduction to the material of this section, see the recent surveys of Kedlaya [Ked16, Ked18]; our notations are consistent with his (except for us $l \neq p$ and λ denotes an arbitrary prime). Throughout this section, k denotes a perfect field, $W(k)$ the ring of Witt vectors, $K(k)$ the field of fractions of $W(k)$, and σ is the canonical lift of absolute Frobenius on k .

Let X/\mathbb{F}_q be a normal variety and \mathbb{F} a fixed algebraic closure of \mathbb{F}_q . Denote by \overline{X} the base change $X \times_{\text{Spec } \mathbb{F}_q} \text{Spec } \mathbb{F}$. We have the following homotopy exact sequence

$$0 \rightarrow \pi_1(\overline{X}) \rightarrow \pi_1(X) \rightarrow \text{Gal}(\mathbb{F}/\mathbb{F}_q) \rightarrow 0$$

(suppressing the implicit geometric point required to define π_1). The profinite group $\text{Gal}(\mathbb{F}/\mathbb{F}_q)$ has a dense subgroup \mathbb{Z} generated by the Frobenius. The inverse image of this copy of \mathbb{Z} in $\pi_1(X)$ is called the *Weil Group* $W(X)$ [Del80, 1.1.7]. The Weil group is given a topology where \mathbb{Z} is discrete; this is not the subspace topology from $\pi_1(X)$.

Definition 3.1. [Del80, 1.1.12] Let X/\mathbb{F}_q be a normal variety and K an l -adic local field. A (*lisse*) *Weil sheaf of rank r with coefficients in K* is a continuous representation $W(X) \rightarrow GL_r(K)$. A (*lisse*) *étale sheaf of rank r with coefficients in K* is a continuous representation $\pi_1(X) \rightarrow GL_r(K)$. We denote the category of Weil sheaves with coefficients in K by $\mathbf{Weil}(X)_K$.

Every lisse étale sheaf yields a lisse Weil sheaf. Conversely, any lisse Weil sheaf with finite determinant is the restriction to $W(X)$ of an étale sheaf [Del80, 1.3.4].

Definition 3.2. Let \mathcal{C} be a K -linear additive category, where K is a field [Sta20, Tag 09MI]. Let L/K be a finite field extension. We define the base-changed category \mathcal{C}_L as follows:

- Objects of \mathcal{C}_L are pairs (M, f) , where M is an object of \mathcal{C} and $f: L \rightarrow \text{End}_{\mathcal{C}} M$ is a homomorphism of K -algebras. We call such an f an L -structure on M .
- Morphisms of \mathcal{C}_L are morphisms of \mathcal{C} that are compatible with the L -structure.

Fact 3.3. *Let \mathcal{C} be a K -linear abelian category and let L/K be a finite field extension. Then there are functors*

$$\text{Ind}_K^L: \mathcal{C} \rightleftarrows \mathcal{C}_L: \text{Res}_K^L$$

called induction and restriction. Restriction is right adjoint to induction. Both functors send semi-simple objects to semi-simple objects. If \mathcal{C} has the structure of a Tannakian category, then so does \mathcal{C}_L . In this case, Ind_K^L preserves rank and Res_K^L multiplies the rank by $[L : K]$.

Proof. The restriction functor is given by forgetting the L -structure. For a description of induction and restriction functors, see [Kri17, Section 3]. The fact about semi-simple objects follows immediately from [Kri17, Corollary 3.12]. For the fact that \mathcal{C}_L is Tannakian, see [Del14]. \square

Given an object M of \mathcal{C} , we will sometimes write M_L or $M \otimes_K L$ for $\text{Ind}_K^L M$.

The category of \mathbb{Q}_l -Weil sheaves, $\mathbf{Weil}(X)$, is naturally an \mathbb{Q}_l -linear neutral Tannakian category, and $\mathbf{Weil}(X)_K \cong (\mathbf{Weil}(X))_K$, where the latter denotes the “based-changed category” as above. We define $\mathbf{Weil}(X)_{\overline{\mathbb{Q}_l}}$ as the 2-colimit of $\mathbf{Weil}(X)_K$ as $K \subset \overline{\mathbb{Q}_l}$ ranges through the finite extensions of \mathbb{Q}_l inside of $\overline{\mathbb{Q}_l}$. Alternatively $\mathbf{Weil}(X)_{\overline{\mathbb{Q}_l}}$ is the category of continuous, finite dimensional representations of $W(X)$ in $\overline{\mathbb{Q}_l}$ -vector spaces where $\overline{\mathbb{Q}_l}$ is equipped with the colimit topology.

Definition 3.4. Let X/\mathbb{F}_q be a smooth variety and let $x \in X^\circ$ be a closed point of X . Given a Weil sheaf L on X with coefficients in $\overline{\mathbb{Q}_l}$, by restriction we get a Weil sheaf L on x . Denote by Fr_x the geometric Frobenius at x . Then $Fr_x \in W(x)$ and we define $P_x(L, t)$, the *characteristic polynomial of L at x* , to be

$$P_x(L, t) := \det(1 - Fr_x t | L_x).$$

Now, let X/k be a scheme of finite type over a perfect field. Berthelot has defined the absolute crystalline site on X : for a reference, see [Sta20, TAG 07I5]. (We implicitly take the crystalline site with respect to $W(k)$ without further comment; in other words, in the formulation of the Stacks Project, $S = \text{Spec } W(k)$ with the canonical divided power structure.) Let $\text{Crys}(X)$ be the category of crystals in *finite locally free $\mathcal{O}_{X/W(k)}$ -modules*. To make this more concrete, we introduce the following notation. A PD test object is a triple

$$(R, I, (\gamma_i))$$

where R is a $W(k)$ algebra with I a nilpotent ideal such that $\text{Spec } R/I$ “is” a Zariski open of X , and (γ_i) is a PD structure on I . Then a crystal in finite locally free modules M on X is a rule with input a PD test object $(R, I, (\gamma_i))$ and output a finitely generated projective R module

$$M_R$$

that is functorial: the pullback maps with respect to morphisms of PD test-objects are isomorphisms. In this formulation, the crystalline structure sheaf $\mathcal{O}_{X/W(k)}$ has the following description: on input $(R, I, (\gamma_i))$, the sheaf $\mathcal{O}_{X/W(k)}$ has as output the ring R .

By functoriality of the crystalline topos, the absolute Frobenius $Frob: X \rightarrow X$ gives a functor $Frob^*: \text{Crys}(X) \rightarrow \text{Crys}(X)$.

Definition 3.5. [Sta20, TAG 07N0] A (non-degenerate) F -crystal on X is a pair (M, F) where M is a crystal in finite locally free modules over the crystalline site of X and $F: Frob^* M \rightarrow M$ is an injective map of crystals.

We denote the category of F -crystals by $FC(X)$; it is a \mathbb{Z}_p -linear category with an internal \otimes but without internal homs or duals in general. There is a object $\mathbb{Z}_p(-1)$, given by the pair $(\mathcal{O}_{X/W(k)}, p)$. We denote by $\mathbb{Z}_p(-n)$ the n th tensor power of $\mathbb{Z}_p(-1)$.

Notation 3.6. [Ked16, Definition 2.1] Let X/k be a scheme of finite type over a perfect field. We denote by $\mathbf{F-Isoc}(X)$ the category of (convergent) F -isocrystals on X .

$\mathbf{F-Isoc}(X)$ is a \mathbb{Q}_p -linear Tannakian category. Denote by $\mathbb{Q}_p(-n)$ the image of $\mathbb{Z}_p(-n)$ and by $\mathbb{Q}_p(n)$ the dual of $\mathbb{Q}_p(-n)$. There is a notion of the rank of an F -isocrystal that satisfies that expected constraints given by \otimes and \oplus . Unfortunately, $\mathbf{F-Isoc}(X)$ is not simply the isogeny category of $FC(X)$; however, it is the isogeny category of the category of F -crystals in *coherent modules* (rather than finite, locally free modules). There is a natural functor $FC(X) \rightarrow \mathbf{F-Isoc}(X)$ [Ked16, 2.2].

For general smooth X/k over a perfect field, it seems that there are two p -adic analogs of a lisse l -adic sheaf: a (convergent) F -isocrystal and an *overconvergent F -isocrystal*. For our purposes (and following [Cre92]), overconvergent F -isocrystals are a better analog.

Notation 3.7. [Ked16, Definition 2.7] Let X/k be a smooth variety over a perfect field. We denote by $\mathbf{F}\text{-Isoc}^\dagger(X)$ the category of *overconvergent F -isocrystals on X* .

The category $\mathbf{F}\text{-Isoc}^\dagger(X)$ is a \mathbb{Q}_p -linear Tannakian category. There is a natural forgetful functor

$$\mathbf{F}\text{-Isoc}^\dagger(X) \rightarrow \mathbf{F}\text{-Isoc}(X)$$

which is fully faithful in general [Ked04, Theorem 1.1] and an equivalence of categories when X is proper.

Notation 3.8. For any finite extension L/\mathbb{Q}_p , we denote by $\mathbf{F}\text{-Isoc}^\dagger(X)_L$ the base-changed category a.k.a. *overconvergent F -isocrystals with coefficients in L* .

Definition 3.9. Let X/k be a smooth scheme over a perfect field. Let L be a p -adic local field and let \mathcal{E} be an object of $\mathbf{F}\text{-Isoc}^\dagger(X)_L$. We say \mathcal{E} is *effective* if the object $\text{Res}_{\mathbb{Q}_p}^L \mathcal{E}$ is in the essential image of the functor

$$FC(X) \rightarrow \mathbf{F}\text{-Isoc}^\dagger(X)$$

Being effective is equivalent to the existence of a ‘‘locally free lattice stable under F ’’. See Lemma 5.8 for the basic result on effectivity.

Let k be a perfect field. Then the category of F -crystals on $\text{Spec}(k)$ is equivalent to the category of finite free $W(k)$ -modules M together with a σ -linear injective map

$$F: M \rightarrow M.$$

(Morphisms in this category are defined to be $W(k)$ -linear maps that commute with the F s.) Similarly, the category of F -isocrystals on $\text{Spec}(k)$ is equivalent to the category of finite dimensional $K(k)$ -vector space V together with a σ -linear bijective map

$$F: V \rightarrow V,$$

where morphisms are defined in the obvious way. The rank of (V, F) is the rank of V as a vector space. Let L/\mathbb{Q}_p be a finite extension. Then $\mathbf{F}\text{-Isoc}(k)_L$ is equivalent to the following category: objects are pairs (V, F) where V is a finite free $K(k) \otimes_{\mathbb{Q}_p} L$ module and F is a $\sigma \otimes 1$ -linear bijective map

$$F: V \rightarrow V$$

and morphisms are maps of $K(k) \otimes_{\mathbb{Q}_p} L$ -modules that commute with F [Kri17, Proposition 5.12]. Note that $K(k) \otimes_{\mathbb{Q}_p} L$ is not necessarily a field. There is also a direct-sum decomposition of abelian categories:

$$\mathbf{F}\text{-Isoc}(k)_L \cong \bigoplus_{\lambda \in \mathbb{Q}_{\geq 0}} \mathbf{F}\text{-Isoc}(k)_L^\lambda$$

which is inherited from the analogous decomposition of $\mathbf{F}\text{-Isoc}(k)$. Here, $\mathbf{F}\text{-Isoc}(k)_L^\lambda$ is the (thick) abelian sub-category with objects isoclinic of slope λ .

Notation 3.10. We denote by $\mathbf{F}\text{-Isoc}^\dagger(X)_{\overline{\mathbb{Q}_p}}$ the 2-colimit of the base-changed categories over all finite extensions $\mathbb{Q}_p \subset L \subset \overline{\mathbb{Q}_p}$, via the functors $\text{Ind}_{\mathbb{Q}_p}^L$. This is the category of *overconvergent F -isocrystals on X with coefficients in $\overline{\mathbb{Q}_p}$* .

Remark 3.11. When k is a perfect field, $\mathbf{F}\text{-Isoc}(k)_{\overline{\mathbb{Q}_p}}$ has the following description. Objects are pairs (V, F) where V is a finite free $K(k) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$ -module and $F: V \rightarrow V$ is a bijective, $\sigma \otimes 1$ -linear map. Morphisms are $K(k) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$ -linear maps that commute with F .

Let L/\mathbb{Q}_p be an algebraic extension, let $k \cong \mathbb{F}_{p^d}$ and let (V, F) be an object of $\mathbf{F}\text{-Isoc}(k)_L$. Then F^d acts as a linear map on V . We let $P((V, F), t)$, the *characteristic polynomial of Frobenius*, be $\det(1 - (F^d)t|V)$. By [Kri17, Proposition 6.1]

$$P((V, F), t) \in L[t]$$

Definition 3.12. Let X/\mathbb{F}_q be a smooth variety, let L/\mathbb{Q}_p be an algebraic extension, and let $\mathcal{E} \in \mathbf{F}\text{-Isoc}^\dagger(X)_L$. Let $x \in X^\circ$ with residue field \mathbb{F}_{p^d} . Define $P_x(\mathcal{E}, t)$, the characteristic polynomial of \mathcal{E} at x , to be $\det(1 - (F^d)t|\mathcal{E}_x)$.

Notation 3.13. [Ked18, Notation 1.1.1] Let X/\mathbb{F}_q be a smooth connected variety. A *coefficient object* is an object either of $\mathbf{Weil}(X)_{\overline{\mathbb{Q}}_l}$ or of $\mathbf{F}\text{-Isoc}^\dagger(X)_{\overline{\mathbb{Q}}_p}$. We informally call the former the *étale case* and the latter the *crystalline case*.

Given an algebraic extension $\mathbb{Q}_l \subset K \subset \overline{\mathbb{Q}}_l$, objects of $\mathbf{Weil}(X)_K$ may be considered as étale coefficient objects. Similarly, given an algebraic extension $\mathbb{Q}_p \subset K \subset \overline{\mathbb{Q}}_p$, then objects of $\mathbf{F}\text{-Isoc}^\dagger(X)_K$ may be considered as crystalline coefficient objects via the induction functor.

Definition 3.14. Let \mathcal{E} be an étale (resp. crystalline) coefficient object. Let K be a subfield of $\overline{\mathbb{Q}}_l$ (resp. of $\overline{\mathbb{Q}}_p$) containing \mathbb{Q}_l (resp. containing \mathbb{Q}_p). The following three equivalent phrases

- \mathcal{E} has *coefficients in* K
- \mathcal{E} has *coefficient field* K
- \mathcal{E} is a *K -coefficient object*

mean that \mathcal{E} may be descended to $\mathbf{Weil}(X)_K$ (resp. $\mathbf{F}\text{-Isoc}^\dagger(X)_K$).

Note that \mathcal{E} having coefficient field L does not preclude \mathcal{E} from having coefficient field K for some sub-field $K \subset L$.

Definition 3.15. [Ked18, Definition 1.1.5] Let X/\mathbb{F}_q be a smooth variety and let \mathcal{F} be a coefficient object on X with coefficients in $\overline{\mathbb{Q}}_\lambda$. We say \mathcal{F} is *algebraic* if $P_x(\mathcal{F}, t) \in \overline{\mathbb{Q}}[t] \subset \overline{\mathbb{Q}}_\lambda$ for all $x \in X^\circ$. Let $E \subset \overline{\mathbb{Q}}_\lambda$ be a number field. We say \mathcal{F} is *E -algebraic* if $P_x(\mathcal{F}, t) \in E[t]$ for all $x \in X^\circ$.

Colloquially, semi-simple coefficient objects are determined by the characteristic polynomials of Frobenius at all closed points. In the étale case, this is a consequence of the Brauer-Nesbitt theorem and the Chebotarev density theorem. In the crystalline case, the argument is more subtle and is due to Tsuzuki [Abel18a, A.4.1]. See also a proof of this fact in [HP18] which is closer in spirit to the original proof and uses a p -adic variant of the Chebotarev density theorem.

Theorem 3.16. [Ked18, Theorem 3.3.1] *Let X/\mathbb{F}_q be a smooth variety. Let \mathcal{F} be a semi-simple coefficient object. Then \mathcal{F} is determined, up to isomorphism, by $P_x(\mathcal{F}, t)$ for all $x \in X^\circ$.*

4. COMPATIBLE SYSTEMS AND COMPANIONS

Definition 4.1. Let X/\mathbb{F}_q be a smooth variety. Let \mathcal{E} and \mathcal{E}' be algebraic coefficient objects on X with coefficients in $\overline{\mathbb{Q}}_\lambda$ and $\overline{\mathbb{Q}}_{\lambda'}$, respectively. Fix a field isomorphism $\iota: \overline{\mathbb{Q}}_\lambda \rightarrow \overline{\mathbb{Q}}_{\lambda'}$. We say \mathcal{E} and \mathcal{E}' are ι -*companions* if

$$\iota(P_x(\mathcal{E}, t)) = P_x(\mathcal{E}', t)$$

for all $x \in X^\circ$. We say that \mathcal{E} and \mathcal{E}' are companions if there exists an isomorphism $\iota: \overline{\mathbb{Q}}_\lambda \rightarrow \overline{\mathbb{Q}}_{\lambda'}$ that makes them ι -companions.

In other words, two coefficient objects are ι -companions if, under ι , the characteristic polynomials of Frobenius match up.

Remark 4.2. For convenience, we spell out the notion of companions for lisse l -adic sheaves. Let X/\mathbb{F}_q be a smooth variety and let L and L' be lisse $\overline{\mathbb{Q}}_l$ and $\overline{\mathbb{Q}}_{l'}$ sheaves respectively. (Here l may equal l' .) We say they are companions if there exists a field isomorphism $\iota: \overline{\mathbb{Q}}_l \rightarrow \overline{\mathbb{Q}}_{l'}$ such that for every closed point x of X , there is an equality of polynomials of Frobenius at x (see Definition 3.4):

$$\iota(P_x(L, t)) = P_x(L', t) \in \overline{\mathbb{Q}}_{l'}[t],$$

and furthermore this polynomial is in $\overline{\mathbb{Q}}[t]$.

Remark 4.3. Note that the ι in Definition 4.1 and Remark 4.2 does not reference the topology of $\overline{\mathbb{Q}}_p$ or $\overline{\mathbb{Q}}_l$. In particular, ι need not be continuous and in fact cannot be continuous if $l \neq p$.

Definition 4.4. Let X/\mathbb{F}_q be a smooth variety and let E be a number field. Then an E -compatible system is a system of lisse E_λ -sheaves $(\mathcal{F}_\lambda)_{\lambda \nmid p}$ over primes $\lambda \nmid p$ of E such that for every $x \in X^\circ$

$$P_x(\mathcal{F}_\lambda, t) \in E[t] \subset E_\lambda[t]$$

and this polynomial is independent of λ .

Definition 4.5. Let X/\mathbb{F}_q be a smooth variety and let λ be a rational prime. Let $(\mathcal{E}_v)_{v \in \Lambda}$ be a collection of $\overline{\mathbb{Q}}_\lambda$ coefficient objects on X , indexed by a finite set Λ . We say they form a *complete set of λ -adic companions* if

- for every $v \in \Lambda$ and for every $\iota \in \text{Aut}_{\mathbb{Q}}(\overline{\mathbb{Q}}_\lambda)$, there exists $v' \in \Lambda$ such that \mathcal{E}_v and $\mathcal{E}_{v'}$ are ι -companions, and
- For every $v, v' \in \Lambda$, \mathcal{E}_v and $\mathcal{E}_{v'}$ are companions.

The following definition of a complete E -compatible system involves *all possible* ι -companions.

Definition 4.6. Let X/\mathbb{F}_q be a smooth variety and let E be a number field. A *complete E -compatible system* (\mathcal{F}_λ) is an E -compatible system together with, for each prime λ of E over p , an object

$$\mathcal{F}_\lambda \in \mathbf{F}\text{-Isoc}^\dagger(X)_{E_\lambda}$$

such that the following two conditions hold.

- (1) For every place λ of E and every $x \in X^\circ$, the polynomial $P_x(\mathcal{F}_\lambda, t) \in E[t] \subset E_\lambda[t]$ is independent of λ .
- (2) For a prime r of \mathbb{Q} , let Λ denote the set of primes of E above r . Then $(\mathcal{F}_\lambda)_{\lambda \in \Lambda}$ form a complete set of r -adic companions for every r .

In particular, let (\mathcal{F}_λ) be a complete E -compatible system and let L be a lisse $\overline{\mathbb{Q}}_l$ -adic sheaf such that L is isomorphic an object of (\mathcal{F}_λ) . Let $\iota: \overline{\mathbb{Q}}_l \rightarrow \overline{\mathbb{Q}}_{\lambda'}$. Then the ι -companion to L exists and is isomorphic to an object in my list (\mathcal{F}_λ) . The most basic example of an E -compatible system is the following.

Example 4.7. Let E/\mathbb{Q} be a totally real number field of degree g and let \mathcal{M} be a Hilbert modular variety parametrizing principally polarized abelian g -folds with multiplication by a given order $\mathcal{O} \subset E$ and some level structure. For most primes \mathfrak{p} of E , \mathcal{M} has a smooth integral canonical model $\tilde{\mathcal{M}}$ over $\mathcal{O}_{\mathfrak{p}}$; moreover there is a universal abelian scheme $\tilde{\mathcal{A}} \rightarrow \tilde{\mathcal{M}}$. Let $\pi: A \rightarrow M$ denote the special fiber of such a smooth canonical model together with the induced abelian scheme. Then $E \hookrightarrow \text{End}_M(A) \otimes \mathbb{Q}$ and in particular the local system $R^1\pi_*(\mathbb{Q}_l)$ admits an action by

$$E \otimes \mathbb{Q}_l \cong \prod_{v|l} E_v$$

Here v ranges over the primes of E over l . Let e_v denote the idempotent projecting $E \otimes \mathbb{Q}_l$ onto E_v . Then $L_v := e_v R^1\pi_*(\mathbb{Q}_l)$ is a rank 2 lisse \mathbb{Q}_l -sheaf with an action of E_v , in other words a rank 2 lisse E_v -sheaf. It follows from the techniques of [Shi67, 11.9, 11.10] that the L_v are all companions. In fact, $(L_v)_{v|l}$ is a complete set of l -adic companions. If $[E: \mathbb{Q}] > 1$ and l splits non-trivially in E , then these will in general be mutually non-isomorphic lisse l -adic sheaves. By ranging over all primes of \mathbb{Q} (using relative crystalline cohomology at p) we obtain a complete E -compatible system.

We recall a conjecture of Deligne from Weil II [Del80, Conjecture 1.2.10].

Conjecture 4.8. Let X/\mathbb{F}_{p^d} be a normal variety with a geometric point $\bar{x} \rightarrow X$. Let $l \neq p$ be a prime. Let \mathcal{L} be an absolutely irreducible l -adic local system with finite determinant on X . The choice of \bar{x} allows us to think of this as a representation $\rho_l: \pi_1(X, \bar{x}) \rightarrow GL(n, \overline{\mathbb{Q}}_l)$. Then

- (1) ρ_l is pure of weight 0.
- (2) There exists a number field E such that for all closed points x of X , the polynomial $P_x(\mathcal{L}, t)$ has all of its coefficients in E . In particular, the eigenvalues of $\rho_l(F_x)$ are all algebraic numbers.
- (3) For each place $\lambda \nmid p$, the roots α of $P_x(\mathcal{L}, t)$ are λ -adic units in \overline{E}_λ .

(4) For each $\lambda|p$, the λ -adic valuations of the roots α satisfy

$$\left| \frac{v(\alpha)}{v(Nx)} \right| \leq \frac{n}{2}$$

where Nx is the size of the residue field of x .

(5) After possibly replacing E by a finite extension, for each $\lambda \nmid p$ there exists a λ -adic local system $\rho_\lambda : \pi_1(X, \bar{x}) \rightarrow GL(n, E_\lambda)$ that is compatible with ρ_l .

(6) After possibly replacing E by a finite extension, for each $\lambda|p$, there exists a crystalline companion to ρ_l .

The following conjecture may be seen as a refinement to (2), (5), and (6) of Conjecture 4.8.

Conjecture 4.9. (Companions) *Let X/\mathbb{F}_q be a smooth variety. Let \mathcal{F} be an irreducible coefficient object on X with algebraic determinant. Then there exists a number field E such that \mathcal{F} fits into a complete E -compatible system.*

The companions conjecture is surprising for the following reason: an l -adic local system is simply a continuous homomorphism from $\pi_1(X)$ to $GL_n(\overline{\mathbb{Q}}_l)$, and the topologies on \mathbb{Q}_l and $\mathbb{Q}_{l'}$ are completely different. Deligne likely made his conjecture out of the hope that such local systems were of *geometric origin*. We summarize what is known about Conjecture 4.8 and Conjecture 4.9.

By work of Deligne, Drinfeld, and Lafforgue, if X is a curve all such local systems are of geometric origin (in the sense of subquotients). Moreover, in this case Chin has proved Part 5 of the conjecture [Chi04, Theorem 4.1]. Abe has recently constructed a sufficiently robust theory of p -adic cohomology to prove a p -adic Langlands correspondence and hence answer affirmatively part 6 of Deligne's conjecture when X is a curve [Abe18a, Abe18b].

Theorem 4.10. (Abe, Lafforgue) *Let C/\mathbb{F}_q be a smooth curve. Then both Deligne's conjecture and the companions conjecture are true for C .*

In higher dimension, much is known but Conjecture 4.9 remains open in general. See [EK12] for a precise chronology of the following theorem, due to Deligne and Drinfeld.

Theorem 4.11. [Dell12, Dri12] *Let X/\mathbb{F}_q be a smooth variety. Let $l \neq p$ be a prime. Let \mathcal{L} be an absolutely irreducible l -adic local system with finite determinant on X . Then (1), (2), (3), and (5) of Conjecture 4.8 are true.*

Theorem 4.12. (Deligne, Drinfeld, Abe-Esnault, Kedlaya) *Let X/\mathbb{F}_q be a smooth variety and let \mathcal{F} be a coefficient object that is absolutely irreducible and has finite determinant. Then for any $l \neq p$, all l -adic companions exist.*

Proof. This follows from Theorem 4.11 together with either [AE19, Theorem 4.2] or [Ked18, Theorem 0.4.1]. \square

Remark 4.13. The “ p -companions” part of the conjecture is *not known* if $\dim C > 1$. Given an irreducible lisse $\overline{\mathbb{Q}}_l$ sheaf L with trivial determinant and a (necessarily non-continuous) isomorphism $\iota: \overline{\mathbb{Q}}_l \rightarrow \overline{\mathbb{Q}}_p$, it is **completely unknown** how to associate a crystalline ι -companion to L .

We remark that Part (4) of the Conjecture 4.8 is not tight even for $n = 2$.

Theorem 4.14. (Abe-Lafforgue) *Let X/\mathbb{F}_q be a smooth variety and let \mathcal{E} be an absolutely irreducible rank 2 coefficient object on C with finite determinant. Then for all $x \in C^\circ$, the eigenvalues α of F_x satisfy*

$$\left| \frac{v(\alpha)}{v(Nx)} \right| \leq \frac{1}{2}$$

for any p -adic valuation v .

Proof. Suppose X is a curve. Then when \mathcal{E} is an l -adic coefficient object, this is [Laf02, Theorem VII.6.(iv)]. For a crystalline coefficient object, we simply apply Theorem 4.10 to construct an l -adic coefficient object.

More generally, [Dri12, Proposition 2.17] and [AE19, Theorem 0.1] function as “Lefschetz theorems” (see [Esn17]) and allow us to reduce to the case of curves. \square

Remark 4.15. A refined version of part (4) of Conjecture 4.8 has been resolved for curves by V. Lafforgue [Laf11, Corollaire 2.2] using automorphic forms. A p -adic variant (which avoids automorphic techniques) was shown directly by Drinfeld-Kedlaya [DK17, Theorem 1.1.5]. They prove that for an indecomposable crystalline coefficient object, the “generic consecutive slopes do not differ by more than 1”, reminiscent of Griffith’s transversality. Kramer-Miller has recently given another proof of the p -adic variant [KM19].

Finally, we make the following simple observation: under strong assumptions on the field of traces, the existence of companions with the same residue characteristic is automatic.

Corollary 4.16. *Let X/\mathbb{F}_q be a smooth variety, E a number field, and \mathcal{F} be an irreducible E -algebraic coefficient object with coefficients in $\overline{\mathbb{Q}}_p$ (resp. $\overline{\mathbb{Q}}_l$). If there is only one prime in E lying above p (resp. l), then all p -adic (resp. l -adic) companions exist.*

Remark. The hypothesis of Corollary 4.16 are certainly satisfied if p (resp. l) is either totally inert or totally ramified in E .

Proof. We assume \mathcal{F} is crystalline; the argument in the étale case will be precisely analogous. Suppose $\mathcal{F} \in \mathbf{F}\text{-Isoc}^\dagger(X)_K$ where K/\mathbb{Q}_p is a p -adic local field; we might as well suppose K is Galois over \mathbb{Q}_p . As there is a unique prime \mathfrak{p} above p in E , there is a unique p -adic completion of E . We now explicitly construct all p -adic companions. For every $g \in \text{Gal}(K/\mathbb{Q}_p)$, consider the object ${}^g\mathcal{F}$: in terms of Definition 3.2, if $\mathcal{F} = (M, f)$, then ${}^g\mathcal{F} := (M, f \circ g^{-1})$. We claim this yields all p -adic companions.

Fix embeddings $E \hookrightarrow K \hookrightarrow \overline{\mathbb{Q}}_p$. Then there is a natural map

$$\text{Gal}(K/\mathbb{Q}_p) \rightarrow \text{Hom}_{\mathbb{Q}\text{-alg}}(E, \overline{\mathbb{Q}}_p)$$

given by precomposing with the fixed embedding. As there is a single \mathfrak{p} over p , this map is surjective. The result follows from Theorem 3.16, and the fact that $P_x({}^g\mathcal{E}, t) = {}^gP_x(\mathcal{E}, t)$ [Kri17, Proposition 6.16]. \square

Remark 4.17. In the crystalline setting of Corollary 4.16, the Newton polygons of all p -adic companions of \mathcal{F} are the same at all closed points. This does not contradict the example [Kos17, Example 2.2]; in his example, p splits completely in the reflex field of the Hilbert modular variety, and the reflex field is the same as the field generated by the characteristic polynomial of Frobenius elements over all closed points.

5. BARSOTTI-TATE GROUPS AND DIEUDONNÉ CRYSTALS

In this section, again let X/k denote a finite type scheme over a perfect field.

Definition 5.1. A *Dieudonné crystal* on X is a triple (M, F, V) where (M, F) is an F -crystal (in finite locally free modules) on X and $V : M \rightarrow \text{Frob}^*M$ is a map of crystals such that $V \circ F = p$ and $F \circ V = p$. We denote the category of Dieudonné Crystals on X by $DC(X)$.

Remark 5.2. Let U/k be a locally complete intersection morphism over a perfect field k . Then the natural forgetful functor $DC(U) \rightarrow FC(U)$ is fully faithful. This follows from the fact that $H_{\text{cris}}^0(U)$ is p -torsion free. This in turn follows from the fact that if $U = \text{Spec}(A)$ is affine, then the p -adic completion of a PD envelope of A is $W(k)$ -flat [dJM99, Lemma 4.7].

Notation 5.3. [CCO14, 1.4.1.3] Let $BT(X)$ denote the category of Barsotti-Tate (“ p -divisible”) groups on X .

For a thorough introduction to p -divisible groups and their contravariant Dieudonné theory, see [Gro74, BBM82, CCO14]. In particular, there exist Dieudonné functors: given a BT group \mathcal{G}/X one can construct a Dieudonné crystal $\mathbb{D}(\mathcal{G})$ over X . According to our conventions, \mathbb{D} is the *contravariant* Dieudonné functor. Moreover, given a p -divisible group \mathcal{G} on X , there exists the (Serre) dual, which we

denote by \mathcal{G}^t (see [CO09, 10.7] or [CCO14, 1.4.1.3]), whose constituent parts are obtained via Cartier duality of finite, locally free group schemes.

Definition 5.4. [CCO14, 3.3] Let \mathcal{G} and \mathcal{H} be Barsotti-Tate groups on X with the same height. Then an *isogeny* is a homomorphism

$$\phi: \mathcal{G} \rightarrow \mathcal{H}$$

whose kernel is represented by a finite, locally free group scheme over X .

Definition 5.5. [CCO14, 1.4.3] Let \mathcal{G} be a Barsotti-Tate group on X . A *quasi-polarization* is an isogeny

$$\phi: \mathcal{G} \rightarrow \mathcal{G}^t$$

that is skew-symmetric in the sense that $\phi^t = -\phi$ under the natural isomorphism $\mathcal{G} \rightarrow (\mathcal{G}^t)^t$.

Notation 5.6. Let \mathcal{G} be a Barsotti-Tate group on X . We denote by $QPol(\mathcal{G})$ the set of quasi-polarizations of \mathcal{G} .

The next theorem is a corollary of [dJ95, Main Theorem 1].

Theorem 5.7. (*de Jong*) Let X/k be a smooth scheme over a perfect field of characteristic p . Then the category $BT(X)$ is anti-equivalent to $DC(X)$ via \mathbb{D} .

The following lemma partially explains the relationship between slopes and effectivity.

Lemma 5.8. Let X/k be a smooth variety over a perfect field of characteristic p and let \mathcal{E} be an F -isocrystal on X . Then

- all of the slopes of \mathcal{E} at all $x \in |X|$ are non-negative if and only if there exists a dense Zariski open $U \subset X$, with complement of codimension at least 2, such that $\mathcal{E}|_U$ is effective (i.e., there exists an F -crystal (M, F) in finite, locally free modules on U such that $(M, F) \otimes_{\mathbb{Q}_p} \cong \mathcal{E}_U$).
- Furthermore, all of the slopes of \mathcal{E} at all points $x \in |X|$ are between 0 and 1 if and only if there exists a dense Zariski open $U' \subset X$, with complement of codimension at least 2, such that $\mathcal{E}_{U'}$ comes from a Dieudonné crystal.

Proof. The “only if” of both statements follows from the following facts.

- (1) If \mathcal{E} is an F -isocrystal on X , then the Newton polygon is an upper semi-continuous function with locally constant right endpoint (a theorem of Grothendieck, see e.g. [Kat79, Theorem 2.3.1]).
- (2) The locus of points where the slope polygon does not coincide with its generic value is pure of codimension 1 [Ked16, Theorem 3.12 (b)].

This “if” of the first statement is implicit in [Kat79, Theorem 2.6.1]; see also [Ked16, Remark 2.3].

Finally, let (M, F) be an F -crystal in finite, locally free modules on U with all Newton slopes ≤ 1 . We explain the small modification to the proof of [Kat79, Theorem 2.6.1] to find an open set $U' \subset U$, with complementary codimension at least 2, and an isogenous F -crystal (M', F) (in finite, locally free modules) on U' that underlies a Dieudonné crystal.

Set $V := F^{-1} \circ p$. Then V does not necessarily stabilize the lattice M . However, the pair (M, V) has the structure of $\sigma^{-1}F$ -isocrystal in the language of [Kat79, p. 115].⁴ Consider the following (à priori infinite) sum, obviously stabilized by V :

$$\tilde{M} := \sum_{n \geq 0} V^n M \subset \mathcal{E}.$$

Then [Kat79, Page 152] implies that \tilde{M} is crystal in *coherent* modules on U . It follows from the fact that $FV = VF = p$ that \tilde{M} is stabilized by F . Running the rest of the argument of [Kat79, Theorem 2.6.1], we obtain an open $U' \subset U$ of complementary codimension at least 2 and a crystal $\mathcal{M}' \subset \mathcal{E}_{U'}$ in

⁴Fortunately, every result in Katz’s paper is written for $\sigma^a F$ -crystals for all $a \neq 0$, not just the positive ones!

finite, locally free modules on U' that is stabilized by F and V .⁵ Therefore, the triple (\mathcal{M}', F, V) is a Dieudonné crystal on U' as desired. \square

Remark 5.9. The F -crystal constructed in Lemma 5.8 is *not unique*.

Remark 5.10. It is *not true* that if (M, F) is an F -crystal with all Newton slopes in $[0, 1]$, then (M, F) underlies a Dieudonné crystal (equivalently, it is not true that (M, F) has all Hodge slopes 0 or 1). Here is an example of an F -crystal over \mathbb{F}_p with Newton slopes all 1 but Hodge slopes 0 and 2: $M = \mathbb{Z}_p^2$, and F is given by the following matrix:

$$F := \begin{pmatrix} 0 & p^2 \\ 1 & 0 \end{pmatrix}$$

The following notion of *duality* of Dieudonné crystals is designed to be compatible with (Serre) duality of BT groups. In particular, the category $DC(X)$ admits a natural anti-involution.

Definition 5.11. [CO09, 2.10] Let $\mathcal{M} = (M, F, V)$ be a Dieudonné crystal on X . Then the *dual* \mathcal{M}^\vee is a Dieudonné crystal on X whose underlying crystal is defined on divided power test rings $(R, I, (\gamma_i))$ as

$$\mathcal{M}^\vee(R, I, (\gamma_i)) = \text{Hom}_R(M(R), R)$$

The operators F and V are defined as follows: if $(R, I, (\gamma_i))$ is a PD test object and $f \in \text{Hom}_R(M(R), R)$, then

$$F(f)(m) = f(Vm)$$

$$V(f)(m) = f(Fm)$$

It is an easy exercise to check that these rules render \mathcal{M}^\vee a Dieudonné crystal.

Remark 5.12. If $\mathcal{G} \rightarrow S$ is a BT group, then $\mathbb{D}(\mathcal{G})^\vee \cong \mathbb{D}(\mathcal{G}^t)$. [BBM82, 5.3.3.1].

We record the following lemma, which is surely well-known, as we could not find a reference. It compares the dual of a Dieudonné crystal with the dual of the associated F -isocrystal.

Lemma 5.13. *Let $\mathcal{M} = (M, F, V)$ be a Dieudonné crystal on X . Then the following two F -isocrystals on X are naturally isomorphic:*

$$\mathcal{M}^\vee \otimes \mathbb{Q} \cong (\mathcal{M} \otimes \mathbb{Q})^*(-1)$$

Proof. There is a natural perfect pairing to the Lefschetz F -crystal

$$\mathcal{M} \otimes \mathcal{M}^\vee \rightarrow \mathbb{Z}_p(-1)$$

as $FV = p$. There is also a perfect pairing

$$(\mathcal{M} \otimes \mathbb{Q}) \otimes (\mathcal{M} \otimes \mathbb{Q})^* \rightarrow \mathbb{Q}_p$$

to the “constant” F -isocrystal. Combining these two pairings shows the result. \square

6. A QUESTION OF GROTHENDIECK

In this section, we answer the question posed in [Gro66, 4.9], see Corollary 6.8. The argument is indeed similar to the arguments of [Gro66, Section 4]; the primary new input is [dJ98, 2.5]. However, some complications arise due to the fact that reduced schemes are not necessarily geometrically reduced over imperfect ground fields.

First, we record the following lemma, which is uniqueness of analytic continuation.

Lemma 6.1. *Let X be an integral noetherian scheme and let Y be a separated scheme. Let $Z \subsetneq X$ be a non-empty closed subscheme. Let $u, v: X \rightarrow Y$ be two morphisms that agree when restricted to the formal scheme $X_{/Z}$. Then $u = v$.*

⁵The argument amounts to taking the double dual, which is automatically a reflexive sheaf, and noting that a reflexive sheaf on a regular scheme is locally free away from a closed subset of codimension at least 3. See also the following remark [Kat79, Page 154].

Proof. Denote by Z_n the n^{th} formal neighborhood of Z . Then the data of the map $X_{/Z} \rightarrow X$ is equivalent to the data of the maps $Z_n \rightarrow X$ [Sta20, Tag 0AI6]. In other words, it suffices to prove that the set of closed immersions $\{Z_n \rightarrow X\}_{n \geq 0}$ are jointly schematically dense. By [GD66, 11.10.1(d)]: it suffices to prove the corresponding statement for every open affine $\text{Spec}(R) \subset X$. Let I be the ideal corresponding to $Z|_{\text{Spec}(R)}$. Then R is a (noetherian) domain. Krull's intersection lemma yields that the map of rings

$$R \rightarrow \varinjlim_n R/I^n$$

is injective and [GD66, 11.10.1(a)] then implies the desired schematic density. \square

We recall a well-known theorem of de Jong and Tate.

Theorem 6.2. (*de Jong-Tate*) *Let X be a noetherian normal integral scheme with generic point η . Let G and H be p -divisible groups over X . Then the natural map*

$$\text{Hom}_X(G, H) \rightarrow \text{Hom}_\eta(G_\eta, H_\eta)$$

is an isomorphism.

Proof. We immediately reduce to the case where $X \cong \text{Spec}(R)$. Then, as R is normal

$$R = \bigcap_{\mathfrak{p}} R_{\mathfrak{p}}$$

where \mathfrak{p} runs over all of the height 1 primes. On the other hand, as R is normal, if \mathfrak{p} is a height 1 prime, then $R_{\mathfrak{p}}$ is a discrete valuation ring. We are therefore reduced to proving the lemma over a discrete valuation ring, which results from [dJ98, Corollary 1.2] when the generic fiber has characteristic p and [Tat67, Theorem 4] when the generic fiber has characteristic 0. \square

Definition 6.3. Let K/k be a field extension. We say that it is *primary* if k is separably closed in K and *regular* if k is algebraically closed in K .

The following proposition directly imitates Conrad's descent-theoretic proof of a theorem of Chow [Con06, Theorem 3.19].

Proposition 6.4. *Let K/k be a primary extension of fields. Let G and H be BT groups on k . Then the natural injective map*

$$\text{Hom}_k(G, H) \rightarrow \text{Hom}_K(G_K, H_K)$$

is an isomorphism.

Proof. As $\text{Hom}_K(G_K, H_K)$ is a finite free \mathbb{Z}_p -module, we immediately reduce to the case that K/k is finitely generated. Let $K' \cong K \otimes_k K$. Then we have a diagram

$$k \rightarrow K \rightrightarrows K'$$

where the double arrows refer to the maps $K \rightarrow K \otimes_k K$ given by $p_1: \lambda \mapsto \lambda \otimes 1$ and $p_2: \lambda \mapsto 1 \otimes \lambda$ respectively. Let $F \in \text{Hom}_K(G_K, H_K)$. By Grothendieck's faithfully flat descent theory, to show that F is in the image of $\text{Hom}_k(G, H)$, it is enough to prove that $p_1^*F = p_2^*F$ under the canonical identifications $p_1^*G \cong p_2^*G$ and $p_1^*H \cong p_2^*H$ [Con06, Theorem 3.1]. There is a distinguished "diagonal point": $\Delta: K' \rightarrow K$, and restricting to the "diagonal point" the two pullbacks agree $F = \Delta^*p_1^*F = \Delta^*p_2^*F$. If we prove that the natural map

$$\text{Hom}_{K'}(G_{K'}, H_{K'}) \xrightarrow{\Delta^*} \text{Hom}_K(G_K, H_K)$$

is injective, we would be done. As K/k is primary and finitely generated, it suffices to treat the following two cases.

Case 1. Suppose K/k is a finite, purely inseparable extension of characteristic p . Then K' is an Artin local ring. Moreover, the ideal $I = \ker(\Delta)$ is nilpotent. As K' is a ring which is killed by p and I is nilpotent, the desired injectivity follows from a lemma of Drinfeld [Kat81, Lemma 1.1.3(2)] in his proof of the Serre-Tate theorem. Alternatively, it directly follows from [CCO14, 1.4.5.7].

Case 2. Suppose K/k is a finitely generated regular extension. Then K' is a noetherian integral domain. The desired injectivity follows from localizing and applying [CCO14, 1.4.4.3]. \square

We now record a p -adic variant of [Gro66, Proposition 4.4].

Lemma 6.5. *Let X/k be a normal, geometrically connected scheme of finite type over a field of characteristic p and let $\pi: A \rightarrow X$ be an abelian scheme. Then the following are equivalent*

- (1) *The BT group $A[p^\infty]$ is isomorphic to the pullback of a BT group G_0/k .*
- (2) *There exists an abelian scheme B_0/k such that A is isomorphic to the pullback of B_0 .*

Proof. That (2) implies (1) is evident, so we prove the reverse implication. We will explain a series of reductions. First of all, as Grothendieck observes in the beginning of the proof of [Gro66, Proposition 4.4], it follows from [Gro66, Proposition 1.2] that it is equivalent to prove the lemma over the generic point $\eta = \text{Spec}(k(X))$.

The hypotheses on X/k imply that $k(X)/k$ is a finitely generated primary extension. The obvious candidate for B_0 is then the Chow $k(X)/k$ trace, denoted by $\text{Tr}_{k(X)/k}(A_\eta)$. This exists, and if (2) holds, this works and is unique up to unique isomorphism by [Con06, Theorem 6.2, Theorem 6.4(2)]. Set $B_0 = \text{Tr}_{k(X)/k} A_\eta$; then B_0 is an abelian variety over k equipped with a homomorphism $B_{0,\eta} \rightarrow A_\eta$ with finite kernel. We will use functorial properties of the Chow trace to proceed with our reductions.

Let k' be the algebraic closure of k in $k(X)$. Then k'/k is a finite, purely inseparable extension. If we prove the theorem for the geometrically integral scheme X/k' , then by the following Proposition 6.6 we will have proven the theorem for X/k . Therefore we reduce to the case that X/k is geometrically integral.

We now explain why we may reduce to the case $k = \bar{k}$. As X/k is now supposed to be geometrically integral, $k(X)/k$ is a regular field extension and [Con06, Theorem 5.4] implies that the construction of $k(X)/k$ trace commutes with arbitrary field extensions E/k . On the other hand, the $k(X)$ -morphism $B_{0,\eta} \rightarrow A_\eta$ is an isomorphism if and only if it is when base-changed to $\bar{k}(X)$.

Next we explain why we may reduce to the case that $A[l] \rightarrow X$ is isomorphic to a trivial étale cover (maintaining the assumption that $k = \bar{k}$). Let $Y \rightarrow X$ be the finite, connected Galois cover that trivializes $A[l]$. Suppose there exists an abelian variety C_0/k such that $A_Y \cong C_{0,Y}$ as abelian schemes over Y . Let $I := \mathbf{Isom}(C_{0,X}, A)$ be the X -scheme of isomorphisms between $C_{0,X}$ and A . Our assumptions that $I(Y)$ is non-empty and that there is an isomorphism $A[p^\infty] \cong G_{0,X}$; our goal is to prove that $I(X)$ is non-empty.

Pick isomorphisms $C_0[p^\infty] \cong G_0$ and $A[p^\infty] \cong G_{0,X}$ once and for all. Then we have a commutative square of sets, where the horizontal arrows are injective because the p -divisible group of an abelian scheme is relatively schematically dense:

$$\begin{array}{ccc} I(X) & \longrightarrow & \text{Aut}_X(G_{0,X}) \\ \downarrow & & \downarrow \\ I(Y) & \longrightarrow & \text{Aut}_Y(G_{0,Y}). \end{array}$$

By Proposition 6.4, the maps $\text{Aut}_k(G_0) \rightarrow \text{Aut}_X(G_{0,X}) \rightarrow \text{Aut}_Y(G_{0,Y})$ are all isomorphisms: indeed, both $k(Y)/k$ and $k(X)/k$ are regular extensions. Let $H = \text{Gal}(Y/X)$. Then H acts naturally on $G_{0,Y}$ and hence acts on $I(Y) \hookrightarrow \text{Aut}_Y(G_{0,Y})$ via conjugation. On the other hand, the action of H on $\text{Aut}_Y(G_{0,Y})$ is trivial because the fixed points are $\text{Aut}_X(G_{0,X})$; therefore the action of H on $I(Y)$ is trivial. Our running assumption was that we had an element of $I(Y)$. Then Galois descent implies that $I(X)$ is non-empty, as desired.

We can find a polarization by [FC90, Remark 1.10(a)] (as X is a noetherian normal scheme), so assume we have one of degree d . Assume that $k = \bar{k}$ and that there exists an $l \geq 3$ such that $A[l] \rightarrow X$ is isomorphic to a trivial étale cover. Let $x \in X(k)$ be a k -point and let $C_0 := A_x$. Denote by $C := C_{0,X}$ be the ‘‘constant’’ abelian scheme with fiber C_0 . There are two induced moduli maps from X to the separated (fine) moduli scheme $\mathcal{A}_{g,d,l}$. Consider the restrictions of these moduli maps to

the formal scheme $X_{/x}$: Serre-Tate theory [Kat81, Theorem 1.2.1] ensures that the two formal moduli maps from $X_{/x}$ to $\mathcal{A}_{g,d,l}$ agree. Then by Lemma 6.1 the two moduli maps agree on the nose, whence the conclusion. \square

Proposition 6.6. *Let K/k be a finite, purely inseparable field extension. Let A/K be an abelian variety such that $A[p^\infty]$ is the pullback of a BT group G_0 over k . Then A is pulled back from an abelian variety over k .*

Proof. Let B be the Chow K/k -trace of A . By definition, there is a natural map $B_K \rightarrow A$ and by [Con06, Theorem 6.6] this map is an isogeny (as $(K \otimes_k K)_{red} \cong K$, see the Terminology and Notations of *ibid.*) with connected Cartier dual; in particular the kernel I has order a power of p . The induced isogeny of BT groups $B[p^\infty]_K \rightarrow G_{0,K} \cong A[p^\infty]$ also has kernel exactly I . Applying Proposition 6.4, we see that this isogeny descends to an isogeny $B[p^\infty] \rightarrow G_0$ over k . Call the kernel of this isogeny H ; then H is a finite k -group scheme with $H_K \cong I$. Finally, B/H is an abelian variety over k and the induced map

$$(B/H)_K \rightarrow A$$

is an isomorphism. (Indeed, after the fact H is trivial.) \square

We will use the following easy refinement.

Corollary 6.7. *Let X/k be a normal, geometrically connected scheme of finite type over a field of characteristic p . Let $p \nmid N$ and d be positive integers. Let $A \rightarrow X$ be an abelian scheme equipped with a polarization λ of degree d and a full level N structure Γ . Suppose $A[p^\infty]$ is the pullback of a BT group G_0 over k . Then (A, λ, Γ) is the pullback of a polarized abelian variety with full level- N structure $(A_0, \lambda_0, \Gamma_0)$ over k .*

Proof. By Proposition 6.5, there exists an abelian variety A_0 over k whose pullback to X is isomorphic to A . Let $K := k(X)$ and note again that K/k is a primary field extension. The group scheme $A_0[N]$ is étale, and a non-trivial étale group scheme cannot become trivial after a primary field extension; therefore the full level- N structure descends. We need to show the polarization descends to a polarization on A_0 . A theorem of Chow [Con06, Theorem 3.19] implies that any homomorphism $A_K \rightarrow A_K^t$ is the pullback of a unique homomorphism $\lambda_0: A_0 \rightarrow A_0^t$. There exists an ample line bundle \mathcal{L} on $A_{\bar{K}}$ such that λ comes from \mathcal{L} in the usual way. On the other hand, the Néron-Severi group of $A_{0,\bar{k}}$ and the Néron-Severi group of $A_{\bar{K}}$ are isomorphic; hence one can find such a line bundle \mathcal{L} (in the same Néron-Severi class) that is defined over \bar{k} . This implies that λ_0 , as a symmetric isogeny, is in fact a polarization as in [FC90, Ch. I, Definition 1.6]. \square

Corollary 6.8. *Let X be a locally noetherian normal scheme and $U \subset X$ be an open dense subset whose complement has characteristic p . Let $A_U \rightarrow U$ be an abelian scheme. Then A_U extends to an abelian scheme over X if and only if $A_U[p^\infty]$ extends to a p -divisible group over X .*

Proof. When X is the spectrum of a discrete valuation ring, this follows from Grothendieck's generalization of Néron-Ogg-Shafarevitch criterion in mixed characteristic [GRR06, Exposé IX, Theorem 5.10] and work of de Jong [dJ98, 2.5] in equal characteristic p . The rest of the argument closely follows [Gro66, pages 73-76].

The reductions in the proof on [Gro66, page 73] immediately bring us to the following case: X is integral, $A_U \rightarrow U$ has a polarization of degree d , and the l -torsion is isomorphic to the trivial étale cover of U . In particular, we have a map $f: U \rightarrow \mathcal{A}_{g,d,l}$, where the latter fine moduli space is a separated scheme of finite type over $\mathbb{Z}[1/l]$.

Let X' denote the closure of the graph of f . Then the valuative criterion of properness together with the case of discrete valuation rings discussed above imply that the morphism $X' \rightarrow X$ is proper. Let X'' be the normalization of X' . As we make no ‘‘Japanese’’ assumption, $X'' \rightarrow X'$ may not be finite; on the other hand, by a theorem of Nagata it has reduced and finite fibers [GD64, Chapitre 0, 23.2.6]. For every point x of X , let $Y := (X''_x)_{red}$ and $Z := (X'_x)_{red}$; then the natural map $Y \rightarrow Z$ has reduced and finite fibers by the preceding sentence. Let B' and B'' be the abelian schemes on X'

and X'' respectively, induced from the map $X'' \rightarrow X' \rightarrow \mathcal{A}_{g,d,l}$. Let $k := k(x)$ and note that Z/k is proper, reduced, and geometrically connected (only the last part is non-trivial: as X is a noetherian normal scheme and $X' \rightarrow X$ is a proper birational morphism, it follows from Zariski's main theorem). It suffices to prove that for every point x of X , the map $Z \rightarrow \mathcal{A}_{g,d,l}$ factors through $x = \text{Spec}(k)$:

$$Z \rightarrow x \rightarrow \mathcal{A}_{g,d,l}$$

Let G be the BT group that exists by assumption on X and G'' the pullback to X'' . Then $B''[p^\infty]$ and G'' are isomorphic on $U \subset X''$. By Theorem 6.2, $B''[p^\infty] \cong G''$. Therefore B_Y'' has constant (with respect to k) BT group over Y .

We may of course assume that $x \in X \setminus U$ and in particular that $\text{char}(k) = p$. To prove the desired statement about the map $Z \rightarrow \mathcal{A}_{g,d,l}$, we claim that we may replace Z by Z'_i , the irreducible components of a scheme Z' that is finite and dominant over Z , and k by $k_i = H^0(Z'_i, \mathcal{O}_{Z'_i})$. Indeed, as Z/k is proper, reduced, and geometrically connected, the ring $K := H^0(Z, \mathcal{O}_Z)$ is a finite, purely inseparable field extension of k . If we prove that each of the maps $Z'_i \rightarrow \mathcal{A}_{g,d,l}$ factors through $\text{Spec}(k_i)$, then it follows that the map $Z \rightarrow \mathcal{A}_{g,d,l}$ factors through $\text{Spec}(K)$. Applying Corollary 6.7 to $\text{Spec}(K)/k$, we deduce that $Z \rightarrow \mathcal{A}_{g,d,l}$ indeed factors through $\text{Spec}(k)$, as desired.

For every maximal point z_i of Z , let Z_i be the closure of z_i in Z (with reduced, induced structure), let y_i be a point of Y over z_i and Y_i the closure of y_i in Y (with induced reduced structure). Let Y'_i be the normalization of Y_i . As $k(y_i)$ is finite over $k(z_i)$ and Z/k is of finite type, it is a consequence of a theorem of E. Noether that $Y'_i \rightarrow Z_i$ is then finite [Eis95, Corollary 13.13]. We take $Z' := \sqcup Y'_i$.

We are therefore left to prove that the morphisms $Y'_i \rightarrow \mathcal{A}_{g,d,l}$ factor through $\text{Spec}(k_i)$, where $k_i = H^0(Y'_i, \mathcal{O}_{Y'_i})$. By construction, the Y'_i/k_i are normal, geometrically connected, of finite type, and the BT group of the associated abelian scheme on Y'_i is constant with respect to k_i . We may therefore directly apply Corollary 6.7. \square

7. LEFSCHETZ THEOREMS I

In this section, we collect together several Lefschetz theorems having to do with morphisms of abelian varieties and p -divisible groups. We first state the simplest case of a theorem of a Lefschetz theorem of Abe-Esnault, which will be useful for us to prove Lemma 7.2, a Lefschetz theorem for homomorphisms of BT groups.

Theorem 7.1. (*Abe-Esnault*) *Let X/k be a smooth projective variety over a perfect field k of characteristic p with $\dim X \geq 2$ and let $C \subset X$ be a smooth projective curve that is the complete intersection of smooth ample divisors. Then for any $\mathcal{E} \in \mathbf{F}\text{-Isoc}(X)_{\overline{\mathbb{Q}}_p}$, the following restriction map is an isomorphism*

$$H^0(X, \mathcal{E}) \rightarrow H^0(C, \mathcal{E})$$

Proof. Note that \mathcal{E} is automatically overconvergent as X is projective. This then follows immediately from the arguments of [AE19, Corollary 2.4]. We comment on two differences. First, while the statements [AE19, Corollary 2.4, Proposition 2.2] are only stated for \mathbb{F} , they only require k be perfect as stated in [AE19, Remark 2.5].⁶ \square

Lemma 7.2. *Let X/k be a smooth projective variety over a field k of characteristic p with $\dim X \geq 2$ and let $U \subset X$ be a Zariski open subset. Let \mathcal{G}_U and \mathcal{H}_U be BT groups on U . Let $C \subset U$ be a smooth projective curve that is the complete intersection of smooth ample divisors of X and denote by \mathcal{G}_C (resp. \mathcal{H}_C) the restriction of \mathcal{G}_U (resp. \mathcal{H}_U) to C . Then the following restriction map*

$$\text{Hom}(\mathcal{G}_U, \mathcal{H}_U) \rightarrow \text{Hom}_C(\mathcal{G}_C, \mathcal{H}_C)$$

is injective with cokernel killed by a power of p .

⁶Indeed, Abe and Esnault pointed out via email that the part of [AC13] they cite for these arguments is Section 1, which only requires k be perfect.

Proof. We first reduce to the case that k is perfect. Let k'/k be a field extension and let $U' := U \times_k k'$. By Theorem 6.2, the natural map

$$\mathrm{Hom}_{U'}(\mathcal{G}_{U'}, \mathcal{H}_{U'}) \rightarrow \mathrm{Hom}_{k'(U)}(\mathcal{G}_{k'(U)}, \mathcal{H}_{k'(U)})$$

is an isomorphism and similarly for D . Let l/k be the perfect closure; it is in particular a primary extension. Therefore $l(U)/k(U)$ and $l(D)/k(D)$ are also primary extensions and hence Proposition 6.4 implies both sides of the equation are unchanged when we replace k by l .

Both the left and the right hand side are torsion-free \mathbb{Z}_p -modules. We have the following diagram

$$\begin{array}{ccc} \mathrm{Hom}_U(\mathcal{G}_U, \mathcal{H}_U) & \xrightarrow{\sim} & \mathrm{Hom}_U(\mathbb{D}(\mathcal{H}_U), \mathbb{D}(\mathcal{G}_U)) \\ \mathrm{res} \downarrow & & \downarrow \otimes \mathbb{Q} \\ \mathrm{Hom}_C(\mathcal{G}_C, \mathcal{H}_C) & & \mathrm{Hom}_U(\mathbb{D}(\mathcal{H}_U) \otimes \mathbb{Q}, \mathbb{D}(\mathcal{G}_U) \otimes \mathbb{Q}) \\ \otimes \mathbb{Q} \downarrow & & \uparrow \wr \\ \mathrm{Hom}_C(\mathbb{D}(\mathcal{H}_C) \otimes \mathbb{Q}, \mathbb{D}(\mathcal{G}_C) \otimes \mathbb{Q}) & \longleftarrow & \mathrm{Hom}_X(\mathcal{F}, \mathcal{E}) \end{array}$$

We explain what \mathcal{E} and \mathcal{F} are. Consider the following diagram:

$$\mathbf{F}\text{-Isoc}^\dagger(X) \xrightarrow{\sim} \mathbf{F}\text{-Isoc}(X) \rightarrow \mathbf{F}\text{-Isoc}(U)$$

The first arrow is an equivalence of categories because X is proper and the second arrow an equivalence of categories by work of Kedlaya and Shiho [Ked16, Theorem 5.1] because the complement of U has codimension at least 2. Hence $\mathbb{D}(\mathcal{G}_U) \otimes \mathbb{Q}$ and $\mathbb{D}(\mathcal{H}_U) \otimes \mathbb{Q}$ have canonical extensions to X , which we denote by \mathcal{E} and \mathcal{F} respectively. The lower right hand vertical arrow is an isomorphism, again by the above equivalence of categories. Moreover we have the isomorphisms

$$\mathbb{D}(\mathcal{G}_C) \otimes \mathbb{Q} \cong \mathcal{E}|_C, \text{ and } \mathbb{D}(\mathcal{H}_C) \otimes \mathbb{Q} \cong \mathcal{F}_C$$

Our goal is to prove that res , an application of the “restrict to C ” functor, is injective with cokernel killed by a power of p . As $\mathrm{Hom}_{D_C}(\mathcal{G}_C, \mathcal{H}_C)$ is a finite free \mathbb{Z}_p -module, it is equivalent to prove that the induced map

$$\mathrm{Hom}_U(\mathcal{G}_U, \mathcal{H}_U) \otimes \mathbb{Q} \rightarrow \mathrm{Hom}_C(\mathbb{D}(\mathcal{H}_C) \otimes \mathbb{Q}, \mathbb{D}(\mathcal{G}_C) \otimes \mathbb{Q})$$

is an isomorphism of \mathbb{Q}_p -vector spaces. By chasing the diagram, this is equivalent to the bottom horizontal arrow being an isomorphism. This arrow is an isomorphism due to Theorem 7.1 (which, as stated, requires k to be perfect). \square

In a similar vein, we write down a Lefschetz theorem for homomorphisms of abelian schemes.

Theorem 7.3. *Let X/k be a smooth projective variety over a field k with $\dim X \geq 2$ and let $U \subset X$ be a Zariski open subset whose complement has codimension at least 2. Let $A_U \rightarrow U$ and $B_U \rightarrow U$ be abelian schemes over U . Let $C \subset U$ be a smooth projective curve that is the complete intersection of smooth, ample divisors of X , and denote by A_C (resp. B_C) the restriction of A_U (resp. B_U) to C . Then the natural restriction map*

$$(7.1) \quad \mathrm{Hom}_U(A_U, B_U) \rightarrow \mathrm{Hom}_C(A_C, B_C)$$

is an isomorphism when tensored with \mathbb{Q} . If the cokernel is non-zero, then $\mathrm{char}(k) = p$ and the cokernel is killed by a power of p .

Proof. We first remark that the above map is well-known to be injective and does not require any positivity property of D (one may immediately reduce to the case of a discrete valuation ring).

We now reduce to the case when k is a finitely generated field. Both sides of Equation 7.1 are finitely generated \mathbb{Z} -modules; hence we may replace k by a subfield finitely generated over the prime field over which everything in Equation 7.1 is defined *without changing either the LHS or the RHS*. Note further that by Grothendieck’s Lefschetz theorem [Gro05, Exposé X, §2, Corollaire 2.6], Zariski-Nagata purity, and induction, the map $\pi_1(C) \rightarrow \pi_1(U)$ is surjective.

By [FC90, Ch. I, Prop. 2.7], the natural map

$$(7.2) \quad \mathrm{Hom}(A_U, B_U) \rightarrow \mathrm{Hom}_k(A_{k(U)}, B_{k(U)})$$

is an isomorphism, and similarly for C . Now, as both sides of Equation 7.1 are finite free \mathbb{Z} -modules, it suffices to examine its completion at all finite places. There are two cases.

Case 1. $\mathrm{char}(k) = 0$. Then ‘‘Tate’s isogeny theorem’’ is true for U and C ; that is, the natural map

$$\mathrm{Hom}_U(A_U, B_U) \otimes \mathbb{Z}_l \rightarrow \mathrm{Hom}_{\pi_1(U)}(T_l(A_U), T_l(B_U))$$

is an isomorphism for every l , and similarly for C ; this is a combination of work of Faltings [FWG⁺92, Theorem 1, Page 211] and Equation 7.2. On the other hand, as noted the restriction map $\pi_1(C) \rightarrow \pi_1(U)$ is surjective; therefore the natural map

$$\mathrm{Hom}_{\pi_1(U)}(T_l(A_U), T_l(B_U)) \rightarrow \mathrm{Hom}_{\pi_1(C)}(T_l(A_C), T_l(B_C))$$

is an isomorphism. Moreover, this is true for all l . Therefore Equation 7.1 is an isomorphism.

Case 2. $\mathrm{char}(k) = p$. Then ‘‘Tate’s isogeny theorem’’, in this case a theorem of Tate-Zarhin-Mori [Mor85, Ch. XII, Théorème 2.5(i), p. 244] together with 7.2, implies that the argument of part (1) works as long as $l \neq p$. Therefore Equation 7.1 is rationally an isomorphism and the cokernel is killed by a power of p . □

Remark 7.4. Daniel Litt indicated to us that there is a more elementary proof of Theorem 7.3 (i.e. not using Faltings’ resolution of the Tate conjecture for divisors on abelian varieties) when $\mathrm{char}(F) = 0$ along the lines of his thesis. He also pointed out that, when $F \cong \mathbb{C}$, the result follows from the theorem of the fixed part.

Example 7.5. Litt constructed an example when Equation 7.1 is not an isomorphism in characteristic p . Let E/\mathbb{F} be a supersingular elliptic curve, so $\alpha_p \subset E$ is a subgroup scheme. Then $E \times E \times E$ contains α_p^3 as a subgroup scheme. There is an injective homomorphism

$$(\alpha_p)_{\mathbb{P}^2} \hookrightarrow (\alpha_p^3)_{\mathbb{P}^2}$$

given as follows: if $[x : y : z]$ are the coordinates on \mathbb{P}^2 and (α, β, γ) are linear coordinates on $\alpha_p^3 \subset \mathbb{A}^3$, then the image of α is cut out by the equations:

$$[x : y : z] = [\alpha : \beta : \gamma]$$

Let $A \rightarrow \mathbb{P}^2$ be the quotient of the constant $E \times E \times E$ family over \mathbb{P}^2 by this varying family of α_p . This family admits a principal polarization and the induced image

$$\mathbb{P}^2 \rightarrow \mathcal{A}_{3,1} \otimes \mathbb{F}$$

is infinite, see the text after [LO98, page 59, 9.4.16]. Suppose there exists a line $H \subset \mathbb{P}^2$, an automorphism ϕ of \mathbb{P}^2 that fixes H pointwise, and a point $p \in \mathbb{P}^2 \setminus H$ with the fibers A_p and $A_{\phi^{-1}(p)}$ non-isomorphic as unpolarized abelian threefolds. Let $B := \phi^*A$, a new family of abelian threefolds over \mathbb{P}^2 . Then $A|_H \cong B|_H$ but $A \not\cong B$ as abelian schemes over \mathbb{P}^2 .

We now explain why we can always find such a triple (H, ϕ, p) . It suffices to find two points $p, q \in \mathbb{P}^2$ such that $A_p \not\cong A_q$; we can then of course find an automorphism that sends p to q and fixes a line. For fixed $g \geq 2$, a theorem of Deligne implies that there is only one isomorphism class of superspecial abelian variety of dimension g over \mathbb{F} [Shi79, Theorem 3.5]. Now, any abelian variety has finitely many polarizations of any fixed degree [NN81, Theorem 1.1]; therefore we can find $p, q \in \mathbb{P}^2(\mathbb{F})$ such that A_p is superspecial and A_q is not, as desired.

The primary application of 7.2 is to show that certain quasi-polarizations lift.

Corollary 7.6. *Let X/k be a smooth projective variety over a field k of characteristic p with $\dim X \geq 2$, let $U \subset X$ be a Zariski open subset with complement of codimension at least 2, and let $C \subset U$ be a*

smooth curve that is the complete intersection of smooth ample divisors of X . Let \mathcal{G}_U be a BT group on U . Then the natural map

$$QPol(\mathcal{G}_U) \rightarrow QPol(\mathcal{G}_C)$$

is injective and moreover for any $\phi_C \in QPol(\mathcal{G}_C)$, there exists $n \geq 0$ such that $p^n \phi_C$ is the image of the above map.

Proof. We have a commutative square

$$\begin{array}{ccc} QPol(\mathcal{G}_U) & \longrightarrow & \text{Hom}_U(\mathcal{G}_U, \mathcal{G}_U^t) \\ \downarrow & & \downarrow \\ QPol(\mathcal{G}_C) & \longrightarrow & \text{Hom}_C(\mathcal{G}_C, \mathcal{G}_C^t) \end{array}$$

where the vertical arrow on the right is injective with torsion cokernel by Lemma 7.2. Now, $QPol(\mathcal{G}_C)$ are those isogenies in $\text{Hom}_C(\mathcal{G}_C, \mathcal{G}_C^t)$ that are anti-symmetric, i.e. those ϕ_C such that $\phi_C^t = -\phi_C$. Given $\phi_C \in QPol(\mathcal{G}_C)$, we know there exists an $n \geq 0$ such that $p^n \phi_C$ is the image of a unique $\psi \in \text{Hom}_U(\mathcal{G}_U, \mathcal{G}_U^t)$; to prove the corollary we must check that ψ is an isogeny and anti-symmetric on all of U .

First of all, we remark that the transpose map commutes with the vertical right hand arrow, “restriction to C ”. As the vertical right hand arrow is injective, the skew-symmetry of $p^n \phi_C$ implies that ψ is automatically skew-symmetric. Moreover, ψ is an isogeny on C ; hence ψ is an isogeny on a Zariski open set $U' \subset U$ by the paragraph after [CCO14, 3.3.9]. (All that is needed is that ψ is an isogeny at a single point $u \in U$.) By a theorem of de Jong [dJ98, 1.2], ψ is an isogeny everywhere because U is integral and normal.⁷ \square

Corollary 7.7. *Let X/k be a smooth projective variety over a field k of characteristic p , let $C \subset X$ be a smooth curve that is the complete intersection of smooth, ample divisors of X , and let $U \supset C$ be a Zariski open set of X . Let $A_C \rightarrow C$ be an abelian scheme and let \mathcal{G}_U be a BT group on U such that $\mathcal{G}_C \cong A_C[p^\infty]$ as BT groups on C . Then there exists a polarizable formal abelian scheme \hat{A} over the formal scheme $X_{/C}$ such that $\hat{A}[p^\infty] \cong \mathcal{G}_{|X_{/D}}$. The formal abelian scheme \hat{A} is (uniquely) algebraizable.*

Proof. First of all, C is normal; then by [FC90, page 6] $A_C \rightarrow C$ is projective. Let ϕ_C be a polarization of A_C ; abusing notation, we also let ϕ_C denote the induced quasi-polarization on the BT group \mathcal{G}_C . By Corollary 7.6, there exists an $n \geq 0$ such that the quasi-polarization $p^n \phi_C$ lifts to a quasi-polarization ψ of \mathcal{G}_U . Applying Serre-Tate [Kat81, Theorem 1.2.1], we therefore get a formal abelian scheme \hat{A} on $X_{/D}$; the fact that the quasi-polarization lifts implies that \hat{A} is polarizable and hence is (uniquely) algebraizable by Grothendieck’s algebraization theorem [Sta20, TAG 089a]. \square

8. LEFSCHETZ THEOREMS II

Good references for this section are [Gro05, Har70].

Definition 8.1. [Gro05, Exposé X, Section 2] We say a pair of a noetherian scheme X and a non-empty closed subscheme Z satisfies the property $Le\!f(X, Z)$ if for any vector bundle E on a Zariski neighborhood $U \supset Z$, the following restriction map is an isomorphism:

$$H^0(U, E) \cong H^0(X_{/Z}, E|_{X_{/Z}})$$

Definition 8.2. ([HM68, p.64] or [Har70, Ch V, §1, p. 190]) Let X be an integral noetherian scheme and $Z \subset X$ be a non-empty, closed subscheme. We say that Z is $\mathcal{G}\mathcal{I}$ in X if the natural embedding

$$K(X) \hookrightarrow K(X_{/Z})$$

from the ring of rational functions to the ring of formal rational functions is an isomorphism.

⁷See [CCO14, 3.3.10] for a thorough discussion of these issues.

The following theorem is an easy consequence of [Har70, Ch V].

Theorem 8.3. *Let X/k be a smooth projective variety. Let $C \subset X$ be a smooth curve which is the complete intersection of smooth, ample divisors. Let $\hat{f}: X_{/C} \rightarrow M$ be a morphism to a quasi-projective variety. Then there exists an open set $W \supset C$ together with a map $f: W \rightarrow M$ such that \hat{f} is the formal completion of f .*

Proof. It follows from [Har70, Ch V, Corollary 2.3, p. 202] that C is G3 in X . Recall that topologically, $X_{/C}$ is homeomorphic to C ; when we make set-theoretic arguments, we pass freely between $X_{/C}$ and C . Embed M in \mathbb{P}_k^m and $\hat{L} := \hat{f}^* \mathcal{O}_M(1)$ be the pullback of the ample class to $X_{/C}$. Then \hat{L} is globally generated. Pick regular sections $(\hat{s}_i)_{i=1}^m$ of $H^0(X_{/C}, \hat{L})$ that define the map to \mathbb{P}^m ; in particular, the \hat{s}_i globally generate. Let $V_i \subset X_{/C}$ be the locus on which $\hat{s}_i \neq 0$. Then \hat{s}_i yields a trivialization over V_i :

$$\hat{s}_i: \mathcal{O}_{X_{/C}}|_{V_i} \rightarrow \hat{L}_{V_i}.$$

As the s_i globally generate \hat{L} , it follows that $\cup_i V_i = X_{/C}$.

With respect to the trivializations given by the \hat{s}_i , the line bundle \hat{L} is defined by the transition functions $\hat{\phi}_{ij} := \frac{\hat{s}_i}{\hat{s}_j}$ on $V_{ij} := V_i \cap V_j$. By definition, $\hat{\phi}_{ij} \in K(X_{/C})$. By the G3 property, there is a unique meromorphic function $\phi_{ij} \in K(X)$ that restricts to $\hat{\phi}_{ij}$ on $X_{/C}$. Call D_{ij} the divisor of zeroes of ϕ_{ij} .

For each $i \in \{1, 2, \dots, m\}$, there exists an open set $U_i \subset X$ such that

- $U_i \cap X_{/C} = V_i$ and
- $U_i \cap D_{ij} = \emptyset$ for $j \neq i$.

Indeed, we can of course find an open set $U_i \subset X$ with $U_i \cap X_{/C} = V_i$. Replace U_i by the complement of $\cup_{j \neq i} D_{ij} \cap U_i$ in U_i ; in other words, remove all of the zeroes of the ϕ_{ij} from U_i for $j \neq i$. The \hat{s}_i are regular (and not merely meromorphic) sections; hence $D_{ij} \cap X_{/C}$ is a subset of the finite set of zeros of \hat{s}_i . Therefore $U_i \cap X_{/C} = V_i$. Set

$$W := \cup_i U_i \supset X_{/C}$$

and $U_{ij} := U_i \cap U_j$. Then on U_{ij} , the functions ϕ_{ij} are invertible. Moreover, the fact that $\hat{\phi}_{ij} \hat{\phi}_{jk} = \hat{\phi}_{ik}$ implies that $\phi_{ij} \phi_{jk} = \phi_{ik}$ by the injectivity of the map $K(X) \hookrightarrow K(X_{/C})$. Therefore we obtain a line bundle L_W on W algebraizing \hat{L} .

By *Lef*(X, C) [Har70, Ch. V, Prop 2.1, p. 200], we deduce that after shrinking W (while still containing C), we may ensure that $H^0(W, L_W) = H^0(X_{/C}, \hat{L})$. This implies that the \hat{s}_i algebraize to sections s_i of $H^0(W, L_W)$. Further replace W by the complement of the common locus of zeroes of the s_i .

Finally, we obtain a morphism $f: W \rightarrow \mathbb{P}^m$. As $f|_{X_{/C}}$ lands in M and X is integral, this implies that the image of g entirely lands in the Zariski closure \bar{M} of $M \subset \mathbb{P}^m$ by schematic density of $X_{/C}$ inside of X . By further shrinking W , we obtain that the map $\hat{f}: X_{/C} \rightarrow M$ uniquely algebraizes to a map $f: W \rightarrow M$, as desired. \square

Corollary 8.4. *Let X/k be a smooth projective variety and let $C \subset X$ be a smooth curve that is the complete intersection of smooth ample divisors. Let $\hat{A} \rightarrow X_{/C}$ be an polarizable abelian scheme over the formal completion of X along C . If there exists $l > 2$, prime to $\text{char}(k)$, such that the l -torsion of $\hat{A}|_C \rightarrow C$ is trivial, then there exists a Zariski open subset of $U \subset X$ containing C , and an abelian scheme $A_U \rightarrow U$ that extends $\hat{A} \rightarrow X_{/C}$.*

Proof. Note that by topological invariance of the étale site, if the l -torsion of $\hat{A}|_C$ is trivial, then so is the l -torsion of \hat{A} . If $l > 2$, then the moduli space $\mathcal{A}_{g,d,l}$ exists as a quasi-projective scheme over $\text{Spec}(\mathbb{Z}[1/l])$. Apply Theorem 8.3 to the induced map $X_{/C} \rightarrow \mathcal{A}_{g,d,l}$. \square

Corollary 8.4 is not true if one does not assume that the l -torsion is trivial for the same reason that Theorem 8.3 is not true with M a Deligne-Mumford stack. In particular, if $C \subset \mathbb{P}^2$ is a smooth plane

curve of positive genus, it is easy to see that there is a non-trivial map $\mathbb{P}^n_{/C} \rightarrow B\mathbb{Z}/2\mathbb{Z}$ that does not algebraize.

D. Litt explained the following lemma to the first author.

Lemma 8.5. *Let X/k be a smooth projective variety with $\dim(X) \geq 2$ and let $U \subset X$ be an open subset whose complement has codimension at least 2. Let $C \subset U$ be a smooth curve that is the complete intersection of smooth ample divisors of X . Let G_U and H_U be finite flat group schemes or Barsotti-Tate groups on U . Suppose the restrictions $G_{X/C} \cong H_{X/C}$. Then $G_U \cong H_U$.*

Proof. If we prove the lemma for finite flat group schemes, then the result for BT groups immediately follows: a homomorphism of BT groups is simply a compatible sequence of homomorphisms of n -truncated BT groups. Therefore we assume G_U and H_U are finite flat group schemes. We remark that the $\text{Lef}(X, C)$ is satisfied by [Har70, Ch. V, Proposition 2.1]. By definition, this means for any vector bundle E on a Zariski neighborhood $V \supset D$, the natural restriction map on sections

$$H^0(V, E) \cong H^0(X_{/C}, E|_{X_{/C}})$$

is an isomorphism. Now, finite flat group schemes are given by their associated (coherent) sheaves of Hopf algebras. Write \mathcal{O}_{G_U} and \mathcal{O}_{H_U} for the Hopf algebras of G_U and H_U respectively. Then $\text{Lef}(X, C)$ implies that, as *vector bundles*, \mathcal{O}_{G_U} and \mathcal{O}_{H_U} are isomorphic on U . We wish to show that \mathcal{O}_{G_U} and \mathcal{O}_{H_U} are isomorphic as *Hopf algebras* on U .

As the Hopf-algebra structures on are given as morphisms of vector bundles, the fact that $\mathcal{O}_{G_U}|_{X_{/C}} \cong \mathcal{O}_{H_U}|_{X_{/C}}$ as sheaves of Hopf algebras implies that \mathcal{O}_{G_U} and \mathcal{O}_{H_U} are isomorphic as sheaves of Hopf algebras on U . In particular, $G_U \cong H_U$ as desired. \square

As a byproduct of the algebraization machinery, we have the following, which is a cousin of Simpson's Theorem 1.1.

Corollary 8.6. *Let X/\mathbb{F}_q be a smooth projective variety with $\dim X \geq 2$ and let $U \subset X$ be an open subset whose complement has codimension at least 2. Let $C \subset U$ be a smooth curve that is the complete intersection of smooth, ample divisors of X . Let $A_C \rightarrow C$ be an abelian scheme of dimension g . Then the following are equivalent.*

- *There exists a Barsotti-Tate group \mathcal{G}_U on U such that $\mathcal{G}_C \cong A_C[p^\infty]$.*
- *There exists an abelian scheme $A_U \rightarrow U$ extending A_C .*

Proof. It suffices to prove the first statement implies the second. By Corollary 7.7, there exists a (polarizable) formal abelian scheme \hat{A} over $X_{/C}$ extending A_C . Picking a polarization, this yields a map $X_{/D} \rightarrow \mathcal{A}_{g,d}$.

By [Ked16, Theorem 5.1], the restriction functor $\mathbf{F}\text{-Isoc}(X) \rightarrow \mathbf{F}\text{-Isoc}(U)$ is an equivalence of categories. Denote the canonical extension of $\mathbb{D}(\mathcal{G}_U) \otimes \mathbb{Q} \in \mathbf{F}\text{-Isoc}(U)$ by $\mathcal{E} \in \mathbf{F}\text{-Isoc}(X)$. Our assumption implies that $\mathcal{E}_C \cong \mathbb{D}(A_C[p^\infty]) \otimes \mathbb{Q}$. Using the equivalence of categories and [Pál15, Theorem 1.2 and Proposition 4.9], it follows that \mathcal{E} is semi-simple.⁸

As $\det(\mathcal{E}|_C) \cong \mathbb{Q}_p(-g)$ it follows that $\det(\mathcal{E}) \cong \mathbb{Q}_p(-g)$. Every irreducible summand \mathcal{E}_i of \mathcal{E} is algebraic because it is on C . We note that for c a closed point of C , we have $P_c(\mathcal{E}, t) \in \mathbb{Q}[t]$.

Pick an isomorphism $\iota: \overline{\mathbb{Q}}_p \rightarrow \overline{\mathbb{Q}}_l$. By [AE19, Theorem 0.3] or [Ked18, Theorem 0.4.1], there is a unique semi-simple ι -companion to \mathcal{E} , which is a lisse $\overline{\mathbb{Q}}_l$ sheaf L . The l -adic sheaf L may be considered as a representation $W(X, \bar{c}) \rightarrow \text{GL}_{2g}(\overline{\mathbb{Q}}_l)$. The map $W(C, \bar{c}) \rightarrow W(X, \bar{c})$ is surjective by the usual Lefschetz theorems and the representation $W(C, \bar{c}) \rightarrow \text{GL}_{2g}(\overline{\mathbb{Q}}_l)$ conjugates into a representation $W(C, \bar{c}) \rightarrow \text{GL}_{2g}(\mathbb{Z}_l)$ because L_C is compatible with $A_C \rightarrow C$. Hence the representation L conjugates into a \mathbb{Z}_l local system on X . Take a connected finite étale cover of X' of X that trivializes the mod- l representation. Let C' and U' denote the inverse image to X' of C and U respectively, and note that C' is still the smooth complete intersection of smooth ample divisors inside of X' . Then we obtain

⁸While [Pál15, Theorem 1.2] only applies when the base is a curve, the main input is Proposition 4.9 and Remark 4.11 of *loc. cit.*, which do not depend on the base being a curve. The key is that \mathcal{E}_c is semi-simple for c a closed point of C .

a map $X'_{/C'} \rightarrow \mathcal{A}_{g,d,l}$, which globalizes to an open neighborhood $V' \subset X'$ of C' by Corollary 8.4. Lemma 8.5 implies that $A_{V'}[p^\infty] \cong \mathcal{G}_{V'}$, because they agree on $X'_{/C'}$. By Corollary 6.8, $A_{V'}$ extends to an abelian scheme $A_{U'} \rightarrow U'$ with $A_{U'}[p^\infty] \cong \mathcal{G}_{U'}$.

Finally, note that the the abelian scheme $A_{U'} \rightarrow U'$, equipped with isomorphisms $A_{U'}|_{C'} \cong A_{C'}$ and $A_{U'}[p^\infty] \cong \mathcal{G}_{U'}$ is *unique up to unique isomorphism*.⁹ Moreover, $A_{U'} \rightarrow U'$ is projective. Therefore we may apply faithfully flat descent to canonically descend to an abelian scheme $A_U \rightarrow U$, as desired. (This argument appears in [Gro66, p. 73, lines 7-12].) \square

Remark 8.7. We note that a slightly stronger variant of Corollary 8.6 is true when \mathbb{F}_q is replaced by a field k of characteristic 0. A Barsotti-Tate group in characteristic 0 is étale and hence satisfies Zariski-Nagata purity. One uses Simpson's Theorem 1.1 to prove that there is a family $A_{X,\mathbb{C}} \rightarrow X_{\mathbb{C}}$ extending $A_{C,\mathbb{C}} \rightarrow C_{\mathbb{C}}$. Then one transports the descent datum of $A_{C,\mathbb{C}}$ to $A_{X,\mathbb{C}}$ using Theorem 7.3. Finally, applying Weil descent [Var02, Appendix A] shows that we may find an abelian scheme $A_X \rightarrow X$ extending $A_C \rightarrow C$, as desired.

Remark 8.8. We note that the key point where we use that the base field is \mathbb{F}_q in Corollary 8.6 is to prove that a specific connected finite étale cover of C extends to a connected finite étale cover of X . We do this via l -adic companions.

Corollary 8.9. *Let X/k be a smooth projective variety over a field of characteristic p with $\dim X \geq 3$ and let $U \subset X$ be an open subset whose complement has codimension at least 3. Let $Z \subset U$ be a smooth complete subvariety of dimension at least 2 that is the complete intersection of smooth, ample divisors of X . Let $A_Z \rightarrow Z$ be an abelian scheme of dimension g . Then the following are equivalent.*

- *There exists a Barsotti-Tate group \mathcal{G}_U on U such that $\mathcal{G}_Z \cong A_Z[p^\infty]$.*
- *There exists an abelian scheme $A_U \rightarrow U$ extending A_Z .*

Proof. The key is that $\pi_1(Z) \rightarrow \pi_1(X)$ is an isomorphism. Therefore, a connected finite étale cover $Z' \rightarrow Z$ trivializing the l -torsion of $A_Z \rightarrow Z$ extends canonically to a connected finite étale cover $X' \rightarrow X$. Then the exact proof of Corollary 8.6 applies. \square

Remark 8.10. Corollary 8.6 is not true if one assumes that $\dim(X) = 1$ and C is a point. For example, let \mathcal{X} be the good reduction modulo p of a moduli space of fake elliptic curves, together with a universal family $\mathcal{A} \rightarrow \mathcal{X}$ of abelian surfaces with quaternionic multiplication. There is a decomposition $\mathcal{A}[p^\infty] \cong \mathcal{G} \oplus \mathcal{G}$, where \mathcal{G} is a height 2, dimension 1, everywhere versally deformed BT group \mathcal{G} . Over any point $x \in \mathcal{X}(\mathbb{F})$, there exists an elliptic curve E/\mathbb{F} such that $E[p^\infty] \cong \mathcal{G}_x$. On the other hand, the induced formal deformation of the elliptic curve $E \rightarrow x$ certainly does not extend to a non-isotrivial elliptic curve over X .

We now have all of the ingredients to prove our main Lefschetz theorem.

Theorem 8.11. *Let X/\mathbb{F}_q be a smooth projective variety. Then there exists an open subset $U \subset X$, whose complement has codimension at least 2, such that the following holds.*

Let \mathcal{E} be a semi-simple object of $\mathbf{F}\text{-Isoc}(X)_{\overline{\mathbb{Q}}_p}$ such that

- *the irreducible constituents of \mathcal{E} have rank 2 and determinant $\overline{\mathbb{Q}}_p(-1)$, and*
- *there exists a smooth projective curve $C \subset U$, the complete intersection of smooth ample divisors in X , and an abelian scheme $A_C \rightarrow C$ of dimension g such that*

$$\mathbb{D}(A_C[p^\infty]) \otimes \overline{\mathbb{Q}}_p \cong \mathcal{E}_C$$

⁹For instance, use [Gro66, Proposition 1.2 on p. 64]. Set $S = X$, let s be the generic point of X (which is the unique associated point of S), let s' be any point of C , and let S' be $\text{Spec}(\widehat{\mathcal{O}}_{S,s'})$ be the completion of the local ring at s' . Then $S' \rightarrow \text{Spec}(\widehat{\mathcal{O}}_{S,s'})$ is faithfully flat, and the Serre-Tate theorem will imply the desired uniqueness.

Then there exists an abelian scheme $B_U \rightarrow U$ of dimension g such that

$$\mathbb{D}(B_U[p^\infty]) \otimes \overline{\mathbb{Q}}_p \cong \mathcal{E}_U.$$

In particular, $A_C \rightarrow C$ extends to an abelian scheme over U after an isogeny.

Proof. We first construct U . It follows from [AE19, Corollary 4.3] that there are only finitely many isomorphism classes of irreducible overconvergent F -isocrystals \mathcal{F}_i on X of rank 2 and with determinant $\overline{\mathbb{Q}}_p(-1)$. We may pick a p -adic local field K over which each such \mathcal{F}_i is defined. By the slope bounds of Theorem 4.14, the two slopes of each such \mathcal{F}_i differ by at most one. As the two slopes add to one, this implies that the two slopes are in the interval $[0, 1]$. Then Lemma 5.8 implies that there is an open subset $U_i \subset X$, whose complement has codimension at least 2 on which \mathcal{F}_i underlies a Dieudonné \mathcal{M}_i in finite, locally free modules with $\mathcal{M}_i \otimes \mathbb{Q}_p \cong \mathcal{F}_i|_{U_i}$ (where on the left hand side we forget V and on the right hand side we forget the K -structure). Let $U := \cap U_i$. In particular every overconvergent F -isocrystal on X that is irreducible of rank 2 and has determinant $\overline{\mathbb{Q}}_p(-1)$ underlies a Dieudonné crystal on U . This is U in the first part of the theorem.

Now, let \mathcal{E} be as in the theorem. We make two claims about \mathcal{E} .

- (1) For every closed point x of X , the polynomial $P_x(\mathcal{E}, t)$ has coefficients in \mathbb{Q} .
- (2) The object \mathcal{E} may be descended to $\mathbf{F}\text{-Isoc}(X)$; that is, it has coefficients in \mathbb{Q}_p .

To prove the first part. Pick an isomorphism $\iota: \overline{\mathbb{Q}}_p \rightarrow \overline{\mathbb{Q}}_l$. By [AE19, Theorem 0.3] or [Ked18, Theorem 0.4.1], there is a unique semi-simple ι -companion to \mathcal{E} , which is a lisse $\overline{\mathbb{Q}}_l$ sheaf L . For c a closed point of C , the polynomial $P_c(L, t) \in \mathbb{Q}[t]$ by the companion relation. Applying [EK11, Proposition 7.1] and induction, we deduce that for all closed points x of X ,

$$P_x(L, t) \in \mathbb{Q}[t]$$

and hence $P_x(\mathcal{E}, t) \in \mathbb{Q}[t]$.

To prove the second part, for every $\sigma \in \text{Aut}_{\overline{\mathbb{Q}}_p}(\overline{\mathbb{Q}}_p)$, as $P_x(\mathcal{E}, t) \in \mathbb{Q}[t]$ for all closed points x of X , Theorem 3.16 implies that ${}^\sigma(\mathcal{E}) \cong \mathcal{E}$. To prove that \mathcal{E} descends to $\mathbf{F}\text{-Isoc}(X)$, we must construct isomorphisms $c_\sigma: {}^\sigma(\mathcal{E}) \cong \mathcal{E}$ that satisfy the cocycle condition by [Sos14, Lemma 4]. Now, the natural map

$$\text{Hom}({}^\sigma \mathcal{E}, \mathcal{E}) \rightarrow \text{Hom}({}^\sigma \mathcal{E}_C, \mathcal{E}_C)$$

is an isomorphism by Theorem 7.1. Note that \mathcal{E}_C descends to $\mathbf{F}\text{-Isoc}(C)$: indeed, $\mathbb{D}(A_C[p^\infty]) \otimes \mathbb{Q} \in \mathbf{F}\text{-Isoc}(C)$. Therefore, the descent data transports to the left and hence $\mathcal{E} \in \mathbf{F}\text{-Isoc}(X)$.

By construction of U , the F -isocrystal \mathcal{E} underlies a Dieudonné crystal (M, F, V) on U whose associated F -isocrystal is isomorphic to \mathcal{E}_U . By Theorem 5.7, there is an associated p -divisible group \mathcal{G}_U , with $\mathbb{D}(\mathcal{G}_U) \cong (M, F, V)$. It then follows that $A_C[p^\infty]$ is isogenous to \mathcal{G}_C : their Dieudonné crystals are isomorphic when viewed as F -isocrystals. Let $K \subset A_C[p^\infty]$ be the kernel of one such isogeny. Then $B_C := A_C/K$ is an abelian scheme with $B_C[p^\infty] \cong \mathcal{G}_C$. Applying Corollary 8.6, we see there exists an abelian scheme $B_U \rightarrow U$ (of dimension g) as desired. As $B_U[p^\infty]$ and \mathcal{G}_U are BT groups on U that are isomorphic on X/C , there are isomorphic by Lemma 8.5. Therefore B_U is compatible with \mathcal{E}_U , as desired. \square

It is natural to wonder whether or not one can remove both the open subset and the isogeny in Theorem 8.11, in further analogy with Theorem 1.1. Johan de Jong suggested the following example, which shows that the answer is “no” for general abelian varieties.

Example 8.12. Let E/\mathbb{F}_q be a supersingular elliptic curve, so $\alpha_p \times \alpha_p \subset E \times E$. Over $\mathbb{A}^2 \setminus (0, 0)$, there is an injective homomorphism

$$\iota: (\alpha_p)_{\mathbb{A}^2 \setminus (0,0)} \hookrightarrow (\alpha_p \times \alpha_p)_{\mathbb{A}^2 \setminus (0,0)} \subset (E \times E)_{\mathbb{A}^2 \setminus (0,0)}$$

given as follows: let (s, t) be linear coordinates on $\alpha_p \times \alpha_p \subset \mathbb{A}^2$ and let (x, y) be the coordinates on $\mathbb{A}^2 \setminus (0, 0)$. Then the above sub-group scheme is defined by the equation

$$[x : y] = [s : t].$$

Let A be the following (principally polarizable) supersingular abelian surface over $\mathbb{A}^2 \setminus (0,0)$:

$$A := (E \times E)_{\mathbb{A}^2 \setminus (0,0)} / \iota(\alpha_p)$$

[MB81]. (The principal polarization might only exist after a finite field extension.) Let $L \subset \mathbb{A}^2$ be a line not passing through the origin. Consider the compactification $\mathbb{A}^2 \setminus (0,0) \subset \mathbb{P}^2$ and let \bar{L} be the closure of L in \mathbb{P}^2 . Then A_L extends to a non-constant supersingular abelian scheme $A_{\bar{L}} \rightarrow \bar{L}$ and $\bar{L} \subset \mathbb{P}^2$ is a smooth, ample divisor. However, it is easy to see that $A_{\bar{L}}$ cannot extend to \mathbb{P}^2 . If it did, the family would have to be supersingular as the geometric monodromy is trivial and there is at least one supersingular fiber. Let $\mathcal{A}_{2,1} \otimes \mathbb{F}$ be the moduli of principally polarized abelian surfaces over \mathbb{F} . Then the supersingular locus of is a union of complete rational curves [Oor74, p. 177] and there are no morphisms $\mathbb{P}^2 \rightarrow \mathbb{P}^1$.

Corollary 8.13. *Let X/\mathbb{F}_q be a smooth projective variety with $\dim(X) \geq 2$. Then there exists an open subset $U \subset X$, whose complement has codimension at least 2, such that the following holds.*

Let L be a rank 2 lisse $\overline{\mathbb{Q}}_l$ sheaf with cyclotomic determinant and infinite geometric monodromy. Suppose there exists a complete set of p -adic companions (\mathcal{E}_i) for L . Then

- *if $C \subset U$ is a smooth proper curve that is the complete intersection of smooth ample divisors and*
- *if L_C comes from an abelian scheme of dimension g on $\pi_C: A_C \rightarrow C$ in the sense of Conjecture 1.2*

then L_U comes from an abelian scheme of dimension g $B_U \rightarrow U$, i.e., Conjecture 1.2 is true for (X, L) .

Proof. By the usual Lefschetz theorems, $W(C) \rightarrow W(X)$ is surjective. Therefore if two lisse $\overline{\mathbb{Q}}_l$ sheaves on X are isomorphic on C , then they are isomorphic. The U in our theorem is the U guaranteed to exist by Theorem 8.11.

Every irreducible summand of (the semi-simple lisse sheaf) $R^1\pi_{C*}\overline{\mathbb{Q}}_l$ is a companion of L_C by assumption, and hence has determinant $\overline{\mathbb{Q}}_l(-1)$. Therefore each of the \mathcal{E}_i has determinant $\overline{\mathbb{Q}}_p(-1)$. Consider $\mathbb{D}(A_C[p^\infty]) \otimes \overline{\mathbb{Q}}_p \in \mathbf{F}\text{-Isoc}(X)_{\overline{\mathbb{Q}}_p}$. This is semisimple by [Pál15]. There is a decomposition:

$$\mathbb{D}(A_C[p^\infty]) \otimes \overline{\mathbb{Q}}_p \cong \bigoplus_i (\mathcal{E}_i|_C)^{m_i}$$

for some integers $m_i \geq 1$ because the LHS is isomorphic to relative crystalline cohomology and [KM74]. Set $\mathcal{E} := \bigoplus_i (\mathcal{E}_i)^{m_i}$ and apply Theorem 8.11. \square

REFERENCES

- [Abe18a] Tomoyuki Abe. Langlands correspondence for isocrystals and the existence of crystalline companions for curves. *J. Amer. Math. Soc.*, 31(4):921–1057, 2018. doi:10.1090/jams/898.
- [Abe18b] Tomoyuki Abe. Langlands program for p -adic coefficients and the petits camarades conjecture. *J. Reine Angew. Math.*, 734:59–69, 2018. doi:10.1515/crelle-2015-0045.
- [AC13] Tomoyuki Abe and Daniel Caro. Theory of weights in p -adic cohomology. *arXiv preprint arXiv:1303.0662*, 2013.
- [AE19] Tomoyuki Abe and Hélène Esnault. A Lefschetz theorem for overconvergent isocrystals with Frobenius structure. *Ann. Scient. Éco. Norm. Sup.*, 52(5):1243–1264, 2019. URL: <https://doi.org/10.24033/asens.2408>, doi:10.24033/asens.2408.
- [BBM82] Pierre Berthelot, Lawrence Breen, and William Messing. *Théorie de Dieudonné cristalline II*, volume 930 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1982. doi:10.1007/BFb0093025.
- [CCO14] Ching-Li Chai, Brian Conrad, and Frans Oort. *Complex multiplication and lifting problems*, volume 195 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2014.
- [Chi04] CheeWhye Chin. Independence of l of monodromy groups. *J. Amer. Math. Soc.*, 17(3):723–747, 2004. doi:10.1090/S0894-0347-04-00456-4.
- [CO09] Ching-Li Chai and Frans Oort. Moduli of abelian varieties and p -divisible groups. 8:441–536, 2009.
- [Con06] Brian Conrad. Chow’s K/k -image and K/k -trace, and the Lang-Néron theorem. *Enseign. Math. (2)*, 52(1-2):37–108, 2006.
- [Cre92] Richard Crew. F -isocrystals and their monodromy groups. *Ann. Sci. École Norm. Sup. (4)*, 25(4):429–464, 1992. URL: http://www.numdam.org/item?id=ASENS_1992_4_25_4_429_0.

- [CS08] Kevin Corlette and Carlos Simpson. On the classification of rank-two representations of quasiprojective fundamental groups. *Compos. Math.*, 144(5):1271–1331, 2008. doi:10.1112/S0010437X08003618.
- [Del80] Pierre Deligne. La conjecture de Weil. II. *Inst. Hautes Études Sci. Publ. Math.*, (52):137–252, 1980. URL: http://www.numdam.org/item?id=PMIHES_1980__52__137_0.
- [Del12] Pierre Deligne. Finitude de l’extension de \mathbb{Q} engendrée par des traces de Frobenius, en caractéristique finie. *Mosc. Math. J.*, 12(3):497–514, 668, 2012.
- [Del14] Pierre Deligne. Semi-simplicité de produits tensoriels en caractéristique p . *Inventiones mathematicae*, 197(3):587–611, 2014.
- [dJ95] A. J. de Jong. Crystalline Dieudonné module theory via formal and rigid geometry. *Inst. Hautes Études Sci. Publ. Math.*, (82):5–96 (1996), 1995. URL: http://www.numdam.org/item?id=PMIHES_1995__82__5_0.
- [dJ98] A. J. de Jong. Homomorphisms of Barsotti-Tate groups and crystals in positive characteristic. *Invent. Math.*, 134(2):301–333, 1998. doi:10.1007/s002220050266.
- [dJM99] A. J. de Jong and W. Messing. Crystalline Dieudonné theory over excellent schemes. *Bull. Soc. Math. France*, 127(2):333–348, 1999. URL: http://www.numdam.org/item?id=BSMF_1999__127_2_333_0.
- [DK17] Vladimir Drinfeld and Kiran S. Kedlaya. Slopes of indecomposable \mathbb{F} -isocrystals. *Pure and Applied Mathematics Quarterly*, 13(1):131–192, 2017. URL: <https://doi.org/10.4310/pamq.2017.v13.n1.a5>, doi: 10.4310/pamq.2017.v13.n1.a5.
- [Dri77] V. G. Drinfel’d. Elliptic modules. II. *Math. USSR, Sb.*, 31:159–170, 1977.
- [Dri83] V. G. Drinfeld. Two-dimensional ℓ -adic representations of the fundamental group of a curve over a finite field and automorphic forms on $GL(2)$. *Am. J. Math.*, 105:85–114, 1983.
- [Dri12] Vladimir Drinfeld. On a conjecture of Deligne. *Moscow Mathematical Journal*, 12(3):515–542, 2012.
- [EG17] Hélène Esnault and Michael Groechenig. Cohomologically rigid local systems and integrality. *arXiv preprint arXiv:1711.06436*, 2017.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry. doi:10.1007/978-1-4612-5350-1.
- [EK11] Hélène Esnault and Moritz Kerz. Notes on Deligne’s letter to Drinfeld dated March 5, 2007. *Notes for the Forschungsseminar in Essen, summer*, 2011. Available at <http://userpage.fu-berlin.de/esnault/preprints/helene/103-110617.pdf>.
- [EK12] Hélène Esnault and Moritz Kerz. A finiteness theorem for Galois representations of function fields over finite fields (after Deligne). *arXiv preprint arXiv:1208.0128*, 2012.
- [Esn17] Hélène Esnault. Survey on some aspects of Lefschetz theorems in algebraic geometry. *Rev. Mat. Complut.*, 30(2):217–232, 2017. doi:10.1007/s13163-017-0223-8.
- [FC90] Gerd Faltings and Ching-Li Chai. *Degeneration of abelian varieties*, volume 22 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1990. With an appendix by David Mumford. doi:10.1007/978-3-662-02632-8.
- [FWG+92] Gerd Faltings, Gisbert Wüstholz, Fritz Grunewald, Norbert Schappacher, and Ulrich Stuhler. *Rational points*. Aspects of Mathematics, E6. Friedr. Vieweg & Sohn, Braunschweig, third edition, 1992. Papers from the seminar held at the Max-Planck-Institut für Mathematik, Bonn/Wuppertal, 1983/1984, With an appendix by Wüstholz. doi:10.1007/978-3-322-80340-5.
- [GD64] Alexandre Grothendieck and Jean Dieudonné. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I. *Inst. Hautes Études Sci. Publ. Math.*, (20):259, 1964. URL: http://www.numdam.org/article/PMIHES_1964__20__5_0.pdf.
- [GD66] Alexandre Grothendieck and Jean Dieudonné. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III. *Inst. Hautes Études Sci. Publ. Math.*, (28):255, 1966. URL: http://www.numdam.org/item?id=PMIHES_1966__28__255_0.pdf.
- [Gro66] Alexandre Grothendieck. Un théorème sur les homomorphismes de schémas abéliens. *Invent. Math.*, 2:59–78, 1966. doi:10.1007/BF01403390.
- [Gro74] Alexandre Grothendieck. *Groupes de Barsotti-Tate et cristaux de Dieudonné*. Les Presses de l’Université de Montréal, Montréal, Que., 1974. Séminaire de Mathématiques Supérieures, No. 45 (Été, 1970). URL: <https://www.imj-prg.fr/~leila.schneps/grothendieckcircle/barsottitate.pdf>.
- [Gro05] Alexander Grothendieck. Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2). 4:x+208, 2005. Séminaire de Géométrie Algébrique du Bois Marie, 1962, Augmenté d’un exposé de Michèle Raynaud. [With an exposé by Michèle Raynaud], With a preface and edited by Yves Laszlo, Revised reprint of the 1968 French original. URL: <https://arxiv.org/abs/math/0511279>.
- [GRR06] Alexandre Grothendieck, Michel Raynaud, and Dock Sang Rim. *Groupes de Monodromie en Géométrie Algébrique: Séminaire de Géométrie Algébrique du Bois-Marie 1967-1969. (SGA 7)*, volume 288. Springer, 2006.
- [Har70] Robin Hartshorne. *Ample subvarieties of algebraic varieties*. Lecture Notes in Mathematics, Vol. 156. Springer-Verlag, Berlin-New York, 1970. Notes written in collaboration with C. Musili.
- [HM68] Heisuke Hironaka and Hideyuki Matsumura. Formal functions and formal embeddings. *J. Math. Soc. Japan*, 20:52–82, 1968. URL: <https://doi.org/10.2969/jmsj/02010052>, doi:10.2969/jmsj/02010052.

- [HP18] Urs Hartl and Ambrus Pál. Crystalline Chebotarev density theorems. *arXiv preprint arXiv:1811.07084v1*, 2018.
- [Kat79] Nicholas M. Katz. Slope filtration of F -crystals. In *Journées de Géométrie Algébrique de Rennes (Rennes, 1978)*, Vol. I, volume 63 of *Astérisque*, pages 113–163. Soc. Math. France, Paris, 1979. Available at <https://web.math.princeton.edu/~nmk/old/f-crystals.pdf>.
- [Kat81] N. Katz. Serre-Tate local moduli. In *Algebraic surfaces (Orsay, 1976–78)*, volume 868 of *Lecture Notes in Math.*, pages 138–202. Springer, Berlin-New York, 1981.
- [Ked04] Kiran S. Kedlaya. Full faithfulness for overconvergent F -isocrystals. In *Geometric aspects of Dwork theory. Vol. I, II*, pages 819–835. Walter de Gruyter, Berlin, 2004.
- [Ked16] Kiran S Kedlaya. Notes on isocrystals. *arXiv preprint arXiv:1606.01321*, 2016. [Online; accessed 11-December-2017].
- [Ked18] Kiran S Kedlaya. Étale and crystalline companions I. *draft*, 2018. [Online; accessed 20-September-2018]. URL: <http://http://kskedlaya.org/papers/companions.pdf>.
- [KM74] Nicholas M. Katz and William Messing. Some consequences of the Riemann hypothesis for varieties over finite fields. *Invent. Math.*, 23:73–77, 1974.
- [KM19] Joe Kramer-Miller. Log decay F -isocrystals on higher dimensional varieties. *arXiv preprint arXiv:1902.04730*, 2019.
- [Kos17] Teruhisa Koshikawa. Overconvergent unit-root F -isocrystals and isotriviality. *Math. Res. Lett.*, 24(6):1707–1727, 2017. doi:10.4310/MRL.2017.v24.n6.a7.
- [Kri17] Raju Krishnamoorthy. Rank 2 local systems, Barsotti-Tate groups, and Shimura curves. *arXiv preprint arXiv:1711.04797*, 2017.
- [Laf02] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Invent. Math.*, 147(1):1–241, 2002. doi:10.1007/s002220100174.
- [Laf11] Vincent Lafforgue. Estimées pour les valuations p -adiques des valeurs propres des opérateurs de Hecke. *Bull. Soc. Math. France*, 139(4):455–477, 2011.
- [Lit18] Daniel Litt. Arithmetic representations of fundamental groups II: finiteness. *arXiv preprint arXiv:1809.03524*, 2018.
- [LO98] Ke-Zheng Li and Frans Oort. *Moduli of supersingular abelian varieties*, volume 1680 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1998. doi:10.1007/BFb0095931.
- [MB81] Laurent Moret-Bailly. Familles de courbes et de variétés abéliennes sur \mathbb{P}^1 . *Sém. sur les pincesaux de courbes de genre au moins deux (ed. L. Szpiro). Astérisques*, 86:109–140, 1981.
- [Mor85] Laurent Moret-Bailly. *Pincesaux de variétés abéliennes. (Pencils of abelian varieties)*., volume 129. Société Mathématique de France (SMF), Paris, 1985.
- [NN81] M. S. Narasimhan and M. V. Nori. Polarizations on an abelian variety. volume 90, pages 125–128, 1981. doi:10.1007/BF02837283.
- [Oor74] Frans Oort. Subvarieties of moduli spaces. *Invent. Math.*, 24:95–119, 1974. URL: <https://doi.org/10.1007/BF01404301>, doi:10.1007/BF01404301.
- [Pál15] Ambrus Pál. The p -adic monodromy group of abelian varieties over global function fields of characteristic p . *arXiv preprint arXiv:1512.03587*, 2015.
- [Shi67] Goro Shimura. Algebraic number fields and symplectic discontinuous groups. *Ann. of Math. (2)*, 86:503–592, 1967. URL: <https://doi.org/10.2307/1970613>, doi:10.2307/1970613.
- [Shi79] Tetsuji Shioda. Supersingular $K3$ surfaces. In *Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, Copenhagen, 1978)*, volume 732 of *Lecture Notes in Math.*, pages 564–591. Springer, Berlin, 1979.
- [Sim91] Carlos T. Simpson. Nonabelian Hodge theory. In *Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990)*, pages 747–756. Math. Soc. Japan, Tokyo, 1991.
- [Sim92] Carlos T. Simpson. Higgs bundles and local systems. *Inst. Hautes Études Sci. Publ. Math.*, (75):5–95, 1992. URL: http://www.numdam.org/item?id=PMIHES_1992__75__5_0.
- [Sos14] Paweł Sosna. Scalar extensions of triangulated categories. *Applied categorical structures*, 22(1):211–227, 2014.
- [ST68] Jean-Pierre Serre and John Tate. Good reduction of abelian varieties. *Ann. of Math. (2)*, 88:492–517, 1968. doi:10.2307/1970722.
- [ST18] Andrew Snowden and Jacob Tsimerman. Constructing elliptic curves from galois representations. *Compositio Mathematica*, 154(10):2045–2054, 2018. URL: <https://arxiv.org/abs/1708.02656>, doi:10.1112/S0010437X18007315.
- [Sta20] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2020.
- [Tat67] John Tate. p -divisible groups. In *Proc. Conf. Local Fields (Driebergen, 1966)*, pages 158–183. Springer, Berlin, 1967.
- [Var02] Yakov Varshavsky. On the characterization of complex Shimura varieties. *Selecta Mathematica*, 8(2):283–314, 2002.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30605, USA

Email address: `a.pal@imperial.ac.uk`

DEPARTMENT OF MATHEMATICS, 180 QUEENS GATE, IMPERIAL COLLEGE, LONDON, SW7 2AZ, UNITED KINGDOM