

The Fyodorov-Bouchaud formula and Liouville conformal field theory

Guillaume Remy

École Normale Supérieure

February 1, 2018

Introduction

Statistical Physics:

- Log-correlated fields
- Gaussian multiplicative chaos (GMC)
- The Fyodorov-Bouchaud formula

Liouville Field Theory:

- Conformal field theory
- Correlation functions and BPZ equation
- Integrability: DOZZ formula

Introduction

DKRV 2014, link between:

Gaussian multiplicative chaos \Leftrightarrow Liouville correlations

Introduction

DKRV 2014, link between:

Gaussian multiplicative chaos \Leftrightarrow Liouville correlations

Consequences:

- Proof of the DOZZ formula KRV 2017
- Proof of the Fyodorov-Bouchaud formula R 2017

\Rightarrow Integrability program for GMC and Liouville theory

Outline

- 1 Definitions, main result and applications
- 2 Proof of the Fyodorov-Bouchaud formula
- 3 The Liouville conformal field theory
- 4 Outlook and perspectives

Gaussian Free Field (GFF)

Gaussian free field X on the unit circle $\partial\mathbb{D}$

$$\mathbb{E}[X(e^{i\theta})X(e^{i\theta'})] = 2 \ln \frac{1}{|e^{i\theta} - e^{i\theta'}|}$$

- $X(e^{i\theta})$ has an infinite variance
- X lives in the space of distributions
- Cut-off approximation X_ϵ

Ex: $X_\epsilon = \rho_\epsilon * X$, $\rho_\epsilon = \frac{1}{\epsilon} \rho(\frac{\cdot}{\epsilon})$, with smooth ρ .

Gaussian multiplicative chaos (GMC)

For $\gamma \in (0, 2)$, define on $\partial\mathbb{D}$ the measure $e^{\frac{\gamma}{2}X}d\theta$

Gaussian multiplicative chaos (GMC)

For $\gamma \in (0, 2)$, define on $\partial\mathbb{D}$ the measure $e^{\frac{\gamma}{2}X}d\theta$

- Cut-off approximation $e^{\frac{\gamma}{2}X_\epsilon}d\theta$
- $\mathbb{E}[e^{\frac{\gamma}{2}X_\epsilon}] = e^{\frac{\gamma^2}{8}\mathbb{E}[X_\epsilon^2]}$
- Renormalized measure: $e^{\frac{\gamma}{2}X_\epsilon - \frac{\gamma^2}{8}\mathbb{E}[X_\epsilon^2]}d\theta$

Gaussian multiplicative chaos (GMC)

For $\gamma \in (0, 2)$, define on $\partial\mathbb{D}$ the measure $e^{\frac{\gamma}{2}X}d\theta$

- Cut-off approximation $e^{\frac{\gamma}{2}X_\epsilon}d\theta$
- $\mathbb{E}[e^{\frac{\gamma}{2}X_\epsilon}] = e^{\frac{\gamma^2}{8}\mathbb{E}[X_\epsilon^2]}$
- Renormalized measure: $e^{\frac{\gamma}{2}X_\epsilon - \frac{\gamma^2}{8}\mathbb{E}[X_\epsilon^2]}d\theta$


Proposition (Kahane 1985)

The following limit exists in the sense of weak convergence of measures, $\forall \gamma \in (0, 2)$:

$$e^{\frac{\gamma}{2}X(e^{i\theta})}d\theta := \lim_{\epsilon \rightarrow 0} e^{\frac{\gamma}{2}X_\epsilon(e^{i\theta}) - \frac{\gamma^2}{8}\mathbb{E}[X_\epsilon^2(e^{i\theta})]}d\theta$$


Illustrations of the GMC measures

Simulations of the measure $e^{\gamma X} dx^2$ on $[0, 1]^2$.



$$\gamma = 1$$

Simulations by Tunan Zhu



$$\gamma = 1.8$$

Moments of the GMC

We introduce:

$$\forall \gamma \in (0, 2), Y_\gamma := \frac{1}{2\pi} \int_0^{2\pi} e^{\frac{\gamma}{2} X(e^{i\theta})} d\theta$$

Existence of the moments of Y_γ :

$$\mathbb{E}[Y_\gamma^p] < +\infty \iff p < \frac{4}{\gamma^2}.$$

The Fyodorov-Bouchaud formula

Theorem (Remy 2017)

Let $\gamma \in (0, 2)$ and $p \in (-\infty, \frac{4}{\gamma^2})$, then:

$$\mathbb{E}[Y_\gamma^p] = \frac{\Gamma(1 - p \frac{\gamma^2}{4})}{\Gamma(1 - \frac{\gamma^2}{4})^p}$$

We also have a density for Y_γ ,

$$f_{Y_\gamma}(y) = \frac{4\beta}{\gamma^2} (\beta y)^{-\frac{4}{\gamma^2}-1} e^{-(\beta y)^{-\frac{4}{\gamma^2}}} \mathbf{1}_{[0, \infty[}(y),$$

where we have set $\beta = \Gamma(1 - \frac{\gamma^2}{4})$.

Some applications

- ➊ Maximum of the Gaussian free field
- ➋ Random matrix theory
- ➌ Tail expansion for Gaussian multiplicative chaos

Application 1: maximum of the GFF

Derivative martingale: work by Duplantier, Rhodes, Sheffield, Vargas.

$\gamma \rightarrow 2$ in our GMC measure (Aru, Powell, Sepúlveda):

$$Y' := -\frac{1}{2\pi} \int_0^{2\pi} X(e^{i\theta}) e^{X(e^{i\theta})} d\theta = \lim_{\gamma \rightarrow 2} \frac{1}{2-\gamma} Y_\gamma.$$

In Y' has the following density:

$$f_{\ln Y'}(y) = e^{-y} e^{-e^{-y}}$$

$\ln Y' \sim \mathcal{G}$ where \mathcal{G} follows a standard Gumbel law

Application 1: maximum of the GFF

Following an impressive series of works (2016):

Theorem (Ding, Madaule, Roy, Zeitouni)

For a reasonable cut-off X_ϵ of the GFF:

$$\max_{\theta \in [0, 2\pi]} X_\epsilon(e^{i\theta}) - 2 \ln \frac{1}{\epsilon} + \frac{3}{2} \ln \ln \frac{1}{\epsilon} \xrightarrow[\epsilon \rightarrow 0]{} \mathcal{G} + \text{In } Y' + C$$

where \mathcal{G} is a standard Gumbel law and $C \in \mathbb{R}$.

Application 1: maximum of the GFF

The Fyodorov-Bouchaud formula implies:

Corollary (Remy 2017)

For a reasonable cut-off X_ϵ of the GFF:

$$\max_{\theta \in [0, 2\pi]} X_\epsilon(e^{i\theta}) - 2 \ln \frac{1}{\epsilon} + \frac{3}{2} \ln \ln \frac{1}{\epsilon} \xrightarrow[\epsilon \rightarrow 0]{} \mathcal{G}_1 + \mathcal{G}_2 + C$$

where $\mathcal{G}_1, \mathcal{G}_2$ are independent Gumbel laws and $C \in \mathbb{R}$.

Application 2: random unitary matrices

$U_N := N \times N$ random unitary matrix

Its eigenvalues $(e^{i\theta_1}, \dots, e^{i\theta_n})$ follow the distribution:

$$\frac{1}{n!} \prod_{k < j} |e^{i\theta_k} - e^{i\theta_j}|^2 \prod_{k=1}^n \frac{d\theta_k}{2\pi}$$

Let $p_N(\theta) = \det(1 - e^{-i\theta} U_N) = \prod_{k=1}^N (1 - e^{i(\theta_k - \theta)})$

Webb (2015): $\forall \alpha \in (-\frac{1}{2}, \sqrt{2})$,

$$\frac{|p_N(\theta)|^\alpha}{\mathbb{E}[|p_N(\theta)|^\alpha]} d\theta \xrightarrow[N \rightarrow \infty]{} e^{\frac{|\alpha|}{2} X(e^{i\theta})} d\theta$$

Application 2: random unitary matrices

Conjecture by Fyodorov, Hiary, Keating (2012):

$$\max_{\theta \in [0, 2\pi]} \ln |p_N(\theta)| - \ln N + \frac{3}{4} \ln \ln N \xrightarrow[N \rightarrow \infty]{} \mathcal{G}_1 + \mathcal{G}_2 + C.$$

Chhaibi, Madaule, Najnudel (2016), tightness of:

$$\max_{\theta \in [0, 2\pi]} \ln |p_N(\theta)| - \ln N + \frac{3}{4} \ln \ln N.$$

With our result it is sufficient to show:

$$\max_{\theta \in [0, 2\pi]} \ln |p_N(\theta)| - \ln N + \frac{3}{4} \ln \ln N \xrightarrow[N \rightarrow \infty]{} \mathcal{G}_1 + \ln Y' + C.$$

Application 3: Tail estimates for GMC

$$\mathbb{E}[\tilde{X}(e^{i\theta})\tilde{X}(e^{i\theta'})] = 2 \ln \frac{1}{|e^{i\theta} - e^{i\theta'}|} + f(e^{i\theta}, e^{i\theta'})$$

Proposition (Rhodes, Vargas 2017)

$\mathcal{O} \subset \partial \mathbb{D}$, $\exists \delta > 0$:

$$\mathbb{P}\left(\int_{\mathcal{O}} e^{\frac{\gamma}{2}\tilde{X}(e^{i\theta})} d\theta > t\right) = \frac{C(\gamma)}{t^{\frac{4}{\gamma^2}}} + o_{t \rightarrow \infty}(t^{-\frac{4}{\gamma^2} - \delta})$$

where $C(\gamma) = \overline{R}_1(\gamma) \int_{\mathcal{O}} e^{(\frac{4}{\gamma^2}-1)f(e^{i\theta}, e^{i\theta})} d\theta$

Application 3: Tail estimates for GMC

$$\mathbb{E}[\tilde{X}(e^{i\theta})\tilde{X}(e^{i\theta'})] = 2 \ln \frac{1}{|e^{i\theta} - e^{i\theta'}|} + f(e^{i\theta}, e^{i\theta'})$$

Corollary (Remy 2017)

$\mathcal{O} \subset \partial \mathbb{D}$, $\exists \delta > 0$:

$$\mathbb{P}\left(\int_{\mathcal{O}} e^{\frac{\gamma}{2}\tilde{X}(e^{i\theta})} d\theta > t\right) = \frac{C(\gamma)}{t^{\frac{4}{\gamma^2}}} + o_{t \rightarrow \infty}(t^{-\frac{4}{\gamma^2} - \delta})$$

$$\text{where } C(\gamma) = \frac{(2\pi)^{\frac{4}{\gamma^2}-1}}{\Gamma(1-\frac{\gamma^2}{4})^{\frac{4}{\gamma^2}}} \int_{\mathcal{O}} e^{(\frac{4}{\gamma^2}-1)f(e^{i\theta}, e^{i\theta})} d\theta$$

Integer moments of the GMC

The computation of Fyodorov and Bouchaud

Fyodorov Y., Bouchaud J.P.: Freezing and extreme value statistics in a Random Energy Model with logarithmically correlated potential, *Journal of Physics A: Mathematical and Theoretical*, Volume 41, Number 37, (2008).

Integer moments of the GMC

For $n \in \mathbb{N}^*$, $n < \frac{4}{\gamma^2}$:

$$\begin{aligned} & \mathbb{E}\left[\left(\frac{1}{2\pi} \int_0^{2\pi} e^{\frac{\gamma}{2}X_\epsilon(e^{i\theta}) - \frac{\gamma^2}{8}\mathbb{E}[X_\epsilon(e^{i\theta})^2]} d\theta\right)^n\right] \\ &= \frac{1}{(2\pi)^n} \int_{[0,2\pi]^n} \mathbb{E}\left[\prod_{i=1}^n e^{\frac{\gamma}{2}X_\epsilon(e^{i\theta_i}) - \frac{\gamma^2}{8}\mathbb{E}[X_\epsilon(e^{i\theta_i})^2]}\right] d\theta_1 \dots d\theta_n \\ &= \frac{1}{(2\pi)^n} \int_{[0,2\pi]^n} e^{\frac{\gamma^2}{4} \sum_{i < j} \mathbb{E}[X_\epsilon(e^{i\theta_i}) X_\epsilon(e^{j\theta_j})]} d\theta_1 \dots d\theta_n \end{aligned}$$

Integer moments of the GMC

For $n \in \mathbb{N}^*$, $n < \frac{4}{\gamma^2}$:

$$\begin{aligned}\mathbb{E}[Y_\gamma^n] &= \frac{1}{(2\pi)^n} \int_{[0,2\pi]^n} e^{\frac{\gamma^2}{4} \sum_{i < j} \mathbb{E}[X(e^{i\theta_i})X(e^{i\theta_j})]} d\theta_1 \dots d\theta_n \\ &= \frac{1}{(2\pi)^n} \int_{[0,2\pi]^n} \prod_{i < j} \frac{1}{|e^{i\theta_i} - e^{i\theta_j}|^{\frac{\gamma^2}{2}}} d\theta_1 \dots d\theta_n \\ &= \frac{\Gamma(1 - n\frac{\gamma^2}{4})}{\Gamma(1 - \frac{\gamma^2}{4})^n}\end{aligned}$$

- Question: can we replace $n \in \mathbb{N}^*$ by a real $p < \frac{4}{\gamma^2}$?

Proof of the Fyodorov-Bouchaud formula

Framework of **conformal field theory**

Belavin A.A., Polyakov A.M., Zamolodchikov A.B.:
Infinite conformal symmetry in two-dimensional quantum
field theory, *Nuclear. Physics.*, B241, 333-380, (1984).

The BPZ differential equation

We introduce the following observable for $t \in [0, 1]$:

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta\right)^p\right]$$

The BPZ differential equation

We introduce the following observable for $t \in [0, 1]$:

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta\right)^p\right]$$

BPZ equation:

$$(t(1-t^2)\frac{\partial^2}{\partial t^2} + (t^2-1)\frac{\partial}{\partial t} + 2(C - (A+B+1)t^2)\frac{\partial}{\partial t} - 4ABt)G(\gamma, p, t) = 0$$

where:

$$A = -\frac{\gamma^2 p}{4}, \quad B = -\frac{\gamma^2}{4}, \quad C = \frac{\gamma^2}{4}(1-p) + 1.$$

Solutions of the BPZ equation

BPZ equation in $t \rightarrow$ hypergeometric equation in t^2

Two bases of solutions:

- $G(\gamma, p, t) = C_1 F_1(t^2) + C_2 t^{\frac{\gamma^2}{2}(p-1)} F_2(t^2)$
- $G(\gamma, p, t) = B_1 \tilde{F}_1(1-t^2) + B_2 (1-t^2)^{1+\frac{\gamma^2}{2}} \tilde{F}_2(1-t^2)$

where:

- $C_1, C_2, B_1, B_2 \in \mathbb{R}$
- $F_1, F_2, \tilde{F}_1, \tilde{F}_2$:= hypergeometric series depending on γ and p .

Change of basis: $(C_1, C_2) \leftrightarrow (B_1, B_2)$.

Link between C_1 , C_2 , B_1 and B_2

Goal: identify C_1 , C_2 , B_1 , B_2

$$G(\gamma, p, t) = \mathbb{E}[(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta)^p]$$

Link between C_1 , C_2 , B_1 and B_2

Goal: identify C_1 , C_2 , B_1 , B_2

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$$

- $C_1 = G(\gamma, p, 0) = (2\pi)^p \mathbb{E}[Y_\gamma^p]$

Link between C_1 , C_2 , B_1 and B_2

Goal: identify C_1 , C_2 , B_1 , B_2

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$$

- $C_1 = G(\gamma, p, 0) = (2\pi)^p \mathbb{E}[Y_\gamma^p]$
- $B_1 = G(\gamma, p, 1) = \mathbb{E}\left[\left(\int_0^{2\pi} |1 - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$

Link between C_1 , C_2 , B_1 and B_2

Goal: identify C_1 , C_2 , B_1 , B_2

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$$

- $C_1 = G(\gamma, p, 0) = (2\pi)^p \mathbb{E}[Y_\gamma^p]$
- $B_1 = G(\gamma, p, 1) = \mathbb{E}\left[\left(\int_0^{2\pi} |1 - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$
- $G(\gamma, p, t) - G(\gamma, p, 0) = O(t^2) \Rightarrow C_2 = 0$

Link between C_1 , C_2 , B_1 and B_2

Goal: identify C_1 , C_2 , B_1 , B_2

$$G(\gamma, p, t) = \mathbb{E}\left[\left(\int_0^{2\pi} |t - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$$

- $C_1 = G(\gamma, p, 0) = (2\pi)^p \mathbb{E}[Y_\gamma^p]$
- $B_1 = G(\gamma, p, 1) = \mathbb{E}\left[\left(\int_0^{2\pi} |1 - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^p\right]$
- $G(\gamma, p, t) - G(\gamma, p, 0) = O(t^2) \Rightarrow C_2 = 0$
- $B_2 = ??$

The value of B_2

Let $h_u(t) = |t - e^{iu}|^{\frac{\gamma^2}{2}}$

- $G(\gamma, p, t) - G(\gamma, p, 1) = p \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu}) + R(t)$

with $c(e^{iu}) = \mathbb{E}\left[\left(\int_0^{2\pi} \frac{|1-e^{i\theta}|^{\frac{\gamma^2}{2}}}{|e^{iu}-e^{i\theta}|^{\frac{\gamma^2}{2}}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^{p-1}\right]$

The value of B_2

Let $h_u(t) = |t - e^{iu}|^{\frac{\gamma^2}{2}}$

- $G(\gamma, p, t) - G(\gamma, p, 1) = p \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu}) + R(t)$

with $c(e^{iu}) = \mathbb{E}\left[\left(\int_0^{2\pi} \frac{|1-e^{i\theta}|^{\frac{\gamma^2}{2}}}{|e^{iu}-e^{i\theta}|^{\frac{\gamma^2}{2}}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^{p-1}\right]$

- $\frac{1}{2\pi} \int_0^{2\pi} du (h_u(t) - h_u(1)) = \hat{F}_1(t) + \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})^2} (1-t^2)^{1+\frac{\gamma^2}{2}} \hat{F}_2(t)$

The value of B_2

Let $h_u(t) = |t - e^{iu}|^{\frac{\gamma^2}{2}}$

- $G(\gamma, p, t) - G(\gamma, p, 1) = p \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu}) + R(t)$

with $c(e^{iu}) = \mathbb{E}\left[\left(\int_0^{2\pi} \frac{|1-e^{i\theta}|^{\frac{\gamma^2}{2}}}{|e^{iu}-e^{i\theta}|^{\frac{\gamma^2}{2}}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^{p-1}\right]$

- $\frac{1}{2\pi} \int_0^{2\pi} du (h_u(t) - h_u(1)) = \hat{F}_1(t) + \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})^2} (1-t^2)^{1+\frac{\gamma^2}{2}} \hat{F}_2(t)$

- $\frac{1}{2\pi} \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu})$
 $= A(1-t) + \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})^2} (1-t^2)^{1+\frac{\gamma^2}{2}} c(1) + o((1-t)^{1+\frac{\gamma^2}{2}})$

The value of B_2

Let $h_u(t) = |t - e^{iu}|^{\frac{\gamma^2}{2}}$

- $G(\gamma, p, t) - G(\gamma, p, 1) = p \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu}) + R(t)$

with $c(e^{iu}) = \mathbb{E}\left[\left(\int_0^{2\pi} \frac{|1-e^{i\theta}|^{\frac{\gamma^2}{2}}}{|e^{iu}-e^{i\theta}|^{\frac{\gamma^2}{2}}} e^{\frac{\gamma}{2}X(\theta)} d\theta\right)^{p-1}\right]$

- $\frac{1}{2\pi} \int_0^{2\pi} du (h_u(t) - h_u(1)) = \hat{F}_1(t) + \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})^2} (1-t^2)^{1+\frac{\gamma^2}{2}} \hat{F}_2(t)$

- $\frac{1}{2\pi} \int_0^{2\pi} du (h_u(t) - h_u(1)) c(e^{iu})$
 $= A(1-t) + \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})^2} (1-t^2)^{1+\frac{\gamma^2}{2}} c(1) + o((1-t)^{1+\frac{\gamma^2}{2}})$

$$\implies B_2 = (2\pi)^p p \frac{\Gamma(-\frac{\gamma^2}{2}-1)}{\Gamma(-\frac{\gamma^2}{4})} \mathbb{E}[Y_\gamma^{p-1}]$$

The shift relation

Change of basis:

$$\begin{pmatrix} B_1 \\ B_2 \end{pmatrix} = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$$

and $C_2 = 0$

$$\Rightarrow B_2 = \frac{\Gamma(-1 - \frac{\gamma^2}{2})\Gamma(\frac{\gamma^2}{4}(1-p) + 1)}{\Gamma(-\frac{\gamma^2}{4})\Gamma(-\frac{\gamma^2 p}{4})} C_1$$

Therefore:

$$\mathbb{E}[Y_\gamma^p] = \frac{\Gamma(1 - p\frac{\gamma^2}{4})}{\Gamma(1 - \frac{\gamma^2}{4})\Gamma(1 - (p-1)\frac{\gamma^2}{4})} \mathbb{E}[Y_\gamma^{p-1}].$$

Negative moments of GMC

The shift relation gives all the negative moments:

$$\mathbb{E}[Y_\gamma^{-n}] = \Gamma\left(1 + \frac{n\gamma^2}{4}\right)\Gamma\left(1 - \frac{\gamma^2}{4}\right)^n, \quad \forall n \in \mathbb{N}.$$

We check:

$$\forall \lambda \in \mathbb{R}, \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \Gamma\left(1 + \frac{n\gamma^2}{4}\right)\Gamma\left(1 - \frac{\gamma^2}{4}\right)^n < +\infty$$

Negative moments \Rightarrow determine the law of Y_γ !

Explicit probability densities

Probability densities for Y_γ^{-1} and Y_γ

$$f_{\frac{1}{Y_\gamma}}(y) = \frac{4}{\beta\gamma^2} \left(\frac{y}{\beta}\right)^{\frac{4}{\gamma^2}-1} e^{-(\frac{y}{\beta})^{\frac{4}{\gamma^2}}} \mathbf{1}_{[0,\infty[}(y)$$

$$f_{Y_\gamma}(y) = \frac{4\beta}{\gamma^2} (\beta y)^{-\frac{4}{\gamma^2}-1} e^{-(\beta y)^{-\frac{4}{\gamma^2}}} \mathbf{1}_{[0,\infty[}(y)$$

where $\gamma \in (0, 2)$ and $\beta = \Gamma(1 - \frac{\gamma^2}{4})$.

Generalizations to log-singularities

1) Degenerate insertions $-\frac{\gamma}{2}$ and $-\frac{2}{\gamma}$

$$\mathbb{E}\left[\left(\frac{1}{2\pi} \int_0^{2\pi} |1 - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta\right)^p\right] = \frac{\Gamma(1 - \frac{\gamma^2}{4}p)}{\Gamma(1 - \frac{\gamma^2}{4})^p} \frac{\Gamma(1 + \frac{\gamma^2}{2})}{\Gamma(1 + \frac{\gamma^2}{4})} \frac{\Gamma(1 + (1-p)\frac{\gamma^2}{4})}{\Gamma(1 + (2-p)\frac{\gamma^2}{4})}$$

$$\mathbb{E}\left[\left(\frac{1}{2\pi} \int_0^{2\pi} |1 - e^{i\theta}|^{\frac{2}{\gamma}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta\right)^p\right] = \frac{\Gamma(1 - \frac{\gamma^2}{4}p)}{\Gamma(1 - \frac{\gamma^2}{4})^p} \frac{\Gamma(1 + \frac{8}{\gamma^2})}{\Gamma(1 + \frac{4}{\gamma^2})} \frac{\Gamma(\frac{4}{\gamma^2} - p + 1)}{\Gamma(\frac{8}{\gamma^2} - p + 1)}$$

Generalizations to log-singularities

1) Degenerate insertions $-\frac{\gamma}{2}$ and $-\frac{2}{\gamma}$

$$\frac{1}{2\pi} \int_0^{2\pi} |1 - e^{i\theta}|^{\frac{\gamma^2}{2}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta \stackrel{\text{law}}{=} Y_\gamma X_1^{-\frac{\gamma^2}{4}}$$

$$\frac{1}{2\pi} \int_0^{2\pi} |1 - e^{i\theta}|^2 e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta \stackrel{\text{law}}{=} Y_\gamma X_2^{-1}$$

Y_γ, X_1, X_2 independent, $X_1 \sim \mathcal{B}(1 + \frac{\gamma^2}{4}, \frac{\gamma^2}{4})$, $X_2 \sim \mathcal{B}(1 + \frac{4}{\gamma^2}, \frac{4}{\gamma^2})$.

$$f_{\mathcal{B}(\alpha, \beta)}(x) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} \mathbf{1}_{[0,1]}(x)$$

Generalizations to log-singularities

2) Arbitrary log-singularity, $\alpha < Q$, $p < \frac{4}{\gamma^2} \wedge \frac{2}{\gamma}(Q - \alpha)$,

$$\begin{aligned} & \mathbb{E}\left[\left(\frac{1}{2\pi} \int_0^{2\pi} \frac{1}{|1 - e^{i\theta}|^{\alpha\gamma}} e^{\frac{\gamma}{2}X(e^{i\theta})} d\theta\right)^p\right] \\ &= \frac{\Gamma(1 - \frac{\gamma^2 p}{4})}{\Gamma(1 - \frac{\gamma^2}{4})^p} \frac{\mathcal{T}_\gamma(\alpha + \frac{\gamma p}{2})^2 \mathcal{T}_\gamma(2\alpha) \mathcal{T}_\gamma(0)}{\mathcal{T}_\gamma(2\alpha + \frac{\gamma p}{2}) \mathcal{T}_\gamma(\alpha)^2 \mathcal{T}_\gamma(\frac{\gamma p}{2})} \end{aligned}$$

$$\ln \mathcal{T}_\gamma(x) = \int_0^\infty \frac{dt}{t} \left(\frac{e^{-\frac{Qt}{2}} - e^{-(Q-x)t}}{(1 - e^{-\frac{\gamma t}{2}})(1 - e^{-\frac{2t}{\gamma}})} - \frac{e^{-t}}{2} \left(\frac{Q}{2} - x \right)^2 - \frac{Q-2x}{2t} \right)$$

$$\text{and } Q = \frac{\gamma}{2} + \frac{2}{\gamma}.$$

Work in progress with Tunan Zhu.

What is Liouville field theory?

Path integral formalism

$$\Sigma = \{X : \mathbb{D} \rightarrow \mathbb{R}\}$$

What is Liouville field theory?

Path integral formalism

$$\Sigma = \{X : \mathbb{D} \rightarrow \mathbb{R}\}$$

For $X \in \Sigma$, energy of $X := \frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2$

What is Liouville field theory?

Path integral formalism

$$\Sigma = \{X : \mathbb{D} \rightarrow \mathbb{R}\}$$

For $X \in \Sigma$, energy of $X := \frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2$

Random field ϕ_G :

$$\mathbb{E}[F(\phi_G)] = \int_{\Sigma} F(X) e^{-\frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2} DX$$

What is Liouville field theory?

Path integral formalism

$$\Sigma = \{X : \mathbb{D} \rightarrow \mathbb{R}\}$$

For $X \in \Sigma$, energy of $X := \frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2$

Random field ϕ_G :

$$\mathbb{E}[F(\phi_G)] = \int_{\Sigma} F(X) e^{-\frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2} DX$$

Informally: $\int |\partial X|^2 dx^2 = - \int X \Delta X dx^2$

$$\Rightarrow \mathbb{E}[\phi_G(x)\phi_G(y)] = -\Delta^{-1}(x, y) \sim \ln \frac{1}{|x-y|}$$

$\Rightarrow \phi_G$ is a GFF!

What is Liouville field theory?

Path integral formalism

$$\Sigma = \{X : \mathbb{D} \rightarrow \mathbb{R}\}$$

For $X \in \Sigma$, energy of X := $\frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2 + \int_{\partial\mathbb{D}} e^{\frac{\gamma}{2} X} ds$

Random field ϕ_L :

$$\mathbb{E}[F(\phi_L)] = \int_{\Sigma} F(X) e^{-\frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2 - \int_{\partial\mathbb{D}} e^{\frac{\gamma}{2} X} ds} DX$$

with $\gamma \in (0, 2)$.

$\Rightarrow \phi_L$ is the **Liouville field**

Correlations of Liouville theory

- Liouville theory is a conformal field theory
- Correlation function of $z_i \in \mathbb{D}$, $\alpha_i \in \mathbb{R}$:

$$\left\langle \prod_{i=1}^N e^{\alpha_i \phi_L(z_i)} \right\rangle_{\mathbb{D}} = \int_{\Sigma} DX \prod_{i=1}^N e^{\alpha_i X(z_i)} e^{-\frac{1}{4\pi} \int_{\mathbb{D}} |\partial X|^2 dx^2 - \int_{\partial \mathbb{D}} e^{\frac{\gamma}{2} X} ds}$$

- Expressed as moments of the Gaussian multiplicative chaos

Correlations of Liouville theory

- Degenerate fields: $e^{-\frac{\gamma}{2}\phi_L(z)}$ and $e^{-\frac{2}{\gamma}\phi_L(z)}$.
- BPZ equation, for $z_1, z \in \mathbb{D}$, $\alpha \in \mathbb{R}$, $\gamma \in (0, 2)$:

$$(z_1, z) \mapsto \langle e^{-\frac{\gamma}{2}\phi_L(z)} e^{\alpha\phi_L(z_1)} \rangle_{\mathbb{D}}$$

is solution of a differential equation.

- $\langle e^{-\frac{\gamma}{2}\phi_L(z)} e^{\alpha\phi_L(z_1)} \rangle_{\mathbb{D}} = \tilde{C} t^{\frac{\alpha\gamma}{2}} (1 - t^2)^{-\frac{\gamma^2}{8}} G(\gamma, p, t)$

BPZ equation on the upper half plane \mathbb{H}

Proposition (Remy 2017)

Let $\gamma \in (0, 2)$ and $\alpha > Q + \frac{\gamma}{2}$. Then:

$$\begin{aligned} & \left(\frac{4}{\gamma^2} \partial_{zz} + \frac{\Delta_{-\frac{\gamma}{2}}}{(z - \bar{z})^2} + \frac{\Delta_\alpha}{(z - z_1)^2} + \frac{\Delta_\alpha}{(z - \bar{z}_1)^2} + \frac{1}{z - \bar{z}} \partial_{\bar{z}} \right. \\ & \left. + \frac{1}{z - z_1} \partial_{z_1} + \frac{1}{z - \bar{z}_1} \partial_{\bar{z}_1} \right) \langle e^{-\frac{\gamma}{2}\phi_L(z)} e^{\alpha\phi_L(z_1)} \rangle_{\mathbb{H}} = 0 \end{aligned}$$

where $Q = \frac{\gamma}{2} + \frac{2}{\gamma}$, $\Delta_\alpha = \frac{\alpha}{2}(Q - \frac{\alpha}{2})$, $\Delta_{-\frac{\gamma}{2}} = -\frac{\gamma}{4}(Q + \frac{\gamma}{4})$.

\Rightarrow differential equation for $G(p, \gamma, t)$.

Outlook and perspectives

Work in progress, analogue for the bulk measure.

For $\gamma \in (0, 2)$, $\alpha \in (\frac{\gamma}{2}, Q)$:

$$\begin{aligned} \mathbb{E}\left[\left(\int_{\mathbb{D}} \frac{1}{|x|^{\gamma\alpha}} e^{\gamma X(x)} dx^2\right)^{\frac{Q-\alpha}{\gamma}}\right] = \\ \gamma^2 \left(\pi \frac{\Gamma(\frac{\gamma^2}{4})}{\Gamma(1 - \frac{\gamma^2}{4})}\right)^{\frac{Q-\alpha}{\gamma}} \cos\left(\frac{\alpha - Q}{\gamma}\pi\right) \frac{\Gamma(\frac{2\alpha}{\gamma} - \frac{4}{\gamma^2})\Gamma(\frac{\gamma}{2}(\alpha - Q))}{\Gamma(\frac{\alpha - Q}{\gamma})} \end{aligned}$$

Outlook and perspectives

Work in progress, analogue for the bulk measure.

For $\gamma \in (0, 2)$, $\alpha \in (\frac{\gamma}{2}, Q)$:

$$\mathbb{E}\left[\left(\int_{\mathbb{D}} \frac{1}{|x|^{\gamma\alpha}} e^{\gamma X(x)} dx^2\right)^{\frac{Q-\alpha}{\gamma}}\right] = \\ \gamma^2 \left(\pi \frac{\Gamma(\frac{\gamma^2}{4})}{\Gamma(1 - \frac{\gamma^2}{4})}\right)^{\frac{Q-\alpha}{\gamma}} \cos\left(\frac{\alpha - Q}{\gamma}\pi\right) \frac{\Gamma(\frac{2\alpha}{\gamma} - \frac{4}{\gamma^2})\Gamma(\frac{\gamma}{2}(\alpha - Q))}{\Gamma(\frac{\alpha - Q}{\gamma})}$$

Liouville theory with action: $\int_{\mathbb{D}} (|\partial X|^2 + e^{\gamma X}) dx^2$

Outlook and perspectives

Integrability program for GMC and Liouville theory:

- GMC on the unit interval $[0, 1]$
- More general Liouville correlations
- Other geometries

Link with planar maps

Conformal bootstrap

Thank you!

- Kupiainen A., Rhodes R., Vargas V.: Local conformal structure of Liouville quantum gravity, arXiv:1512.01802.
- Kupiainen A., Rhodes R., Vargas V.: Integrability of Liouville theory: Proof of the DOZZ formula, arXiv:1707.08785.
- Remy G.: The Fyodorov-Bouchaud formula and Liouville conformal field theory, arXiv:1710.06897.