




Contents lists available at ScienceDirect

Advances in Mathematics

journal homepage: www.elsevier.com/locate/aim

Hecke algebras for p -adic reductive groups and Local Langlands Correspondences for Bernstein blocks

Anne-Marie Aubert^a, Yujie Xu^{b,*}^a Sorbonne Université and Université Paris Cité, CNRS, IMJ-PRG, F-75005 Paris, France^b M.I.T., 77 Massachusetts Avenue, Cambridge, MA, USA

ARTICLE INFO

Article history:

Received 6 December 2022

Received in revised form 22

September 2023

Accepted 26 October 2023

Available online 17 November 2023

Communicated by Pramod Achar

*Keywords:*Representation of p -adic group

Hecke algebra

Bernstein center

Local Langlands Correspondence

ABSTRACT

We study the endomorphism algebras attached to Bernstein components of reductive p -adic groups and construct a local Langlands correspondence with the appropriate set of enhanced L -parameters, using certain “desiderata” properties for the LLC for supercuspidal representations of proper Levi subgroups. We give several applications of our LLC to various reductive groups with Bernstein blocks cuspidally supported on general linear groups.

In particular, for Levi subgroups of maximal parabolic subgroups of the split exceptional group G_2 , we compute the explicit weight functions for the corresponding Hecke algebras, and show that they satisfy a conjecture of Lusztig’s. Some results from §4 are used by the same authors to construct a full local Langlands correspondence in [9]. Moreover, we prove a “reduction to depth-zero” result for regular Bernstein blocks (i.e., blocks for which the supercuspidal support of each irreducible representation is regular).

© 2023 Elsevier Inc. All rights reserved.

* Corresponding author.

E-mail addresses: anne-marie.aubert@imj-prg.fr (A.-M. Aubert), yujiexu@mit.edu (Y. Xu).

Contents

1.	Introduction	2
2.	Hecke algebras and Bernstein center	7
2.1.	General framework	7
2.2.	Bernstein blocks	16
3.	Local Langlands correspondence for Bernstein blocks	20
3.1.	Axiomatic construction of the correspondence	20
3.2.	Matching of simple modules of extended affine Hecke algebras	26
4.	Applications to G_2	28
4.1.	Explicit Hecke algebra parameters	32
4.2.	Intertwining algebras	37
5.	Applications to other groups	43
References		44

1. Introduction

1.1. Background

Let F be a non-archimedean local field. Let $\mathbf{G}$ be a connected reductive group defined over F , and G its group of F -points. Let M be a Levi subgroup of a parabolic subgroup P of G .

Let $\mathfrak{s} = [M, \sigma]_G$ be the inertial class attached to the pair (M, σ) , where σ is an irreducible supercuspidal representation of M . Recall that this means that $\mathfrak{s}$ is the G -conjugacy class of $(M, \mathfrak{X}_{\text{nr}}(M) \cdot \sigma)$, where $\mathfrak{X}_{\text{nr}}(M) \cdot \sigma$ is the orbit of σ under $\mathfrak{X}_{\text{nr}}(M)$ —the group of unramified characters of M . Let $\mathfrak{B}(G)$ be the set of such $\mathfrak{s}$. We denote by $\text{Irr}^{\mathfrak{s}}(G)$ the Bernstein series of irreducible representations of G whose cuspidal support lies in $\mathfrak{s}$ (see §2.1 for the precise definition).

Let $W_G^{\mathfrak{s}}$ denote the extended finite Weyl group $N_G(\mathfrak{s}_M)/M$, where $\mathfrak{s}_M = [M, \sigma]_M$, and let $W_G^{\mathfrak{s}, x}$ be the stabilizer of $x \in \text{Irr}^{\mathfrak{s}_M}(M)$ in $W_G^{\mathfrak{s}}$. By [54], there exists a collection $(\natural_x)_x$ of 2-cocycles for $x \in \text{Irr}^{\mathfrak{s}_M}(M)$,

$$\natural_x : W_G^{\mathfrak{s}, x} \times W_G^{\mathfrak{s}, x} \longrightarrow \mathbb{C}^{\times}, \quad (1.1.1)$$

such that we have a bijection

$$\xi_G^{\mathfrak{s}} : \text{Irr}^{\mathfrak{s}}(G) \longrightarrow (\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\natural}, \quad (1.1.2)$$

where $(\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\natural}$ is a *twisted extended quotient* in the sense of [1, §2.1] (see (1.3.1) for the precise definition).

A parallel picture to (1.1.2) exists on the Galois side. Let W_F be the absolute Weil group of F and I_F its inertia subgroup. Let $M^{\vee}$ be the Langlands dual group of M , i.e. it is a complex reductive group with root datum dual to that of M . It is equipped

with an action of W_F , and we write ${}^L M := M^\vee \rtimes W_F$. The group $M^\vee$ acts on the set of *cuspidal* M -relevant enhanced L -parameters for M —a terminology based on Lusztig’s notion of cuspidal pairs (see Definition 3.1.8 for more details). Let $\Phi_e^c(M)$ denote the set of $M^\vee$ -conjugacy classes of cuspidal enhanced L -parameters for M .

Let $Z_{M^\vee \rtimes I_F}$ be the center of $M^\vee \rtimes I_F$. The group $\mathfrak{X}_{\text{nr}}({}^L M) := (Z_{M^\vee \rtimes I_F})_{W_F}^\circ$, which is naturally isomorphic to the group $\mathfrak{X}_{\text{nr}}(M)$ (see [25, §3.3.1]), acts naturally on the set of cuspidal M -relevant enhanced L -parameters for M . We denote by $\mathfrak{s}^\vee = [{}^L M, \varphi_c, \varrho_c]_{G^\vee}$ the $G^\vee$ -conjugacy class of the orbit of $(\varphi_c, \varrho_c) \in \Phi_e^c(M)$ under the action of $\mathfrak{X}_{\text{nr}}({}^L M)$. Let $\mathfrak{B}^\vee(G)$ be the set of such $\mathfrak{s}^\vee$.

In [7], the first author, with Moussaoui and Solleveld, constructed a partition-à la Bernstein—of the set $\Phi_e(G)$ of G -relevant enhanced Langlands parameters for G :

$$\Phi_e(G) = \bigsqcup_{\mathfrak{s}^\vee \in \mathfrak{B}^\vee(G)} \Phi_e^{\mathfrak{s}^\vee}(G), \quad (1.1.3)$$

where $\Phi_e^{\mathfrak{s}^\vee}(G)$ consists of the enhanced Langlands parameters for G whose cuspidal support lies in $\mathfrak{s}^\vee$. Let $\mathfrak{s}_M^\vee := [{}^L M, \varphi_c, \varrho_c]_{M^\vee} \in \mathfrak{B}^\vee(M)$. Analogous to the group side $W_G^{\mathfrak{s}}$, we denote by $W_{G^\vee}^{\mathfrak{s}_M^\vee}$ the stabilizer of $\mathfrak{s}_M^\vee$ in $N_{G^\vee}(M^\vee)/M^\vee$, and by $W_{G^\vee}^{\mathfrak{s}_M^\vee, y}$ the stabilizer of $y \in \Phi_e^{\mathfrak{s}_M^\vee}(M)$ in $W_{G^\vee}^{\mathfrak{s}_M^\vee}$. By [7, Theorem 9.3], there is a bijection

$$\xi_{G^\vee}^{\mathfrak{s}_M^\vee} : \Phi_e^{\mathfrak{s}_M^\vee}(G) \longrightarrow (\Phi_e^{\mathfrak{s}_M^\vee}(M) // W_{G^\vee}^{\mathfrak{s}_M^\vee})_{L_{\mathfrak{h}}}, \quad (1.1.4)$$

where the right-hand side $(\Phi_e^{\mathfrak{s}_M^\vee}(M) // W_{G^\vee}^{\mathfrak{s}_M^\vee})_{L_{\mathfrak{h}}}$ is a twisted extended quotient with respect to a collection $(L_{\mathfrak{h}_y})_y$ of 2-cocycles

$$L_{\mathfrak{h}_y} : W_{G^\vee}^{\mathfrak{s}_M^\vee, y} \times W_{G^\vee}^{\mathfrak{s}_M^\vee, y} \rightarrow \mathbb{C}^\times. \quad (1.1.5)$$

1.2. Main results

Axiomatic setup: we suppose the existence of a map

$$\begin{array}{ccc} \mathfrak{L}^{\mathfrak{s}_M} : & \text{Irr}^{\mathfrak{s}_M}(M) & \longrightarrow \Phi_e^c(M) \\ & \sigma & \mapsto (\varphi_\sigma, \varrho_\sigma) \end{array} \quad (1.2.1)$$

such that the following properties are satisfied for any $\sigma \in \text{Irr}^{\mathfrak{s}_M}(M)$:

(1) For any $\chi \in \mathfrak{X}_{\text{nr}}(M)$, we have

$$(\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}) = \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma),$$

where $\chi \mapsto \chi^\vee$ is the canonical isomorphism $\mathfrak{X}_{\text{nr}}(M) \xrightarrow{\sim} \mathfrak{X}_{\text{nr}}({}^L M)$.

(2) For any $w \in W(M)$, we have

$${}^{w^\vee}(\varphi_\sigma, \varrho_\sigma) \simeq (\varphi_{w^\vee \sigma}, \varrho_{w^\vee \sigma}),$$

where $w \mapsto w^\vee$ is the canonical isomorphism $W(M) \xrightarrow{\sim} W(M^\vee)$.

We suppose that the collections of 2-cocycles $\natural$ and ${}^L \natural$ satisfy the following

$${}^L \natural_{\chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)} = \natural_{\chi \otimes \sigma} \quad \text{for any } \sigma \in \mathfrak{s} \text{ and any } \chi \in \mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma), \quad (1.2.2)$$

where $\mathfrak{X}_{\text{nr}}(M, \sigma) := \{\chi \in \mathfrak{X}_{\text{nr}}(M) : \chi \otimes \sigma \simeq \sigma\}$. We establish the following result.

Theorem 1. (Theorem 3.1.32)

(1) *There is a natural isomorphism*

$$\mathfrak{e}: \text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}} \xrightarrow{\sim} \Phi_e^{\mathfrak{s}_M^\vee}(M) // W_{G^\vee}^{\mathfrak{s}}. \quad (1.2.3)$$

(2) *The map*

$$\mathfrak{L} := (\xi_{G^\vee}^{\mathfrak{s}})^{-1} \circ \mathfrak{e} \circ \xi_G^{\mathfrak{s}}: \text{Irr}^{\mathfrak{s}}(G) \longrightarrow \Phi_e^{\mathfrak{s}^\vee}(G) \quad (1.2.4)$$

is a bijection.

We suppose in the rest of this introduction that the group $\mathbf{G}$ splits over a tamely ramified extension of F and that the residual characteristic p of F does not divide the order of the Weyl group of G . Then there exists a compact mod center subgroup $\tilde{K}_M$ of M and an irreducible representation ρ_M^d of it such that $\sigma = \text{ind}_{\tilde{K}_M}^M \rho_M^d$.

Let $\mathcal{H}^{\mathfrak{s}}(G)$ denote the endomorphism algebra of the Bernstein progenerator of $\mathfrak{s}$ (see (2.1.5)) and let $\mathcal{H}(G, \rho_{\mathcal{D}})$ be the intertwining algebra of an $\mathfrak{s}$ -type $(K_{\mathcal{D}}, \rho_{\mathcal{D}})$. We prove in Proposition 2.1.58 that the algebras $\mathcal{H}^{\mathfrak{s}}(G)$ and $\mathcal{H}(G, \rho_{\mathcal{D}})$ are isomorphic.

From now on, we suppose that σ is *regular* in the sense of [31], which allows us to attach a supercuspidal Langlands parameter $\varphi_\sigma: W_F \rightarrow {}^L M$ to σ . Applying Theorem 1 to the map $\mathfrak{L}^{\mathfrak{s}_M}: \sigma \mapsto (\varphi_\sigma, 1)$ as in (1.2.1), we obtain the following result:

Proposition 1.2.5. *When the L -packet of σ is a singleton, the properties (3.1.23) and (3.1.23) are always satisfied.*

On the other hand, the construction of $\tilde{K}_M$ involves notably a depth zero supercuspidal irreducible representation σ^0 of a Levi subgroup M^0 of a twisted Levi subgroup G^0 of G . We denote by $\mathfrak{s}^0 = [M^0, \sigma^0]_{G^0}$ the inertial class of σ^0 .

Suppose that p is good for G (in the sense of [19]) and does not divide the order of the fundamental group of $\mathbf{G}_{\text{der}}$, and that the representation σ is regular.

Theorem 2. (Theorem 2.2.14) *There is a bijection*

$$\varrho_{s^0}^s : \text{Irr}^s(G) \longrightarrow \text{Irr}^{s^0}(G^0), \quad (1.2.6)$$

which induces a bijection

$$\text{Irr}(\mathcal{H}^s(G)) \longrightarrow \text{Irr}(\mathcal{H}^{s^0}(G^0)) \quad (1.2.7)$$

between the sets of equivalence classes of simple modules for the algebras $\mathcal{H}^s(G)$ and $\mathcal{H}^{s^0}(G^0)$.

Theorem 2 proves the validity of [5, Conjecture 1.1], under the above assumption on p , for all regular supercuspidal representations of M . The bijection $\varrho_{s^0}^s$ is defined as

$$\varrho_{s^0}^s := (\xi_{G^0}^{s^0})^{-1} \circ \mathfrak{l}_\sigma \circ \xi_G^s, \quad (1.2.8)$$

where $(\text{Irr}^{s_{M^0}}(M^0) // W_{G^0}^{s^0})_{\natural^0}$ is the twisted extended quotient with respect to a certain collection $\natural^0$ of 2-cocycles, the definition of which is recalled in (2.2.13), and

$$\mathfrak{l}_\sigma : (\text{Irr}^{s_M}(M) // W_G^s)_{\natural} \longrightarrow (\text{Irr}^{s_{M^0}}(M^0) // W_{G^0}^{s^0})_{\natural^0} \quad (1.2.9)$$

is the isomorphism constructed in [5].

In Section 4, we study in greater detail the case when G is the exceptional group of type G_2 . Recall that for split p -adic groups, the principal series case, i.e. $M = T$, is done in [46] and [3], therefore it suffices to consider the cases where $M \simeq \text{GL}_2(F)$ is a maximal Levi subgroup. The $G_2(F)$ -covers of the supercuspidal types in M were computed explicitly in [14] when M corresponds to the long simple root of G_2 , and in [20] when M corresponds to the short simple root of G_2 , but the intertwining algebras of these types were still unknown. We compute these intertwining algebras later in §4.2.2, and in particular, by computing their parameters explicitly, we show that they satisfy a conjecture of Lusztig's in [40, 1.(a)].

Acknowledgments

The authors would like to thank Maarten Solleveld and the anonymous referee for valuable comments on a previous version of the manuscript. Y.X. was supported by the National Science Foundation under Award No. 2202677 at MIT.

1.3. Notations and definitions

Let F be a non-archimedean local field. Let $\mathfrak{o}_F$ denote the ring of integers of F , $\mathfrak{p}_F$ the maximal ideal in $\mathfrak{o}_F$ and $k_F := \mathfrak{o}_F / \mathfrak{p}_F$ the residue field of F . We denote by q_F the

cardinality of k_F . Let $\text{val}_F: F \rightarrow \mathbb{Z} \cup \{\infty\}$ be a valuation of F and let ν_F the character of $F^\times$ defined by $\nu_F(a) := q_F^{-\text{val}_F(a)}$ for any $a \in F^\times$.

We fix a separable closure F_{sep} of F . Denote by $W_F \subset \text{Gal}(F_{\text{sep}}/F)$ the absolute Weil group of F and I_F its inertia subgroup. We denote by F_{nr} the maximal unramified extension of F inside F_{sep} and by Fr_F the element of $\text{Gal}(F_{\text{nr}}/F)$ that induces the automorphism $a \mapsto a^q$ on the residue field $\bar{k}_F$ of F_{nr} . Then $W_F = I_F \rtimes \langle \text{Fr}_F \rangle$. Let I_F^+ denote the wild inertia group of F (i.e. the maximal pro- p open normal subgroup of I_F). We have $I_F^t = \text{Gal}(F_{\text{sep}}/F_t) \simeq I_F/I_F^+$, where F_t is the tame closure of F in F_{sep} . The group I_F^t is pro-cyclic and we denote by ζ_F a generator of it. Let $W'_F := W_F \times \text{SL}_2(\mathbb{C})$ be the Weil-Deligne group of F .

Let $\mathbf{G}$ be a connected reductive algebraic group defined over F , and $G := \mathbf{G}(F)$ its F -rational points. We denote by $\mathbf{G}_{\text{der}}$ the derived group of $\mathbf{G}$. Let $\mathbf{G}_{\text{sc}}$ (resp. $\mathbf{G}_{\text{ad}}$) be the simply connected cover (resp. adjoint quotient) of $\mathbf{G}_{\text{der}}$. Let $Z_{\mathbf{G}}$ be the center of $\mathbf{G}$, and $\mathbf{A}_{\mathbf{G}}$ the maximal F -split torus contained in $Z_{\mathbf{G}}$.

Fix a maximal torus $\mathbf{T}$ of $\mathbf{G}$, and let $(X, R, Y, R^\vee)$ denote the root datum of $\mathbf{G}$ with respect to $\mathbf{T}$. Thus $X = X^*(\mathbf{T})$ is the character group of $\mathbf{T}$, and $R \subset X$ is the set of weights of $\mathbf{T}$ on the Lie algebra $\mathfrak{g}$ of $\mathbf{G}$. Fix $\Delta \subset R$ a system of simple roots. When R is irreducible, the root with maximal height (with respect to Δ) will be denoted $\tilde{\alpha}$. Write $\tilde{\alpha} = \sum_{\gamma \in \Delta} c_\gamma \gamma$ for positive integers c_γ . A prime number p is said to be *good for $\mathbf{G}$* if it does not divide any c_γ . We may simply list the *bad*, i.e. not good, primes: $p = 2$ is bad unless R is of type A , $p = 3$ is bad if R is of type G_2, F_4, E_n , and $p = 5$ is bad if R is of type E_8 . The prime p is good for a general R just in case it is good for each irreducible component of R .

Suppose that H is a group, H_1 a subgroup of H and h an element of H . We set ${}^h H_1 := hH_1h^{-1}$. If π is a representation of H_1 , we denote by ${}^h \pi$ the representation $h_1 \mapsto \pi(h^{-1}h_1h)$ of ${}^h H_1$. We denote by $\text{Irr}(H)$ the set of equivalence classes of irreducible representations of H .

The category of right modules over an algebra $\mathcal{A}$ is denoted $\mathcal{A}\text{-Mod}$. We write $\text{Irr}(\mathcal{A})$ for the set of equivalence classes of simple modules of $\mathcal{A}$.

1.3.1. Twisted extended quotients

Let Γ be a group acting on a topological space X and let Γ_x denote the stabilizer in Γ of $x \in X$. Let $\natural = (\natural_x)_{x \in X}$ be a collection of 2-cocycles

$$\natural_x: \Gamma_x \times \Gamma_x \rightarrow \mathbb{C}^\times,$$

such that $\natural_{\gamma x}$ and $\gamma_* \natural_x$ define the same class in $H^2(\Gamma_{\gamma x}, \mathbb{C}^\times)$, where $\gamma_*: \Gamma_x \rightarrow \Gamma_{\gamma x}$ sends α to $\gamma\alpha\gamma^{-1}$. Let $\mathbb{C}[\Gamma_x, \natural_x]$ be the group algebra of Γ_x twisted by $\natural_x$. We set

$$\widetilde{X}_{\natural} := \{(x, \tau) : x \in X, \tau \in \text{Irr } \mathbb{C}[\Gamma_x, \natural_x]\},$$

and topologize $\widetilde{X}_{\natural}$ by decreeing that a subset of $\widetilde{X}_{\natural}$ open if its projection to the first coordinate is open in X .

We require, for every $(\gamma, x) \in \Gamma \times X$, an algebra isomorphism

$$\phi_{\gamma,x}: \mathbb{C}[\Gamma_x, \mathfrak{h}_x] \rightarrow \mathbb{C}[\Gamma_{\gamma x}, \mathfrak{h}_{\gamma x}]$$

satisfying the conditions

- (a) if $\gamma x = x$, then $\phi_{\gamma,x}$ is conjugation by an element of $\mathbb{C}[\Gamma_x, \mathfrak{h}_x]^\times$;
- (b) $\phi_{\gamma',\gamma x} \circ \phi_{\gamma,x} = \phi_{\gamma'\gamma,x}$ for all $\gamma', \gamma \in \Gamma$ and $x \in X$.

Define a Γ -action on $\widetilde{X}_{\mathfrak{h}}$ by $\gamma \cdot (x, \tau) := (\gamma x, \tau \circ \phi_{\gamma,x}^{-1})$. The *spectral twisted extended quotient* of X by Γ with respect to $\mathfrak{h}$ is defined to be

$$(X//\Gamma)_{\mathfrak{h}} := \widetilde{X}_{\mathfrak{h}}/\Gamma. \quad (1.3.1)$$

In the case when the 2-cocycles $\mathfrak{h}_x$ are trivial, we write simply $X//\Gamma$ for $(X//\Gamma)_{\mathfrak{h}}$ and refer to it as the *spectral extended quotient* of X by Γ .

2. Hecke algebras and Bernstein center

2.1. General framework

2.1.1. Let $\mathfrak{R}(G)$ denote the category of all smooth complex representations of G . It is an abelian category admitting arbitrary coproducts. Let M be a Levi subgroup of a parabolic subgroup P of G . We denote by M_1 the subgroup of M generated by all its compact subgroups. Recall that a character of M is said to be unramified if it is trivial on M_1 , and let $\mathfrak{X}_{\text{nr}}(M)$ be the group of unramified characters of M . Let σ be an irreducible supercuspidal smooth representation of M . We write $\mathfrak{s} := [M, \sigma]_G$ for the G -conjugacy class of the pair $(M, \mathfrak{X}_{\text{nr}}(M) \cdot \sigma)$, it is called a *Bernstein inertial class*. Let $\mathfrak{B}(G)$ denote the set of Bernstein inertial classes $\mathfrak{s}$. We set $\mathfrak{s}_M := [M, \sigma]_M$.

We denote by $\mathfrak{R}^{\mathfrak{s}}(G)$ the full subcategory of $\mathfrak{R}(G)$ whose objects are the representations (π, V) such that every irreducible G -subquotient of π is equivalent to a subquotient of a parabolically induced representation $i_P^G(\sigma')$, where i_P^G is the functor of normalized parabolic induction and $\sigma' \in \mathfrak{X}_{\text{nr}}(M) \cdot \sigma$. We write $\text{Irr}^{\mathfrak{s}}(G)$ for the class of irreducible objects in $\mathfrak{R}^{\mathfrak{s}}(G)$, i.e. representations whose supercuspidal support lies in $\mathfrak{s}$.

2.1.2. The categories $\mathfrak{R}^{\mathfrak{s}}(G)$ are indecomposable and split the full smooth category $\mathfrak{R}(G)$ in a direct product (see [11, Proposition 2.10]):

$$\mathfrak{R}(G) = \prod_{\mathfrak{s} \in \mathfrak{B}(G)} \mathfrak{R}^{\mathfrak{s}}(G).$$

If $\Pi^{\mathfrak{s}}$ is a progenerator of $\mathfrak{R}^{\mathfrak{s}}(G)$, then the functor $V \mapsto \text{Hom}_G(\Pi^{\mathfrak{s}}, V)$ is an equivalence from $\mathfrak{R}^{\mathfrak{s}}(G)$ to the algebra $\text{End}_G(\Pi^{\mathfrak{s}})$ (see for instance [47, § 1.1]).

Let $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$ and let V be the underlying vector space for the supercuspidal representation σ of M and σ_1 an irreducible component of the restriction of σ to M_1 . We denote by $\text{ind}_{M_1}^M$ the functor of compact induction. As noticed in [47, § 1.2], the isomorphism class of

$$\Pi_M^{\mathfrak{s}_M} := \text{ind}_{M_1}^M(\sigma_1) \quad (2.1.3)$$

is independent of the choice of σ_1 . It was shown by Bernstein that

$$\Pi_G^{\mathfrak{s}} := i_P^G(\Pi_M^{\mathfrak{s}_M}) \quad (2.1.4)$$

is a progenerator of $\mathfrak{R}^{\mathfrak{s}}(G)$ (see [47, §1.6]). We write

$$\mathcal{H}^{\mathfrak{s}}(G) := \text{End}_G(\Pi_G^{\mathfrak{s}}). \quad (2.1.5)$$

Hence we have an equivalence of categories of right modules

$$\mathfrak{R}^{\mathfrak{s}}(G) \simeq \mathcal{H}^{\mathfrak{s}}(G) - \text{Mod}. \quad (2.1.6)$$

Let $B := \mathbb{C}[M/M_1]$ and $V_B := V \otimes_{\mathbb{C}} B$. Then $i_P^G(V_B)$ is also a progenerator of $\mathfrak{R}^{\mathfrak{s}}(G)$, and we have an equivalence of categories of right modules given by

$$\begin{aligned} \mathcal{E}: \quad \mathfrak{R}^{\mathfrak{s}}(G) &\longrightarrow \text{End}_G(i_P^G(V_B)) - \text{Mod} \\ \mathcal{V} &\longmapsto \text{Hom}_G(i_P^G(V_B), \mathcal{V}). \end{aligned} \quad (2.1.7)$$

2.1.8. Consider

$$\mathfrak{X}_{\text{nr}}(M, \sigma) := \{\chi \in \mathfrak{X}_{\text{nr}}(M) : \chi \otimes \sigma \simeq \sigma\}, \quad (2.1.9)$$

which is a finite subgroup of $\mathfrak{X}_{\text{nr}}(M)$.

Remark 2.1.10. In the case where $M = \text{GL}_n(F)$ with n a positive integer, there is a simple type (J, λ) in the sense of [12, (5.5.10)] such that the restriction of the supercuspidal representation σ to J contains λ . The order of $\mathfrak{X}_{\text{nr}}(M, \sigma)$ is $n/e(L|F)$, where $e(L|F)$ is the ramification index of the extension L/F involved in the definition of (J, λ) (see [12, (6.0.1) and (6.2.5)]).

We denote by $\mathcal{O}$ the orbit of σ under the action of $\mathfrak{X}_{\text{nr}}(M)$. The map $\chi \mapsto \chi \otimes \sigma$ defines a bijection

$$\mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma) \xrightarrow{\sim} \mathcal{O} = \{\chi \otimes \sigma : \chi \in \mathfrak{X}_{\text{nr}}(M)\} = \text{Irr}^{\mathfrak{s}_M}(M). \quad (2.1.11)$$

We set $W(M) := N_G(M)/M$, where $N_G(M)$ denotes the normalizer of M in G , and define

$$W_G^s := W(M, \mathcal{O}) := \{n \in N_G(M) : {}^n\mathcal{O} \simeq \mathcal{O}\} / M. \quad (2.1.12)$$

Recall that $\mathbf{A}_M$ is the maximal split torus contained in the center of $\mathbf{M}$. We denote by $\Sigma(A_M) \subset X^*(A_M)$ the set of nonzero weights occurring in the adjoint representation of A_M on the Lie algebra of G , and by $\Sigma_{\text{red}}(A_M)$ be the set of indivisible elements therein. (Recall that a root γ in a root system Σ is called *indivisible* if $\frac{1}{2}\gamma \notin \Sigma$.)

For every $\gamma \in \Sigma_{\text{red}}(A_M)$, let $M_\gamma \supset M$ denote the centralizer of $\ker \gamma$ in G (it is a Levi subgroup of G whose semisimple rank is one larger than that of M). Let μ^G be the Harish-Chandra μ -function for G (see [52, §1] or [55, §V.2]). The restriction of μ^G to $\mathcal{O}$ is a rational $W(M, \mathcal{O})$ -invariant function on $\mathcal{O}$ [55, Lemma V.2.1]. By [26, Proposition 1.3], the set

$$\Sigma_{\mathcal{O}, \mu} := \{\gamma \in \Sigma_{\text{red}}(A_M) : \mu^{M_\gamma} \text{ has a zero on } \mathcal{O}\} \quad (2.1.13)$$

is a root system. Let $W_{\mathcal{O}}$ denote the Weyl group of $\Sigma_{\mathcal{O}, \mu}$.

Let $P = MN$ be a parabolic subgroup of G with Levi factor M . Denote by $\Sigma(P)$ the subset of $\Sigma(A_M)$ of roots which act on the Lie algebra of N . Let $\Sigma_{\mathcal{O}, \mu}(P) := \Sigma_{\mathcal{O}, \mu} \cap \Sigma(P)$. By [26, 1.12], the group $W(M, \mathcal{O})$ decomposes as

$$W(M, \mathcal{O}) = W_{\mathcal{O}} \rtimes R(\mathcal{O}), \quad (2.1.14)$$

where

$$R(\mathcal{O}) := \{w \in W(M, \mathcal{O}) : w(\Sigma_{\mathcal{O}, \mu}(P)) = \Sigma_{\mathcal{O}, \mu}(P)\}. \quad (2.1.15)$$

The action of every element w of W_G^s can be lifted to a transformation $\tilde{w}$ of $\mathfrak{X}_{\text{nr}}(M)$. Let $W(M, \sigma, \mathfrak{X}_{\text{nr}}(M))$ be the group of permutations of $\mathfrak{X}_{\text{nr}}(M)$ generated by $\mathfrak{X}_{\text{nr}}(M, \sigma)$ and the $\tilde{w}$'s. We have

$$W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)) / \mathfrak{X}_{\text{nr}}(M, \sigma) \simeq W_G^s. \quad (2.1.16)$$

Let $K(B) := \mathbb{C}(\mathfrak{X}_{\text{nr}}(M))$ denote the quotient field of $B := \mathbb{C}[\mathfrak{X}_{\text{nr}}(M)]$. Let

$$\mathbb{C}[W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)), \kappa]$$

be the twisted group algebra of $W(M, \sigma, \mathfrak{X}_{\text{nr}}(M))$ with basis elements t_w that multiply as $t_w t_{w'} = \kappa(w, w') t_{ww'}$. By [54, Corollary 5.8], there is a 2-cocycle

$$\kappa : W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)) \times W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)) \rightarrow \mathbb{C}^\times, \quad (2.1.17)$$

such that we have an algebra isomorphism

$$K(B) \otimes_B \text{End}_G(i_P^G(V_B)) \simeq K(B) \rtimes \mathbb{C}[W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)), \kappa]. \quad (2.1.18)$$

Here the symbol $\rtimes$ denotes the crossed product: as a vector space, it just means the tensor product, with multiplication rules determined by the action of $W(M, \sigma, \mathfrak{X}_{\text{nr}}(M))$ on $K(B)$. Note that the cocycle κ is trivial on $W_{\mathcal{O}}$.

Remark 2.1.19. If $R(\mathcal{O})$ has order at most 2, the intertwining operators can be normalized such that the cocycle κ is trivial (see [54, Proposition 5.2 & above Lemma 5.7]). This is indeed the case for $G = G_2(F)$.

For any $\chi \in \mathfrak{X}_{\text{nr}}(M)$, let $W_G^{\mathfrak{s}, \chi \otimes \sigma}$ denote the stabilizer of $\chi \otimes \sigma$ in $W_G^{\mathfrak{s}}$. Let $\mathfrak{d}_{\chi}$ be the 2-cocycle denoted $\mathfrak{d}_{\sigma'}$ in [54, (9.13)]. Let $(\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{d}}$ denote the twisted extended quotient (as in § 1.3.1) with respect to the collection $\mathfrak{d}$ of the 2-cocycles $\mathfrak{d}_{\chi}$.

Proposition 2.1.20. *There is a bijection*

$$\xi_G^{\mathfrak{s}}: \text{Irr}^{\mathfrak{s}}(G) \longrightarrow (\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{d}}. \quad (2.1.21)$$

Proof. By [54, Theorem 9.7], there are bijections

$$\text{Irr}^{\mathfrak{s}}(G) \xleftarrow{\mathcal{E}} \text{Irr}(\text{End}_G(i_P^G(V_B))) \xleftarrow{\zeta} \text{Irr}(\mathbb{C}[\mathfrak{X}_{\text{nr}}(M)] \rtimes \mathbb{C}[W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)), \kappa],$$

where $\mathcal{E}$ is induced by the equivalence of categories defined in (2.1.7). On the other hand, by [54, Lemma 9.8], $\text{Irr}(\mathbb{C}[\mathfrak{X}_{\text{nr}}(M)] \rtimes \mathbb{C}[W(M, \sigma, \mathfrak{X}_{\text{nr}}(M)), \kappa]$ is canonically isomorphic to $(\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{d}}$, where $\mathfrak{s}_M := [M, \sigma]_M$. $\square$

Corollary 2.1.22. *Let $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$. There is a bijection*

$$\text{Irr}(\mathcal{H}^{\mathfrak{s}}(G)) \xrightarrow{1-1} (\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{d}}. \quad (2.1.23)$$

Proof. The result follows from the proof of Proposition 2.1.20 by using (2.1.6). $\square$

Remark 2.1.24. As observed in [54, (10.12)], if the restriction of σ to M_1 is multiplicity free, we have

$$\Pi_G^{\mathfrak{s}} = (i_P^G(V_B))^{\mathfrak{X}_{\text{nr}}(M, \sigma)} \quad \text{and} \quad \text{End}_G(i_P^G(V_B)) \simeq \mathcal{H}^{\mathfrak{s}}(G) \otimes_{\mathbb{C}} \text{Mat}_{[M: M_{\sigma}]}(\mathbb{C}), \quad (2.1.25)$$

where $\text{Mat}_{[M: M_{\sigma}]}(\mathbb{C})$ is the algebra of square matrices of size $[M: M_{\sigma}]$ (the index of M_{σ} in M) with entries in $\mathbb{C}$. Note that if σ is generic, then its restriction to M_1 is multiplicity free (see [48, Remark 1.6.1.3]). In particular, if σ is a supercuspidal irreducible representation of a proper Levi subgroup M of G_2 , since M is isomorphic to either $F^{\times} \times F^{\times}$ or $\text{GL}_2(F)$, the representation σ is generic, and hence its restriction to M_1 is multiplicity free.

2.1.1. Theory of types

We fix a Haar measure on G . Let $\mathcal{H}(G)$ be the space of locally constant, compactly supported functions $f: G \rightarrow \mathbb{C}$ and view $\mathcal{H}(G)$ as a $\mathbb{C}$ -algebra via convolution relative to the Haar measure. The algebra $\mathcal{H}(G)$ is called the *Hecke algebra of G* .

Let (ρ, V_ρ) be a smooth representation of a compact open subgroup K of G , and let $(\tilde{\rho}, V_{\tilde{\rho}})$ denote its contragredient. We define $\mathcal{H}(G, \rho)$ to be the space of compactly supported functions $f: G \rightarrow \text{End}_G(V_{\tilde{\rho}})$ such that

$$f(kgk') = \rho(k)f(g)\rho(k'), \quad \text{where } k, k' \in K \text{ and } g \in G. \quad (2.1.26)$$

The convolution product gives $\mathcal{H}(G, \rho)$ the structure of a unitary associative $\mathbb{C}$ -algebra. The algebra $\mathcal{H}(G, \rho)$ is called the *$\tilde{\rho}$ -spherical Hecke algebra* or the *intertwining algebra of (K, ρ)* .

Let $e_\rho \in \mathcal{H}(G)$ be the function defined by

$$e_\rho(g) := \begin{cases} \frac{\dim \rho}{\text{meas}(K)} \text{tr}(\rho(g^{-1})) & \text{if } g \in K, \\ 0 & \text{if } g \in G, g \notin K. \end{cases} \quad (2.1.27)$$

Then e_ρ is idempotent, and $e_\rho \star \mathcal{H}(G) \star e_\rho$ is a sub-algebra of $\mathcal{H}(G)$ with unit e_ρ . By [13, (2.12)], there is a canonical isomorphism

$$\mathcal{H}(G, \rho) \otimes_{\mathbb{C}} \text{End}_{\mathbb{C}}(V_\rho) \rightarrow e_\rho \star \mathcal{H}(G) \star e_\rho. \quad (2.1.28)$$

The algebras $\mathcal{H}(G, \rho)$ and $e_\rho \star \mathcal{H}(G) \star e_\rho$ are therefore canonically Morita equivalent. Hence, we get an equivalence of categories:

$$\mathcal{H}(G, \rho) - \text{Mod} \simeq e_\rho \star \mathcal{H}(G) \star e_\rho - \text{Mod}. \quad (2.1.29)$$

Let $\mathfrak{R}_\rho(G)$ be the full subcategory of $\mathfrak{R}(G)$ whose objects are those V satisfying $V = \mathcal{H}(G) \star e_\rho \star V$, i.e. $\mathfrak{R}_\rho(G)$ is generated over G by the subspace $e_\rho \star V$.

Definition 2.1.30.

- (1) The pair (K, ρ) is called an *$\mathfrak{s}$ -type for G* if the following holds: for any $\pi \in \text{Irr}(G)$, we have $\pi \in \mathfrak{R}^\mathfrak{s}(G)$ if and only if π contains ρ .
- (2) A *supercuspidal type for G* is an $\mathfrak{s}$ -type where $\mathfrak{s} = [G, \sigma]_G$.

If (K, ρ) is an $\mathfrak{s}$ -type for G , then $\mathfrak{R}_\rho(G) = \mathfrak{R}^\mathfrak{s}(G)$ by [13, (4.1)–(4.2)], where $\mathfrak{R}^\mathfrak{s}(G)$ is equivalent to the category of modules for $\mathcal{H}(G, \rho)$ by [13, Theorem 3.5]:

$$\mathfrak{R}^\mathfrak{s}(G) \simeq \mathcal{H}(G, \rho) - \text{Mod}. \quad (2.1.31)$$

Combining (2.1.31) and (2.1.6), we obtain an equivalence

$$\mathcal{H}^s(G)\text{-Mod} \simeq \mathcal{H}(G, \rho) \text{-Mod}. \quad (2.1.32)$$

Let (K_M, ρ_M) be an $\mathfrak{s}_M$ -type for $\mathfrak{s}_M \in \mathfrak{B}(M)$. If the pair (K, ρ) is a G -cover of (K_M, ρ_M) as defined in [13, Definition 8.1], then K decomposes with respect to M in the sense of [13, Definition 6.1] (in particular, $K_M = K \cap M$ and $\rho_M = \rho|_{K_M}$) and the equivalence of categories (2.1.31) commutes with parabolic induction and parabolic restriction in the appropriate sense (see [13, Corollary 8.4]).

Proposition 2.1.33. *Let (K_M, ρ_M) be an $\mathfrak{s}_M$ -type for $\mathfrak{s}_M \in \mathfrak{B}(M)$, such that $\Pi_M^{\mathfrak{s}_M} \simeq \text{c-Ind}_{K_M}^M(\rho_M, V_{\rho_M})$. Let (K, ρ) be a G -cover of (K_M, ρ_M) . Then*

$$\Pi_G^{\mathfrak{s}} \simeq \text{c-Ind}_K^G(\rho, V_\rho). \quad (2.1.34)$$

As a consequence, we have

$$\mathcal{H}^s(G) := \text{End}_G(\Pi_G^{\mathfrak{s}}) \simeq \mathcal{H}(G, \rho). \quad (2.1.35)$$

Proof. See [17, Lemma B.3]. $\square$

2.1.36. In this section, in order to be able to apply the constructions of [56] and [34], we assume that $\mathbf{G}$ splits over a tamely ramified extension of F , and that p does not divide the order of the Weyl group of G . By a *Levi subgroup of $\mathbf{G}$* , we mean an F -subgroup of $\mathbf{G}$ which is a Levi factor of a parabolic F -subgroup of $\mathbf{G}$. Let L/F be a finite extension. By a *twisted L -Levi subgroup of $\mathbf{G}$* , we mean an F -subgroup $\mathbf{G}'$ of $\mathbf{G}$ such that $\mathbf{G}' \otimes_F L$ is a Levi subgroup of $\mathbf{G} \otimes_F L$. If L/F is tamely ramified, then $\mathbf{G}'$ is called a *tamely ramified twisted Levi subgroup of $\mathbf{G}$* . A *tamely ramified twisted Levi sequence in $\mathbf{G}$* is a finite sequence $\vec{\mathbf{G}} = (\mathbf{G}^0, \mathbf{G}^1, \dots, \mathbf{G}^d)$ of twisted E -Levi subgroups of $\mathbf{G}$, with E/F tamely ramified (see [56, p. 586]).

Let $\mathcal{B}(\mathbf{G}, F)$ denote the (enlarged) building of G :

$$\mathcal{B}(\mathbf{G}, F) = \mathcal{B}(\mathbf{G}/Z_{\mathbf{G}}, F) \times X_*(Z_G) \otimes_{\mathbb{Z}} \mathbb{R}, \quad (2.1.37)$$

where $X_*(Z_G)$ is the set of F -algebraic cocharacters of Z_G . Recall that when $\mathbf{G}'$ is a tamely ramified twisted Levi subgroup of $\mathbf{G}$, there is a family of natural embeddings of $\mathcal{B}(\mathbf{G}', F)$ into $\mathcal{B}(\mathbf{G}, F)$.

For x a point in $\mathcal{B}(\mathbf{G}, F)$, let $G_{x,0}$ denote the associated parahoric subgroup, and let $G_{x,0+}$ denote the pro- p unipotent radical of $G_{x,0}$. In general, for r a positive real number, $G_{x,r}$ is the corresponding Moy-Prasad filtration subgroup of $G_{x,0}$.

Definition 2.1.38. [34, § 7.1] A *depth-zero G -datum* is a triple

$$((\mathbf{G}, \mathbf{M}), (y, \iota), (K_M, \rho_M)) \quad (2.1.39)$$

satisfying the following

- $\mathbf{G}$ is a connected reductive group over F , and $\mathbf{M}$ is a Levi subgroup of $\mathbf{G}$;
- y is a point in $\mathcal{B}(M)$ such that $M_{y,0}$ is a maximal parahoric subgroup of M , and $\iota: \mathcal{B}(M) \hookrightarrow \mathcal{B}(G)$ is a 0-generic embedding relative to y (see [34, Definition 3.2]);
- K_M is a compact open subgroup of M containing $M_{y,0}$ as a normal subgroup, and ρ_M is an irreducible smooth representation of K_M such that $\rho_M|_{M_{y,0}}$ contains the inflation to $M_{y,0}$ of a cuspidal representation of $M_{y,0}/M_{y,0+}$.

Let $\vec{\mathbf{G}} = (\mathbf{G}^0, \mathbf{G}^1, \dots, \mathbf{G}^d)$ be a tamely ramified twisted Levi sequence in $\mathbf{G}$. To $\vec{\mathbf{G}}$, we associate a sequence of Levi subgroups $\vec{\mathbf{M}} = (\mathbf{M}^0, \dots, \mathbf{M}^d)$, where $\mathbf{M}^i$ is a Levi subgroup of $\mathbf{G}^i$ given as the centralizer of $A_{\mathbf{M}^0}$ in $\mathbf{G}^i$, with $A_{\mathbf{M}^0}$ the maximal F -split torus of the center $Z_{\mathbf{M}^0}$ of M^0 .

Definition 2.1.40. A $\mathbf{G}$ -datum is a 5-tuple

$$\mathcal{D} = ((\vec{\mathbf{G}}, \mathbf{M}^0), (y, \{\iota\}), \vec{r}, (K_{M^0}, \rho_{M^0}), \vec{\phi}) \quad (2.1.41)$$

satisfying the following:

- D1.** $\vec{\mathbf{G}} = (\mathbf{G}^0, \mathbf{G}^1, \dots, \mathbf{G}^d)$ is a tamely ramified twisted Levi sequence in $\mathbf{G}$, and $\mathbf{M}^0$ a Levi subgroup of $\mathbf{G}^0$. Let $\vec{\mathbf{M}}$ be associated to $\vec{\mathbf{G}}$ as above;
- D2.** y is a point in $\mathcal{B}(M^0)$, and $\{\iota\}$ is a commutative diagram of $\vec{s}$ -generic embeddings of buildings relative to y in the sense of [34, Definition 3.5], where $\vec{s} = (0, r_0/2, \dots, r_{d-1}/2)$;
- D3.** $\vec{r} = (r_0, r_1, \dots, r_d)$ is a sequence of real numbers satisfying $0 < r_0 < r_1 < \dots < r_{d-1} \leq r_d$ if $d > 0$, and $0 \leq r_0$ if $d = 0$;
- D4.** (K_{M^0}, ρ_{M^0}) is such that $\mathcal{D}^0 := ((\mathbf{G}^0, \mathbf{M}^0), (y, \iota), (K_{M^0}, \rho_{M^0}))$ is a depth zero G^0 -datum;
- D5.** $\vec{\phi} = (\phi_0, \phi_1, \dots, \phi_d)$ is a sequence of quasi-characters, where ϕ_i is a quasi-character of G^i such that ϕ_i is G^{i+1} -generic of depth r_i relative to x for all $x \in \mathcal{B}(G^i)$ in the sense of [56, § 9].

The construction. For a given $\mathbf{G}$ -datum $\mathcal{D}$ as in (2.1.41), we write

$$K_{\mathcal{D}^0} := K_{M^0} G_{\iota(y),0}^0. \quad (2.1.42)$$

We recall that $G_{\iota(y),0}^0$ is the parahoric subgroup of G^0 associate to the building point $\iota(y)$. Let $G_{\iota(y),0+}^0$ be the pro- p unipotent radical of $G_{\iota(y),0}^0$.

By [34, Proposition 4.3(b)], we have

$$K_{\mathcal{D}^0}/G_{\iota(y),0+}^0 \simeq K_{M^0}/M_{y,0+}^0, \quad (2.1.43)$$

and we define $\rho_{\mathcal{D}^0}$ to be the representation of $K_{\mathcal{D}^0}$ obtained by composing the isomorphism (2.1.43) with ρ_{M^0} .

Definition 2.1.44. A *Kim-Yu type* in the sense of [34, §7.4], which builds on earlier construction in [56], is a pair $(K_{\mathcal{D}}, \rho_{\mathcal{D}})$ where

- $K_{\mathcal{D}}$ is an open compact subgroup given by

$$K_{\mathcal{D}} := K_{\mathcal{D}^0} G_{\iota(y), s_0}^1 \cdots G_{\iota(y), s_{d-1}}^d \quad (2.1.45)$$

- $\rho_{\mathcal{D}}$ is an irreducible representation of $K_{\mathcal{D}}$.

To $\vec{\mathbf{G}}$, we associate a tamely ramified twisted Levi sequence $\vec{\mathbf{M}} = (\mathbf{M}^0, \dots, \mathbf{M}^d)$ of $\mathbf{M}$, where $\mathbf{M}^i$ is the centralizer of $A_{\mathbf{M}}$ in $\mathbf{G}^i$. Consider

$$\mathcal{D}_M := (\vec{\mathbf{M}}, y, \vec{r}, \rho_{M^0}, \vec{\phi}). \quad (2.1.46)$$

When $K_{M^0} = M_y^0$ the datum $\mathcal{D}_M$ gives a supercuspidal type in M as follows.

Let $K_M^d := K_{\mathcal{D}} \cap M$. Let $\tilde{K}_M^d$ denote the normalizer in M of K_M^d . This group $\tilde{K}_M^d$ is a compact modulo center subgroup of M . Let $\rho_M^d := \rho_{\mathcal{D}}|_{K_M^d}$ and consider

$$\sigma_{\mathcal{D}_M} := \text{ind}_{\tilde{K}_M^d}^M \rho_M^d. \quad (2.1.47)$$

Theorem 2.1.48. [34] Suppose that $K_{M^0} = M_y^0$. Then

- (1) (K_M^d, ρ_M^d) is a supercuspidal type on M (as in Definition 2.1.30), and $\sigma_{\mathcal{D}_M}$ is an irreducible supercuspidal representation of M ;
- (2) $(K_{\mathcal{D}}, \rho_{\mathcal{D}})$ is a G -cover of (K_M^d, ρ_M^d) , and thus is an $\mathfrak{s}$ -type for $\mathfrak{s} = [M, \sigma_{\mathcal{D}_M}]_G$.

Proposition 2.1.49. Let $\mathcal{D}$ and $\dot{\mathcal{D}}$ be two G -data

$$\mathcal{D} = ((\vec{\mathbf{G}}, \mathbf{M}^0), (y, \iota), \vec{r}, (K_{M^0}, \rho_{M^0}), \vec{\phi}) \text{ and } \dot{\mathcal{D}} = ((\vec{\mathbf{G}}, \dot{\mathbf{M}}^0), (\dot{y}, \dot{\iota}), \vec{r}, (K_{\dot{M}^0}, \rho_{\dot{M}^0}), \vec{\phi})$$

such that $K_{M^0} = M_y^0$ and $K_{\dot{M}^0} = \dot{M}_{\dot{y}}^0$. Let $\mathfrak{s} := [M, \sigma_{\mathcal{D}_M}]_G$, and $\dot{\mathfrak{s}} := [M, \sigma_{\dot{\mathcal{D}}_M}]_G$.

Then we have $\mathfrak{s} = \dot{\mathfrak{s}}$ if and only if there exists $g \in G$ such that

$${}^g K_{M^0} = K_{\dot{M}^0} \quad \text{and} \quad {}^g (\rho_{M^0} \otimes \phi) \simeq \rho_{\dot{M}^0} \otimes \dot{\phi}, \quad (2.1.50)$$

where $\phi := \prod_{i=0}^d (\phi_i|_{M^0})$ and $\dot{\phi} := \prod_{i=0}^d (\dot{\phi}_i|_{\dot{M}^0})$.

Proof. It is a reformulation of [34, Theorem 10.3]. Indeed, when $K_{M^0} = M_y^0$ and $K_{\dot{M}^0} = \dot{M}_{\dot{y},0}^0$, we have $\mathfrak{s} = \dot{\mathfrak{s}}$ if and only if the types $(K_{\mathcal{D}}, \rho_{\mathcal{D}})$ and $(K_{\dot{\mathcal{D}}}, \rho_{\dot{\mathcal{D}}})$ are equivalent in the sense of [34, Definition 10.1]. Note that [34, Theorems 10.2 and 10.3] still hold without assuming the hypothesis $C(\vec{\mathbf{G}})$ of [28, Remark 2.49 & above], since [31, §3.5] shows that [28, Theorems 6.6 and 6.7] are valid without assuming $C(\vec{\mathbf{G}})$. $\square$

Remark 2.1.51. If $G = M$, it follows from [28, Theorems 6.6 and 6.7] that $\mathfrak{s}_M = \dot{\mathfrak{s}}_M$ if and only if the data $\mathcal{D}_M = (\vec{M}, y, \vec{r}, \rho_{M^0}, \vec{\phi})$ and $\dot{\mathcal{D}}_M = (\vec{M}, \dot{y}, \vec{r}, \rho_{\dot{M}^0}, \vec{\phi})$ are equivalent in the sense of [28, Definition 5.3].

2.1.52. For any $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$, consider

$$N_G^{\mathfrak{s}} := \{n \in N_{\mathbf{G}}(\mathbf{M})(F) : {}^n\sigma \simeq \chi \otimes \sigma \text{ for some } \chi \in \mathfrak{X}_{\text{nr}}(M)\}. \quad (2.1.53)$$

Corollary 2.1.54. We suppose that p does not divide the order of the Weyl group of G . Let $\mathfrak{s} \in \mathfrak{B}(G)$ be an arbitrary Bernstein inertial class. For every $n \in N_G^{\mathfrak{s}}$, there exists an $m \in M$ such that

$${}^{mn}K_{M^0} = K_{M^0} \quad \text{and} \quad {}^{mn}(\rho_{M^0} \otimes \phi) \simeq \rho_{M^0} \otimes \phi,$$

where $\phi := \prod_{i=0}^d (\phi_i|_{M^0})$.

Proof. By [22], we have $\sigma = \sigma_{\mathcal{D}_M}$, for some M -datum $\mathcal{D}_M = (\vec{M}, y, \vec{r}, \rho_{M^0}, \vec{\phi})$. Let $n \in N_G^{\mathfrak{s}}$. Thus ${}^n\sigma \simeq \chi \otimes \sigma$ for some $\chi \in \mathfrak{X}_{\text{nr}}(M)$. Let $\dot{\mathcal{D}}_M := (\vec{M}, y, \vec{r}, \rho_{M^0} \otimes \chi|_{K_{M^0}}, \vec{\phi})$. By (2.1.47), we have

$$\chi \otimes \sigma = (\text{ind}_{\widetilde{K}_M}^M \rho_M^d) \otimes \chi \simeq \text{ind}_{\widetilde{K}_M}^M (\rho_M^d \otimes \chi|_{\widetilde{K}_M}). \quad (2.1.55)$$

Since χ is unramified, we have $\text{ind}_{\widetilde{K}_M}^M (\rho_M^d \otimes \chi|_{\widetilde{K}_M}) = \sigma_{\dot{\mathcal{D}}_M}$. Therefore $\chi \otimes \sigma \simeq \sigma_{\dot{\mathcal{D}}_M}$. Applying Remark 2.1.51 to the M -data ${}^n\mathcal{D}_M$ and $\dot{\mathcal{D}}_M$, one can see that these data are equivalent. Hence there exists an $m \in M$ such that

$${}^{mn}K_{M^0} = K_{M^0} \quad \text{and} \quad {}^{mn}(\rho_{M^0} \otimes \phi) \simeq \rho_{M^0} \otimes \chi|_{K_{M^0}} \otimes \phi = \rho_{M^0} \otimes \phi,$$

where the last equality holds because χ is trivial on K_{M^0} . Here $\phi := \prod_{i=0}^d (\phi_i|_{M^0})$. $\square$

2.1.56. As shown in [28], it follows from the original construction in [56] that ρ_M^d is of the form $\rho_M^d = \rho_{M^0} \otimes \kappa$, where the representation $\kappa = \kappa_G$ depends only on $\vec{\phi}$. Suppose $K_{M^0} = M_y^0$. Let $\widetilde{K}_{M^0}$ denote the normalizer of K_{M^0} in M^0 . Consider

$$\sigma^0 := \text{ind}_{\widetilde{K}_{M^0}}^{M^0} \rho_{M^0}. \quad (2.1.57)$$

The representation σ^0 is a depth-zero irreducible supercuspidal representation of M^0 . Let $\mathfrak{s}^0 := [M^0, \sigma^0]_{G^0}$. Let $\Pi_{G^0}^{\mathfrak{s}^0}$ be defined as in (2.1.4), i.e. it is a progenerator for the category $\mathfrak{R}^{\mathfrak{s}^0}(G^0)$. Set $\mathcal{H}^{\mathfrak{s}^0}(G^0) := \text{End}(\Pi_{G^0}^{\mathfrak{s}^0})$.

Proposition 2.1.58.

(1) The algebras $\mathcal{H}^{\mathfrak{s}^0}(G^0)$ and $\mathcal{H}(G^0, \rho_{\mathcal{D}^0})$ are isomorphic.

(2) The algebras $\mathcal{H}^s(G)$ and $\mathcal{H}(G, \rho_{\mathcal{D}})$ are isomorphic.

Proof. We verify the assumptions in Proposition 2.1.33. Firstly, $(K_{\mathcal{D}^0}, \rho_{\mathcal{D}^0})$ is a G^0 -cover of (K_{M^0}, ρ_{M^0}) (see [34, §7.1]) and $(K_{\mathcal{D}}, \rho_{\mathcal{D}})$ is a G -cover of (K_M^d, ρ_M^d) (see [34, Theorem 7.5]).

Secondly, since σ^0 and σ are supercuspidal irreducible representations, an element m^0 of M^0 intertwines ρ_{M^0} if and only if $m^0 \in \tilde{K}_{M^0}$, and an element m of M intertwines ρ_M if and only if $m \in \tilde{K}_M$. Then the proof of [17, Lemma B.4] applies, and shows that $\Pi_M^{\mathfrak{s}_M} \simeq \text{c-Ind}_{K_M}^M(\rho_M, V_{\rho_M})$. Thus the result follows from Proposition 2.1.33. $\square$

Proposition 2.1.59. *If $W_G^{\mathfrak{s}} = \{1\}$, then there is an algebra isomorphism*

$$\mathcal{H}(G, \rho_{\mathcal{D}}) \simeq \mathcal{H}(M, \rho_M^d),$$

which preserves support of functions, and the algebra $\mathcal{H}(G, \rho_{\mathcal{D}})$ is commutative.

Proof. Since $W_G^{\mathfrak{s}} = \{1\}$, we have $N_G(\mathfrak{s}) \subset M$. Then the first assertion follows from [13, (12.1)]. On the other hand, the algebra $\mathcal{H}(M, \rho_M^d)$ is commutative (see for instance [13, (5.6)]). $\square$

Remark 2.1.60. Applying Proposition 2.1.59 to the group G^0 , we see that if $W_{G^0}^{\mathfrak{s}_0} = \{1\}$, then there is an algebra isomorphism

$$\mathcal{H}(G^0, \rho_{\mathcal{D}^0}) \simeq \mathcal{H}(M^0, \rho_{M^0}^0) \quad (2.1.61)$$

that preserves support of functions; thus the algebra $\mathcal{H}(G^0, \rho_{\mathcal{D}^0})$ is also commutative.

2.2. Bernstein blocks

We assume that G splits over a tamely ramified extension, and that the residual characteristic p of F is odd, good for G and does not divide the order of the fundamental groups of $\mathbf{G}_{\text{der}}$. Let $\mathbf{M}$ be an F -rational Levi subgroup of an F -rational parabolic subgroup of $\mathbf{G}$. Then p satisfies the same assumptions with respect to $\mathbf{M}$, i.e. p is good for M and does not divide the order of the fundamental groups of $\mathbf{M}_{\text{der}}$.

Let $(\mathbf{S}, \theta)$ be a pair consisting of a tamely ramified torus $\mathbf{S}$ in $\mathbf{M}$, and a character $\theta: S \rightarrow \mathbb{C}^\times$. For any positive real number r , consider

$$\Sigma_r^{\mathbf{S}, \theta} := \{ \gamma \in \Sigma(\mathbf{M}, \mathbf{S}) : (\theta \circ N_{E/F})(\gamma^\vee(E_r^\times)) = 1 \}. \quad (2.2.1)$$

We have $\Sigma_s^{\mathbf{S}, \theta} \subset \Sigma_r^{\mathbf{S}, \theta}$ for $s < r$. Set $\Sigma_{r+}^{\mathbf{S}, \theta} := \bigcap_{s>r} \Sigma_s^{\mathbf{S}, \theta}$. Then $r \mapsto \Sigma_r^{\mathbf{S}, \theta}$ defines a $\text{Gal}(F_{\text{sep}}/F)$ -invariant filtration. Let $r_{d-1} > r_{d-2} > \cdots > r_0 > 0$ denote the breaks of the filtration, i.e. the r 's such that $\Sigma_r^{\mathbf{S}, \theta} \neq \Sigma_{r+}^{\mathbf{S}, \theta}$. We set $r_{-1} := 0$, and let r_d be the depth of θ . We have $r_d \geq r_{d-1}$. For each i such that $0 \leq i \leq d$, we denote by $\mathbf{M}^i$ the connected

reductive subgroup of $\mathbf{M}$ with maximal torus $\mathbf{S}$ and root system $\Sigma_{r_{i-1}+}^{\mathbf{S},\theta}$. By definition, the root system of $\mathbf{M}^d$ is $\Sigma(\mathbf{M}, \mathbf{S})$, and thus $\mathbf{M}^d = \mathbf{M}$. The $\mathbf{M}^i$'s are tame twisted Levi subgroups of $\mathbf{M}$ by [31, Lemma 3.6.1]. Moreover, the root system of $\mathbf{M}^0$ is $\Sigma_{0+}^{\mathbf{S},\theta}$; if the latter is empty, we have $\mathbf{M}^0 = \mathbf{S}$.

Denote $M^i := \mathbf{M}^i(F)$. By [31, Proposition 3.6.7], the pair $(\mathbf{S}, \theta)$ has a Howe factorization with respect to a sequence $(\phi_{-1}, \phi_0, \dots, \phi_d)$ of characters, where $\phi_{-1}: S \rightarrow \mathbb{C}^\times$ and $\phi_i: M^i \rightarrow \mathbb{C}^\times$ for $0 \leq i \leq d$. More precisely, we have

$$\theta = \prod_{i=-1}^d \phi_i, \quad (2.2.2)$$

- for any $i \in \{0, \dots, d\}$, the character ϕ_i is trivial on $(\mathbf{M}^i)_{\text{sc}}$, has depth r_i and is $\mathbf{M}^{i+1}$ -generic for any $i \neq d$;
- ϕ_d is trivial if $r_d = r_{d-1}$, and has depth r_d otherwise.

2.2.3. From now on, we make the following assumptions: $\mathbf{S}$ is an elliptic maximal torus of $\mathbf{M}$; the splitting extension of $\mathbf{S}$ is tamely ramified; $\mathbf{S}$ is maximally unramified inside $\mathbf{M}^0$, i.e. S coincides with its maximal unramified subtorus as in [31, Definition 3.4.2]; θ is k_F -regular with respect to M^0 , in the sense of [32, Definition 3.1.1].

For any point x in the building of M , let $[x]$ be the projection of x onto the reduced building $\mathcal{B}_{\text{red}}(M)$. Let M_x (resp. $M_{[x]}$) be the subgroup of M fixing x (resp. $[x]$). Recall that $M_{[x]} = N_M(M_{x,0})$ by [56, Lemma 3.3]. As in [31, Lemma 3.4.3], we can then associate to S a vertex $[y]$ of $\mathcal{B}_{\text{red}}(M)$, which is the unique $\text{Gal}(F^{\text{nr}}/F)$ -fixed point in the apartment $\mathcal{A}_{\text{red}}(S, F^{\text{nr}})$ of $\mathcal{B}_{\text{red}}(M)$.

Let S_b be the unique maximal bounded subgroup of S (which is also the unique maximal compact subgroup of S). Denoting by $\mathfrak{S}^\circ$ the connected Néron model of $\mathbf{S}$, we write $S_0 := \mathfrak{S}^\circ(\mathfrak{o}_f) \subset S_b$ (see [31, §3.1] for more details). Let $\mathbf{M}_{y,0}^0$ be the connected reductive k_F -group such that

$$\mathbb{M}_{y,0}^0 := \mathbf{M}_{y,0}^0(k_F) = M_{y,0}^0/M_{y,0+}^0. \quad (2.2.4)$$

There exists an elliptic maximal k_F -torus $\mathbf{S}$ of $\mathbf{M}_{y,0}^0$ such that for every unramified extension F' of F , the image of $\mathbf{S}(F')_0$ in $\mathbf{M}(F')_{y,0}/\mathbf{M}(F')_{y,0+}$ is equal to $\mathbf{S}(k_{F'})$ (see [31, Lemma 3.4.4]). By [31, Lemma 3.4.14], the character $\phi_{-1}|_{S_0}$ factors through a regular character $\overline{\phi_{-1}}$ of $\mathbf{S} := \mathbf{S}(k_F)$ as defined in [31, Definition 3.4.16]. In particular, $\overline{\phi_{-1}}$ is in general position in the sense of [21, Definition 5.15]. Then it follows from [21, Proposition 7.4, Theorem 8.3] that the Deligne-Lusztig character

$$(-1)^{r(\mathbf{M}_{y,0}^0) - r(\mathbf{S})} R_{\mathbf{S}}^{\mathbf{M}_{y,0}^0}(\overline{\phi_{-1}}) \quad (2.2.5)$$

can be represented by a *cuspidal*, i.e. which cannot be obtained from a proper parabolic induction, $\mathbb{M}_{y,0}^0$ -module $\kappa_{S, \phi_{-1}}$, where $r(?)$ denotes the k_F -rank of $?$. Then $\kappa_{S, \phi_{-1}}$ is

irreducible (see [21, Definition 5.15]), and its pull-back to $M_{y,0}^0$ extends uniquely to a representation $\kappa_{S,\phi_{-1}}$ of $SM_{y,0}^0$. We define

$$\rho_{\mathbf{S},\theta} := \text{Ind}_{SM_{y,0}^0}^{M_{[y]}^0} \kappa_{S,\phi_{-1}} \quad \text{and} \quad \sigma^0 := \text{c-Ind}_{M_{[y]}^0}^{M^0} \rho_{\mathbf{S},\theta}. \quad (2.2.6)$$

Then σ^0 is a depth-zero irreducible regular supercuspidal representation of M^0 (see [31, Definition 3.4.19 & Proposition 3.4.20]). Set $\mathfrak{s}_{M^0} := [M^0, \sigma^0]_{M^0}$.

More generally, we define an irreducible supercuspidal representation σ of M by using the twisted Yu construction of [23]. As observed in [32, §3.4], it has the same effect as using the original Yu construction from [56] applied to the character $\theta \cdot \epsilon$, where $\epsilon: S \rightarrow \{\pm 1\}$ is the product of the characters $\epsilon_y^{M^i/M^{i-1}}$ of [23, Theorem 3.4]. The representation σ is regular, i.e. satisfies [31, Definition 3.7.13]. Then $\chi \otimes \sigma$ is regular for any $\chi \in \mathfrak{X}_{\text{nr}}(M)$, and we say that the inertial class $\mathfrak{s} = [M, \sigma]_G$ is regular.

2.2.7. For $\mathfrak{s}_M = [M, \sigma]_M$ and $\mathfrak{s}_{M^0} = [M^0, \sigma^0]_{M^0}$, the map

$$\begin{aligned} \mathfrak{f}: \text{Irr}^{\mathfrak{s}_M}(M) &\longrightarrow \text{Irr}^{\mathfrak{s}_{M^0}}(M^0), \\ \sigma \otimes \chi &\longmapsto \sigma^0 \otimes \chi|_{M^0}, \quad \chi \in \mathfrak{X}_{\text{nr}}(M), \end{aligned} \quad (2.2.8)$$

is an isomorphism of varieties by [41, Theorem 6.1]. Let $\mathcal{O}^0$ be the orbit of σ^0 under the action of $\mathfrak{X}_{\text{nr}}(M^0)$. Let $W_{G^0}^{\mathfrak{s}^0} = W_{\mathcal{O}^0} \rtimes R(\mathcal{O}^0)$ be the decomposition analogous to (2.1.14). Then (2.2.8) and (2.1.11), applied to both $\mathfrak{s}_M$ and $\mathfrak{s}_{M^0}$, show that the orbits $\mathcal{O}$ and $\mathcal{O}^0$ are isomorphic. The following is a consequence of [5, 7.3, 9.3].

Lemma 2.2.9 (Adler-Mishra). *Suppose that p is good for G and does not divide the order of the fundamental group of $\mathbf{G}_{\text{der}}$. Let $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$ be a regular inertial class. Then (1) there is a group isomorphism*

$$\mathfrak{w}_\sigma: W_G^{\mathfrak{s}} \longrightarrow W_{G^0}^{\mathfrak{s}^0}, \quad (2.2.10)$$

where $\mathfrak{s}^0 = [M^0, \sigma^0]_{G^0}$, and $\mathfrak{f}$ is equivariant with respect to $\mathfrak{w}_\sigma$.

(2) there is an isomorphism

$$\mathfrak{l}_\sigma: (\text{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{h}} \longrightarrow (\text{Irr}^{\mathfrak{s}_{M^0}}(M^0) // W_{G^0}^{\mathfrak{s}^0})_{\mathfrak{h}^0}. \quad (2.2.11)$$

The collection $\mathfrak{h}^0$ of 2-cocycles is defined as follows. For $x \in \text{Irr}^{\mathfrak{s}_M}(M)$, let $W_G^{\mathfrak{s},x}$ denote the stabilizer of x in $W^{\mathfrak{s}}$. Since $\mathfrak{f}$ is equivariant with respect to $\mathfrak{w}_\sigma$, the latter restricts to an isomorphism

$$\mathfrak{w}_\sigma|_{W_G^{\mathfrak{s},x}}: W_G^{\mathfrak{s},x} \longrightarrow W_{G^0}^{\mathfrak{s}^0, \mathfrak{f}(x)}, \quad (2.2.12)$$

and every 2-cocycle $\mathfrak{h}_x: W_G^{\mathfrak{s},x} \times W_G^{\mathfrak{s},x} \longrightarrow \mathbb{C}^\times$ defines a 2-cocycle

$$\mathfrak{h}_{\mathfrak{f}(x)}^0: W_{G^0}^{\mathfrak{s}^0, \mathfrak{f}(x)} \times W_{G^0}^{\mathfrak{s}^0, \mathfrak{f}(x)} \longrightarrow \mathbb{C}^\times. \quad (2.2.13)$$

Consequently, we obtain in Theorem 2.2.14(2) new cases of [5, Conjecture 1.1].

Theorem 2.2.14. *Suppose that p is good for G and does not divide the order of the fundamental group of $\mathbf{G}_{\text{der}}$. Let $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$ be a regular inertial class.*

(1) *Then*

$$(\xi_{G^0}^{\mathfrak{s}^0})^{-1} \circ \mathfrak{l}_\sigma \circ \xi_G^{\mathfrak{s}}: \text{Irr}^{\mathfrak{s}}(G) \longrightarrow \text{Irr}(G^0)^{\mathfrak{s}^0} \quad (2.2.15)$$

is a bijection.

(2) *We have a bijection*

$$\text{Irr}(\mathcal{H}^{\mathfrak{s}}(G)) \longrightarrow \text{Irr}(\mathcal{H}^{\mathfrak{s}^0}(G^0)). \quad (2.2.16)$$

Proof. (1) This follows from the fact that the map ξ_G defined in (2.1.21) and the analogous map

$$\xi_{G^0}^{\mathfrak{s}^0}: \text{Irr}(G)^{\mathfrak{s}^0} \longrightarrow (\text{Irr}^{\mathfrak{s}^0}(G^0) // W_{G^0}^{\mathfrak{s}^0})_{\mathfrak{h}}, \quad (2.2.17)$$

are isomorphisms.

(2) By [54, Theorem 9.7] applied to both G and G^0 , we have

$$\text{Irr}^{\mathfrak{s}}(G) \cong \text{Irr}(\text{End}(i_P^G(V_B))) \quad \text{and} \quad \text{Irr}^{\mathfrak{s}^0}(G^0) \cong \text{Irr}(\text{End}(i_{P^0}^{G^0}(V_{B^0}))).$$

Thus by (1), we have $\text{Irr}(\text{End}(i_P^G(V_B))) \cong \text{Irr}^{\mathfrak{s}}(G) \cong \text{Irr}^{\mathfrak{s}^0}(G^0) \cong \text{Irr}(\text{End}(i_{P^0}^{G^0}(V_{B^0})))$. Then the result follows by applying Corollary 2.1.22 to both $\mathfrak{s}$ and $\mathfrak{s}^0$. $\square$

Remark 2.2.18. We show in Theorem 4.2.23 that the algebras $\mathcal{H}^{\mathfrak{s}}(G)$ and $\mathcal{H}^{\mathfrak{s}^0}(G^0)$ are isomorphic when $G = G_2$ and M is a maximal Levi subgroup.

2.2.19. We end this section with brief recollections on *non-singular* supercuspidal representations in the sense of [32], as we will consider *non-singular Bernstein blocks* in §3.1.50.

Let $\theta: \mathbf{S}(k_F) \rightarrow \overline{\mathbb{Q}}_\ell$ be a non-singular character. Let $N_{\mathbf{G}}(\mathbf{S})(k_F)_\theta$ denote the stabilizer of the pair $(\mathbf{S}, \theta)$. By [32, Proposition 2.3.3], the character θ extends to the group $N_{\mathbf{G}}(\mathbf{S})(k_F)_\theta$. Let $\mathbf{U} \subset \mathbf{G}$ be the unipotent radical of a k_F -rational Borel subgroup of $\mathbf{G}$, containing $\mathbf{S}$, and let $Y_{\mathbf{U}}$ be the corresponding Deligne-Lusztig variety. Let $\kappa_{(\mathbf{S}, \theta)}$ be the isomorphism class of the representation $H_c^{d_{\mathbf{U}}}(Y_{\mathbf{U}}, \overline{\mathbb{Q}}_\ell)_\theta$. For simplicity of expositions, we only describe the depth-zero situation. The representation $\pi_{(\mathbf{S}, \theta)} := \text{c-Ind}_{G_x}^G \inf_{G_x}^{G_x} \kappa_{(\mathbf{S}, \theta)}$ is supercuspidal but not necessarily irreducible. When it is indeed irreducible, we return to the case where θ is regular as in [31]. When $\pi_{(\mathbf{S}, \theta)}$ is reducible (e.g. when θ is non-singular non-regular), it decomposes as [32, 3.3.3]

$$\pi_{(\mathbf{S}, \theta)} = \sum_{\varrho \in \text{Irr}(\mathbf{N}_{\mathbf{G}}(\mathbf{S})(k_F)_{\theta}, \theta)} \dim(\varrho) \pi_{(\mathbf{S}, \theta, \varrho)}^{\epsilon}, \quad (2.2.20)$$

where the constituents $\pi_{(\mathbf{S}, \theta, \rho)}^{\epsilon} := \text{c-Ind}_{G_x}^G \inf_{\mathbb{G}_x}^{G_x} \kappa_{(\mathbf{S}, \theta, \varrho)}^{\epsilon}$, constructed from $\kappa_{(\mathbf{S}, \theta, \varrho)}^{\epsilon}$ as in [32, Definition 2.7.6], are irreducible *non-singular* supercuspidal representations. Here $\text{Irr}(\mathbf{N}_{\mathbf{G}}(\mathbf{S})(k_F)_{\theta}, \theta)$ denotes the set of irreducible representations of $\mathbf{N}_{\mathbf{G}}(\mathbf{S})(k_F)_{\theta}$ whose restriction to $\mathbf{S}(k_F)$ is θ -isotypic ϵ is a fixed coherent splitting of the family of 2-cocycles $\{\eta_{\Psi, \mathbf{U}}\}$ as in [32, §2.4]. The positive-depth supercuspidal representations can be described similarly, by applying Yu's construction [56] to the representation $\pi_{(G^0, \mathbf{S}, \phi_{-1})}$ of G^0 associated to the pair $(\mathbf{S}, \phi_{-1})$ [32, (3.2)].

3. Local Langlands correspondence for Bernstein blocks

3.1. Axiomatic construction of the correspondence

Let $G^{\vee}$ denote the Langlands dual group of G , i.e. the complex reductive group with root datum dual to that of G . Let $Z_{G^{\vee}}$ be the center of $G^{\vee}$ and $G_{\text{ad}}^{\vee}$ the quotient $G^{\vee}/Z_{G^{\vee}}$. The L -group of G is defined to be ${}^L G := G^{\vee} \rtimes W_F$. Similarly, $M^{\vee}$ denotes the Langlands dual group of M . Let $Z_{M^{\vee} \rtimes I_F}$ be the center of $M^{\vee} \rtimes I_F$, and define

$$\mathfrak{X}_{\text{nr}}({}^L M) := (Z_{M^{\vee} \rtimes I_F})_{W_F}^{\circ}. \quad (3.1.1)$$

The group $\mathfrak{X}_{\text{nr}}({}^L M)$ is naturally isomorphic to the group $\mathfrak{X}_{\text{nr}}(M)$. We denote the isomorphism $\mathfrak{X}_{\text{nr}}(M) \xrightarrow{\sim} \mathfrak{X}_{\text{nr}}({}^L M)$ by $\chi \mapsto \chi^{\vee}$.

3.1.2. An L -parameter is a continuous morphism $\varphi: W'_F \rightarrow {}^L G$ such that

- $\varphi(w)$ is semisimple for each $w \in W_F$;
- the restriction $\varphi|_{\text{SL}_2(\mathbb{C})}$ is a morphism of complex algebraic groups.

An L -parameter φ is said to be *discrete* if $\varphi(W'_F)$ is not contained in any proper Levi subgroup of $G^{\vee}$. The group $G^{\vee}$ acts on the set of L -parameters. We denote by $\Phi(G)$ the set of $G^{\vee}$ -classes of G -relevant L -parameters. Attached to each L -parameter φ for G , we define several (possibly disconnected) complex reductive groups as follows.

Set $Z_{G^{\vee}}(\varphi) := Z_{G^{\vee}}(\varphi(W'_F))$. Let $Z_{G_{\text{sc}}^{\vee}}^1(\varphi)$ be the inverse image of $Z_{G^{\vee}}(\varphi)/Z_{G^{\vee}}(\varphi) \cap Z_{G^{\vee}}$ (viewed as a subgroup of $G_{\text{ad}}^{\vee}$) under the quotient map $G_{\text{sc}}^{\vee} \rightarrow G_{\text{ad}}^{\vee}$. Then we set

$$\mathcal{G}_{\varphi} := Z_{G^{\vee}}^1(\varphi|_{W_F}). \quad (3.1.3)$$

We also define the following component group

$$\mathcal{S}_{\varphi} := Z_{G^{\vee}}^1(\varphi)/Z_{G^{\vee}}^1(\varphi)^{\circ}. \quad (3.1.4)$$

An *enhancement* of φ is an irreducible representation ϱ of S_φ . Pairs (φ, ϱ) are called *enhanced L -parameters* (for G and its inner forms).

3.1.5. Let $G^\vee$ act on the set of enhanced L -parameters via

$$g \cdot (\varphi, \varrho) = (g\varphi g^{-1}, g \cdot \varrho). \quad (3.1.6)$$

We denote by $\Phi_e(G)$ the set $G^\vee$ -conjugacy classes of enhanced L -parameters. We define an action of $\mathfrak{X}_{\text{nr}}({}^L M)$ on $\Phi_e(M)$ as follows. Given $(\varphi, \varrho) \in \Phi_e(M)$ and $\xi \in \mathfrak{X}_{\text{nr}}({}^L M)$, we define $(\xi\varphi, \varrho) \in \Phi_e(M)$ by $\xi\varphi := \varphi$ on $I_F \times \text{SL}_2(\mathbb{C})$ and $(\xi\varphi)(\text{Fr}_F) := \tilde{\xi}\varphi(\text{Fr}_F)$. Here $\tilde{\xi} \in Z_{M^\vee \rtimes I_F}^\circ$ represents z .

For an L -parameter φ of G , we denote by u_φ the image of $(1, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix})$ under φ . By [7, (92)], we have $u_\varphi \in \mathcal{G}_\varphi^\circ$ and

$$\mathcal{S}_\varphi \simeq \mathcal{Z}_{\mathcal{G}_\varphi}(u_\varphi) / \mathcal{Z}_{\mathcal{G}_\varphi}(u_\varphi)^\circ := A_{\mathcal{G}_\varphi^\circ}(u_\varphi). \quad (3.1.7)$$

Let ϱ be an irreducible representation of $A_{\mathcal{G}_\varphi^\circ}(u_\varphi)$. The pair (u_φ, ϱ) is said to be *cuspidal* if there is an $\mathcal{G}^\circ$ -equivariant *cuspidal* (in the sense of Lusztig [35]) local system on the $\mathcal{G}^\circ$ -conjugacy class of u_φ .

Definition 3.1.8. An enhanced L -parameter (φ, ϱ) is said to be *cuspidal* if φ is discrete and (u_φ, ϱ) is a cuspidal pair in $\mathcal{G}_\varphi$.

3.1.9. From now on, we use the subscript c to denote “cuspidal”. Let (φ_c, ϱ_c) be a cuspidal enhanced L -parameter for M , we denote by

$$\mathfrak{s}^\vee := [{}^L M, \varphi_c, \varrho_c]_{G^\vee} \quad (3.1.10)$$

the $G^\vee$ -conjugacy class of $({}^L M, \mathcal{O}^\vee)$, where $\mathcal{O}^\vee$ is the orbit of (φ_c, ϱ_c) under the action of $\mathfrak{X}_{\text{nr}}({}^L M)$. Let $\mathfrak{B}^\vee(G)$ be the set of such $\mathfrak{s}^\vee$. Set $\mathfrak{s}_M^\vee := [{}^L M, \varphi_c, \varrho_c]_{M^\vee}$. Let

$$\mathcal{N}_{G^\vee}^{\mathfrak{s}^\vee} := \{n \in \mathcal{N}_{G^\vee}(M^\vee) : {}^n(\varphi_c, \varrho_c) \simeq (\varphi_c, \varrho_c) \otimes \chi^\vee \text{ for some } \chi^\vee \in \mathfrak{X}_{\text{nr}}({}^L M)\}. \quad (3.1.11)$$

Denote

$$W_{G^\vee}^{\mathfrak{s}^\vee} := W(M^\vee, \mathcal{O}^\vee) := \mathcal{N}_{G^\vee}^{\mathfrak{s}^\vee} / M^\vee. \quad (3.1.12)$$

The group $W_{G^\vee}^{\mathfrak{s}^\vee}$ is a finite extended Weyl group, i.e. it decomposes as

$$W_{G^\vee}^{\mathfrak{s}^\vee} = W_{\mathcal{O}^\vee} \rtimes R(\mathcal{O}^\vee), \quad (3.1.13)$$

where $W_{\mathcal{O}^\vee}$ is a finite Weyl group, and $R(\mathcal{O}^\vee)$ is a finite abelian group (see (3.2.8)).

3.1.14. Let $\Phi_e^c(M)$ denote the set of $M^\vee$ -conjugacy classes of cuspidal enhanced L -parameters for M . By [7, (115)], there is a decomposition of $\Phi_e(G)$ into series of enhanced L -parameters indexed by the set $\mathfrak{B}^\vee(G)$:

$$\Phi_e(G) = \bigsqcup_{\mathfrak{s}^\vee \in \mathfrak{B}^\vee(G)} \Phi_e^{\mathfrak{s}^\vee}(G), \quad (3.1.15)$$

where $\Phi_e^{\mathfrak{s}^\vee}(G)$ consists of enhanced L -parameters whose *cuspidal support* lies in $\mathfrak{s}^\vee$. Moreover, for any $(\varphi_c, \varrho_c) \in \Phi_e^c(M)$, we have

$$\Phi_e^{\mathfrak{s}^\vee M}(M) = \mathfrak{X}_{\text{nr}}({}^L M) \cdot (\varphi_c, \varrho_c). \quad (3.1.16)$$

For $\mathfrak{s}^\vee = [{}^L M, \varphi_c, \varrho_c]_{G^\vee} \in \mathfrak{B}^\vee(G)$, there exists a bijection [7, Theorem 9.3]

$$\xi_{G^\vee}^{\mathfrak{s}^\vee} : \Phi_e^{\mathfrak{s}^\vee}(G) \longrightarrow (\Phi_e^{\mathfrak{s}^\vee M}(M) // W_{G^\vee}^{\mathfrak{s}^\vee})_{L\mathfrak{h}}, \quad (3.1.17)$$

where $L\mathfrak{h} = ({}^L\mathfrak{h}_z)_{z \in \mathfrak{X}_{\text{nr}}({}^L M) / \mathfrak{X}_{\text{nr}}({}^L M, (\varphi_c, \varrho_c))}$ is a collection of 2-cocycles, and

$$\mathfrak{X}_{\text{nr}}({}^L M, (\varphi_\sigma, \varrho_\sigma)) := \{z \in \mathfrak{X}_{\text{nr}}({}^L M) : z \cdot (\varphi_\sigma, \varrho_\sigma) = (\varphi_\sigma, \varrho_\sigma)\}. \quad (3.1.18)$$

3.1.19. Let $\eta : M \rightarrow \widetilde{M}$ be an F -morphism of connected reductive F -groups with abelian kernel and cokernel. Then by [16, §1.4], η induces a map from the root datum of $\widetilde{M}^\vee$ to that of $M^\vee$. We denote by $\eta^\vee : \widetilde{M}^\vee \rightarrow M^\vee$ the associated morphism of algebraic groups as in [16, §2.1].

$$\begin{aligned} {}^L\eta : {}^L\widetilde{M} &\rightarrow {}^L M \\ (\widetilde{m}, w) &\mapsto (\eta^\vee(\widetilde{m}), w) \text{ for } \widetilde{m} \in M^\vee. \end{aligned} \quad (3.1.20)$$

We recall [16, Desideratum 10.3(5)]: Let $\widetilde{\varphi} : W_F \times W_F \rightarrow \widetilde{M}^\vee$ be an L -parameter for $\widetilde{M}$ and set $\varphi := {}^L\eta \circ \widetilde{\varphi}$. Then for any $\widetilde{\pi} \in \Pi_{\widetilde{\varphi}}$, the representation $\widetilde{\pi} \circ \eta$ is the direct sum of finitely many irreducible representations belonging to Π_φ .

3.1.21. Let $\text{Irr}^{\text{sc}}(M) \subset \text{Irr}(M)$ denote the set of equivalence classes of irreducible supercuspidal representations of M . Let $g \in N_{\mathbf{G}}(\mathbf{M})(F)$. For any $\sigma \in \text{Irr}^{\text{sc}}(M)$, we have ${}^g\sigma \in \text{Irr}^{\text{sc}}(M)$. We denote by c_g the isomorphism

$$c_g : (M, \sigma) \xrightarrow{\sim} (M, {}^g\sigma). \quad (3.1.22)$$

Let ${}^L c_g : {}^L M \rightarrow {}^L M$ be the morphism defined by (3.1.20). Let $w \mapsto w^\vee$ be the canonical isomorphism from $W(M) := N_{\mathbf{G}}(\mathbf{M})(F)/M$ to $W(M^\vee) := N_{G^\vee}(M^\vee)/M^\vee$ defined in [1, Proposition 3.1]. Let n_w (resp. $n_{w^\vee}$) be a representative of w (resp. $w^\vee$) in $N_{\mathbf{G}}(\mathbf{M})(F)$ (resp. $N_{G^\vee}(M^\vee)$). With these notations, we have $c_{n_w}^\vee(m^\vee) = {}^{n_{w^\vee}}m^\vee$.

For $\sigma \in \text{Irr}(M)$, since the equivalence class of ${}^{n_w}\sigma$ does not depend on the choice of the representative n_w , we will simply denote it by ${}^w\sigma$. Similarly, we use the notation $w^\vee$, instead of $n_{w^\vee}$, to denote the action of $N_{G^\vee}(M^\vee)$ on $\Phi_e(M)$.

Property 3.1.23. *Let M be a Levi subgroup of G . Let $\mathfrak{s}_M := [M, \sigma]_M \in \mathfrak{B}(M)$. There exists a map*

$$\begin{aligned} \mathfrak{L}^{\mathfrak{s}_M}: \quad \text{Irr}^{\mathfrak{s}_M}(M) &\longrightarrow \Phi_e^c(M) \\ \sigma &\longmapsto (\varphi_\sigma, \varrho_\sigma) \end{aligned} \quad (3.1.24)$$

such that the following properties are satisfied for any $\sigma \in \text{Irr}^{\mathfrak{s}_M}(M)$:

(1) For any $\chi \in \mathfrak{X}_{\text{nr}}(M)$, we have

$$(\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}) = \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma), \quad (3.1.25)$$

where $\chi \mapsto \chi^\vee$ is the canonical isomorphism $\mathfrak{X}_{\text{nr}}(M) \xrightarrow{\sim} \mathfrak{X}_{\text{nr}}({}^L M)$.

(2) For any $w \in W(M)$, we have

$$w^\vee(\varphi_\sigma, \varrho_\sigma) \simeq (\varphi_{w\sigma}, \varrho_{w\sigma}), \quad (3.1.26)$$

where $w \mapsto w^\vee$ is the canonical isomorphism $W(M) \xrightarrow{\sim} W(M^\vee)$.

Remark 3.1.27.

- (1) Property 3.1.23(1) is closely related to [16, Desideratum 10.3.(2)].
- (2) Property 3.1.23(2) is a stronger version of [16, Desideratum 10.3.(5)] for $\eta = c_g$, and can be viewed as an analogue of [25, Conjecture 5.2.4] for enhanced L -parameters for supercuspidal representations. In the special case where the L -packet of σ is a singleton, Property 3.1.23(2) is in fact equivalent to [16, Desideratum 10.3.(5)] for $\eta = c_g$.

Lemma 3.1.28. *Let $\mathfrak{s} = [M, \sigma]_M \in \mathfrak{B}(G)$ satisfy Property 3.1.23. Then there is a group isomorphism*

$$\mathfrak{r}: W_G^{\mathfrak{s}} \xrightarrow{\sim} W_{G^\vee}^{\mathfrak{s}^\vee}, \quad \text{where } \mathfrak{s}^\vee := [{}^L M, \mathfrak{L}^{\mathfrak{s}_M}(\sigma)]_{G^\vee}. \quad (3.1.29)$$

Moreover, $\mathfrak{L}^{\mathfrak{s}_M}$ is equivariant with respect to $\mathfrak{r}$.

Proof. Let $w \in W_G^{\mathfrak{s}} \subset W(M)$. By the definition of $W_G^{\mathfrak{s}}$, we have ${}^w\sigma \simeq \chi \otimes \sigma$ for some $\chi \in \mathfrak{X}_{\text{nr}}(M)$. By Property 3.1.23, we have

$$w^\vee(\varphi_\sigma, \varrho_\sigma) \simeq (\varphi_{w\sigma}, \varrho_{w\sigma}) \simeq (\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}) \simeq \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma). \quad (3.1.30)$$

Thus $w^\vee \in W_{G^\vee}^{\mathfrak{s}^\vee}$, and the map $w \mapsto w^\vee$ gives a group morphism from $W_G^{\mathfrak{s}}$ to $W_{G^\vee}^{\mathfrak{s}^\vee}$. Reversing the argument, we see that it is an isomorphism. $\square$

We suppose that Property 3.1.23 holds, and that the collections of 2-cocycles $\mathfrak{h}$ and ${}^L\mathfrak{h}$ satisfy the following

$${}^L\mathfrak{h}_{\chi^\vee} = \mathfrak{h}_\chi \quad \text{for any } \sigma \in \mathfrak{s} \text{ and any } \chi \in \mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma), \quad (3.1.31)$$

Theorem 3.1.32.

(1) *We have a canonical isomorphism*

$$\mathfrak{e}: (\text{Irr}^{\mathfrak{s}_M}(M)//W_G^{\mathfrak{s}})_{\mathfrak{h}} \xrightarrow{\sim} (\Phi_e^{\mathfrak{s}_M}(M)//W_{G^\vee}^{\mathfrak{s}^\vee})_{{}^L\mathfrak{h}}. \quad (3.1.33)$$

(2) *There is a bijection*

$$\mathfrak{L} := (\xi_{G^\vee}^{\mathfrak{s}^\vee})^{-1} \circ \mathfrak{e} \circ \xi_G^{\mathfrak{s}}: \text{Irr}^{\mathfrak{s}}(G) \longrightarrow \Phi_e^{\mathfrak{s}^\vee}(G). \quad (3.1.34)$$

Proof. Let $\mathfrak{s} = [M, \sigma]_G$. Consider $(\varphi_\sigma, \varrho_\sigma) := \mathfrak{L}^{\mathfrak{s}_M}(\sigma)$. Then the isomorphism $\mathfrak{X}_{\text{nr}}(M) \simeq \mathfrak{X}_{\text{nr}}({}^L M)$ combined with (3.1.16) shows that

$$\Phi_e^{\mathfrak{s}_M}(M) = \{\chi^\vee \otimes (\varphi_\sigma, \varrho_\sigma) : \chi \in \mathfrak{X}_{\text{nr}}(M)\}. \quad (3.1.35)$$

By Property 3.1.23(1) and (2.1.11), we have

$$\Phi_e^{\mathfrak{s}_M}(M) = \{(\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}) : \chi \in \mathfrak{X}_{\text{nr}}(M)\} \simeq \text{Irr}^{\mathfrak{s}_M}(M). \quad (3.1.36)$$

Recall that $W_G^{\mathfrak{s}, \chi \otimes \sigma}$ denotes the stabilizer of $\chi \otimes \sigma$ in $W_G^{\mathfrak{s}}$. By construction of the extended quotients in (1.3.1), we have

$$(\text{Irr}^{\mathfrak{s}_M}(M)//W_G^{\mathfrak{s}})_{\mathfrak{h}} = \bigsqcup_{\chi \in \mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma)} (\text{Irr}(\mathbb{C}[W_G^{\mathfrak{s}, \chi \otimes \sigma}, \mathfrak{h}_\chi])/W_G^{\mathfrak{s}}). \quad (3.1.37)$$

For $\chi \in \mathfrak{X}_{\text{nr}}(M)$, let $W_{G^\vee}^{\mathfrak{s}^\vee, \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)}$ denote the stabilizer of $\chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)$ in $W_{G^\vee}^{\mathfrak{s}^\vee}$. By Property 3.1.23(1), we have

$$\mathfrak{X}_{\text{nr}}(M^\vee, (\varphi_\sigma, \varrho_\sigma)) \simeq \mathfrak{X}_{\text{nr}}(M, \sigma). \quad (3.1.38)$$

Again by (1.3.1), we have

$$(\Phi_e^{\mathfrak{s}_M}(M)//W_{G^\vee}^{\mathfrak{s}^\vee})_{{}^L\mathfrak{h}} = \bigsqcup_{\chi^\vee \in \mathfrak{X}_{\text{nr}}(M^\vee)/\mathfrak{X}_{\text{nr}}(M^\vee, (\varphi_\sigma, \varrho_\sigma))} \text{Irr}(\mathbb{C}[W_{G^\vee}^{\mathfrak{s}^\vee, \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)}, {}^L\mathfrak{h}_{\chi^\vee}])/W_{G^\vee}^{\mathfrak{s}^\vee}. \quad (3.1.39)$$

For any $w \in W_G^{\mathfrak{s}, \chi \otimes \sigma}$, by Property 3.1.23, we have

$$w^\vee(\chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)) = w^\vee(\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}) = (\varphi_{w(\chi \otimes \sigma)}, \varrho_{w(\chi \otimes \sigma)}) = (\varphi_{\chi \otimes \sigma}, \varrho_{\chi \otimes \sigma}). \quad (3.1.40)$$

Thus the morphism $\mathfrak{r}$ from (3.1.29) restricts to an isomorphism

$$W_G^{\mathfrak{s}, \chi \otimes \sigma} \xrightarrow{\sim} W_{G^\vee}^{\mathfrak{s}^\vee, \chi^\vee \cdot (\varphi_\sigma, \varrho_\sigma)}. \quad (3.1.41)$$

Combined with (3.1.37) and (3.1.39), we obtain an isomorphism

$$\mathfrak{c}: (\mathrm{Irr}^{\mathfrak{s}^M}(M) // W_G^{\mathfrak{s}})_{\mathfrak{h}} \longrightarrow (\Phi_e^{\mathfrak{s}^\vee M}(M) // W_{G^\vee}^{\mathfrak{s}^\vee})_{L_{\mathfrak{h}}}. \quad (3.1.42)$$

Then (2) follows from the combination of (1) with Proposition 2.1.20 and (3.2.15). $\square$

Remark 3.1.43. When G is a split classical group, F has characteristic zero and $\mathfrak{L}^{\mathfrak{s}^M}$ is the LLC defined by Arthur in [8], then Lemma 3.1.28 was proved in [42, Theorem 4.1], and Theorem 3.1.32 follows from [43, §3.2 & 3.3].

Remark 3.1.44. The 2-cocycles in $\mathfrak{h}$ and ${}^L\mathfrak{h}$ are expected to be often trivial: (1) They are trivial if the groups $R(\mathcal{O})$ from (2.1.14) and $R(\mathcal{O}^\vee)$ from (3.1.13) have cardinality at most 2, and hence when M is a Levi subgroup of a maximal parabolic subgroup of G .

(2) The 2-cocycles are also trivial when G is a symplectic group or a split special orthogonal group by [26, Theorem 7.7] on the group side and [43, §4.5] on the Galois side. They are trivial for principal series representations of split groups by [46, Corollary 7.9 and Theorem 8.2] on the group side and [3, Theorem 13.1] on the Galois side.

(3) However, there exist cases when these 2-cocycles are not trivial: e.g. see [2, Example 5.5] for an example where $\mathfrak{h}$ is non-trivial, and [7, Example 9.4] for an example where ${}^L\mathfrak{h}$ is non-trivial.

3.1.45. Let σ be a regular supercuspidal irreducible representation of M . Let $\varphi_\sigma: W_F \rightarrow {}^L M$ be the L -parameter of σ as constructed in [32].

Proposition 3.1.46. Let $\mathfrak{s} = [M, \sigma]_G$ be such that the L -packet for σ is a singleton. Let $\mathfrak{s}^\vee = [M^\vee, (\varphi_\sigma, 1)]_{G^\vee}$. Assume that the collections of 2-cocycles $\mathfrak{h}$ and ${}^L\mathfrak{h}$ are both trivial. We have the following bijection

$$\mathfrak{L}: \mathrm{Irr}^{\mathfrak{s}}(G) \longrightarrow \Phi_e^{\mathfrak{s}^\vee}(G). \quad (3.1.47)$$

Proof. Let $\mathfrak{L}^{\mathfrak{s}^M}$ be the map

$$\mathfrak{L}^{\mathfrak{s}^M}: \sigma \mapsto (\varphi_\sigma, 1). \quad (3.1.48)$$

By Proposition 3.1.51, Property 3.1.23(1) is satisfied.

The validity of [16, Desideratum 10.3(5)] has been established in [15, Theorem 1.1] when the L -parameter $\tilde{\varphi}$ is supercuspidal, and G is quasi-split. Since the L -packet for σ

is a singleton, Property 3.1.23 (2) holds by Remark 3.1.27. The result thus follows from Corollary 3.1.32. $\square$

Remark 3.1.49. Proposition 3.1.46 is sufficient for the case of $G = G_2$ (as also exemplified in [9]), since M is either GL_2 or a torus, both only having singleton L -packets for their supercuspidals.

3.1.50. Suppose now σ is a *non-singular* supercuspidal irreducible representation of M . Let $\varphi_\sigma: W_F \rightarrow {}^L M$ be the L -parameter of σ defined in [32, §4.1], with enhancement ϱ_σ .

More precisely, consider the $\pi_{(\mathbf{S}, \theta, \varrho)}^\epsilon$ recalled in § 2.2.19. We fix a coherent splitting ϵ . We recall the construction of the non-singular L -parameter $\varphi_{(\mathbf{S}, \theta)}$ in [32, §4.1] (we drop ϱ from the notation as the L -parameter does not depend on ϱ), which is given as a composition $W_F \xrightarrow{\varphi_S} {}^L S \xrightarrow{{}^L j} {}^L G$. Here $\varphi_S: W_F \rightarrow {}^L S$ is Langlands parameter for the character θ , and ${}^L j: {}^L S \rightarrow {}^L G$ is a certain L -embedding arising from the χ -data as part of some *torally wild L -packet datum* $(\mathbf{S}, \hat{j}, \chi_0, \theta)$ as [32].

Proposition 3.1.51. *The map $\mathfrak{L}^{\mathbf{S}^M}: \sigma \mapsto (\varphi_\sigma, \varrho_\sigma)$ satisfies Property 3.1.23 (1).*

Proof. By [32, Proposition 3.4.6], we have $\chi \otimes \pi_{(\mathbf{S}, \theta, \varrho)}^\epsilon = \pi_{(\mathbf{S}, \chi \cdot \theta, \chi \otimes \rho)}^\epsilon$. On the other hand, by [30, Proposition 4.5.3], we have $\varphi_{(\mathbf{S}, \chi \cdot \theta)} = (\varphi_\chi \cdot \varphi_S) \circ {}^L j$ (since the L -embedding ${}^L j$ stays the same after twisting by χ), where $\varphi_\chi: W_F \rightarrow Z_{\hat{G}}$ is the corresponding 1-cocycle as defined [30]. $\square$

Remark 3.1.52. Property 3.1.23 (2) of the map $\mathfrak{L}^{\mathbf{S}^M}: \sigma \mapsto (\varphi_\sigma, \varrho_\sigma)$ follows from [33, Conjecture 2.12], which is expected to hold for LLC for non-singular supercuspidal representations. The authors intend to return to this question in future work.

3.2. Matching of simple modules of extended affine Hecke algebras

In this section, we use Corollary 3.1.32 to obtain a bijection between simple modules of extended affine Hecke algebras for the p -adic group and Galois sides, assuming Property 3.1.23. Note that this is a completely reasonable assumption, as many groups satisfy this property. Therefore, our main results give a new approach to constructing local Langlands correspondences “inductively”, building from LLC’s on the Levi subgroups’.

3.2.1. We now recall the construction of a (possibly twisted) extended affine Hecke algebra $\mathcal{H}^{\mathbf{S}^\vee}(G^\vee)$ constructed in [6]. Consider

$$\mathcal{M}_{\varphi_c} := Z_{M^\vee}(\varphi_c(W_F)). \quad (3.2.2)$$

Let $\mathcal{A}_{\varphi_c}$ be the identity component of the center of $\mathcal{M}_{\varphi_c}$. We set

$$\mathcal{J}_\varphi = \mathcal{J}_\varphi^{G^\vee} := \mathbf{Z}_{G^\vee}(\varphi(I_F)). \quad (3.2.3)$$

Let $\Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})$ be the set of $\alpha \in X^*(\mathcal{A}_{\varphi_c}) \setminus \{0\}$ which appear in the adjoint action of $\mathcal{A}_{\varphi_c}$ on the Lie algebra of $\mathcal{J}_\varphi^\circ$. It is a root system by [6, Proposition 3.9]. Let $\Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})^+$ be the positive root system defined by an F -rational Borel subgroup of $\mathcal{J}_\varphi^\circ$. Let Δ be a basis of the reduced root system $\Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})_{\text{red}}$. Let $a \in \mathcal{A}_{\varphi_c}$ be such that $\alpha(a^{-1})$ is an eigenvalue of $\text{Ad}(\varphi(\text{Fr}))$ for any $\alpha \in \Delta$. We define $\varphi_a \in \Phi(M)$ by

$$\varphi_a|_{I_F \times \text{SL}_2(\mathbb{C})} := \varphi_c|_{I_F \times \text{SL}_2(\mathbb{C})} \quad \text{and} \quad \varphi_a(\text{Fr}_F) := a \cdot \varphi_c(\text{Fr}_F). \quad (3.2.4)$$

By [6, Proposition 3.9], we have $\Sigma(\mathcal{G}_{\varphi_a}, \mathcal{A}_{\varphi_c})_{\text{red}} = \Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})_{\text{red}}$, where $\mathcal{G}_{\varphi_a} = \mathbf{Z}_{G^\vee}(\varphi_a(W_F))$. Consider

$$\mathfrak{X}_{\text{nr}}(M^\vee, \varphi_a) := \{x \in \mathfrak{X}_{\text{nr}}(M^\vee) : (z\varphi_a)_{M^\vee} = (\varphi_a)_{M^\vee}\}. \quad (3.2.5)$$

Set $T_{\mathcal{O}^\vee} := \mathfrak{X}_{\text{nr}}(M^\vee)/\mathfrak{X}_{\text{nr}}(M^\vee, \varphi_a)$. For each $\alpha \in \Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})_{\text{red}}$, let $m_\alpha \in \mathbb{Z}_{>0}$ be the smallest integer such that

$$\ker(m_\alpha \alpha) \supset \{a' \in \mathcal{A}_{\varphi_c} : (a'\varphi_a)_{M^\vee} = (\varphi_a)_{M^\vee}\}. \quad (3.2.6)$$

We set

$$\Sigma_{\mathcal{O}^\vee} := \{m_\alpha : \alpha \in \Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})_{\text{red}}\} \subset X^*(T_{\mathcal{O}^\vee}). \quad (3.2.7)$$

Then $W_{\mathcal{O}^\vee}$ from (3.1.13) is the finite Weyl group of $\Sigma_{\mathcal{O}^\vee}$, and

$$R(\mathcal{O}^\vee) := \{w \in W(M^\vee, \mathcal{O}^\vee) : w \cdot \Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})^+ = \Sigma(\mathcal{J}_\varphi^\circ, \mathcal{A}_{\varphi_c})^+\}. \quad (3.2.8)$$

Let

$$\lambda^\vee : \Sigma_{\mathcal{O}^\vee} \rightarrow \mathbb{Z}_{\geq 0} \quad \text{and} \quad \lambda^{*\vee} : \{m_\alpha \alpha \in \Sigma_{\mathcal{O}^\vee} : (m_\alpha \alpha)^\vee \in 2X_*(T_{\mathfrak{s}^\vee})\} \rightarrow \mathbb{Z}_{\geq 0} \quad (3.2.9)$$

be the two parameter functions defined in the proof of [6, Lemma 3.12]. Recall from [6, (28)] that $\lambda^{*\vee}(\alpha) = \lambda^\vee(\alpha)$ unless α is a short root in a type B root subsystem of $R_{\mathcal{O}^\vee}$.

The algebra $\mathcal{H}^{\mathfrak{s}^\vee}(G^\vee)$ is defined to be

$$\mathcal{H}^{\mathfrak{s}^\vee}(G^\vee) := \mathcal{H}_{\text{aff}}(\mathcal{O}^\vee, \Sigma_{\mathcal{O}^\vee}, \lambda^\vee, \lambda^{*\vee}, z) \rtimes \mathbb{C}[R(\mathcal{O}^\vee), \kappa^\vee], \quad (3.2.10)$$

where z is a positive real number, $\mathcal{H}_{\text{aff}}(G^\vee, \mathfrak{s}^\vee) := \mathcal{H}_{\text{aff}}(\mathcal{O}^\vee, \Sigma_{\mathcal{O}^\vee}, \lambda^\vee, \lambda^{*\vee}, z)$ is the corresponding affine Hecke algebra with affine Weyl group $W_{\mathcal{O}^\vee} \ltimes X^*(T_{\mathcal{O}^\vee})$, and $\kappa^\vee$ a 2-cocycle on $R(\mathcal{O}^\vee)$.

Theorem 3.2.11. *We suppose that the collections of 2-cocycles $\natural$ and ${}^L\natural$ are both trivial. Let $\mathfrak{s} = [M, \sigma]_G$ and $\mathfrak{s}^\vee = [{}^LM, \varphi_\sigma, 1]_{G^\vee}$. There is a bijection*

$$\mathrm{Irr}(\mathcal{H}^{\mathfrak{s}}(G)) \longleftrightarrow \mathrm{Irr}(\mathcal{H}^{\mathfrak{s}^\vee}(G^\vee)). \quad (3.2.12)$$

Proof. By Corollary 2.1.22, we have a bijection

$$\mathrm{Irr}(\mathcal{H}^{\mathfrak{s}}(G)) \longrightarrow (\mathrm{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}})_{\natural}. \quad (3.2.13)$$

By Corollary 3.1.32, we have

$$\mathrm{Irr}^{\mathfrak{s}_M}(M) // W_G^{\mathfrak{s}} \simeq \Phi_e^{\mathfrak{s}_M^\vee}(M) // W_{G^\vee}^{\mathfrak{s}^\vee}. \quad (3.2.14)$$

On the other hand, by (3.1.17) we have a bijection

$$\Phi_e^{\mathfrak{s}_M^\vee}(M) // W_{G^\vee}^{\mathfrak{s}^\vee} \longrightarrow \Phi_e^{\mathfrak{s}^\vee}(G). \quad (3.2.15)$$

Finally, by [6, Theorem 3.18], there is a bijection

$$\Phi_e^{\mathfrak{s}^\vee}(G) \xrightarrow{\sim} \mathrm{Irr}(\mathcal{H}^{\mathfrak{s}^\vee}(G^\vee)). \quad (3.2.16)$$

Combining equations (3.2.13), (3.2.14), (3.2.15) and (3.2.16), we obtain the desired bijection. $\square$

Corollary 3.2.17. *With the same assumption as in Theorem 3.2.11.*

$$\mathrm{Irr}(\mathcal{H}^{\mathfrak{s}^\vee}(G^\vee)) \longrightarrow \mathrm{Irr}(\mathcal{H}^{(\mathfrak{s}^0)^\vee}((G^0)^\vee)).$$

Proof. This follows from combining Theorems 3.2.11 and 2.2.14. $\square$

4. Applications to G_2

In this section, we introduce notations and background specifically for the G_2 case.

4.0.1. General background

4.0.1. Let $\mathfrak{a}_M$ be the real Lie algebra of A_M , and $\mathfrak{a}_M^*$ its dual. Let $\mathfrak{a}_{M,\mathbb{C}}^*$ be the complexification of $\mathfrak{a}_M^*$. Let $|\cdot|_F$ be the modulus of F . Let $H_M : M \rightarrow \mathfrak{a}_M^*$ be such that $q^{-\langle H_M(m), \alpha \rangle} = |\alpha(m)|_F$ for every rational character α of M and every $m \in M$. Note that the kernel of H_M is equal to M_1 (recall from § 2.1.1). Consider

$$M_\sigma := \bigcap_{\chi \in \mathfrak{X}_{\mathrm{nr}}(M, \sigma)} \ker \chi, \quad (4.0.2)$$

which has finite index in M_1 . Recall $\mathfrak{X}_{\text{nr}}(M, \sigma)$ from (2.1.9), and we have

$$\text{Irr}(M_\sigma/M_1) \simeq \mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma) \quad \text{and} \quad \mathbb{C}[M_\sigma/M_1] \simeq \mathbb{C}[\mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma)]. \quad (4.0.3)$$

Set $(M_\sigma/M_1)^\vee := \text{Hom}_{\mathbb{Z}}(M_\sigma/M_1, \mathbb{Z})$. Composing H_M with the $\mathbb{R}$ -linear extension of $H_M(M_\sigma/M_1) \rightarrow \mathbb{Z}$ gives an embedding

$$H_M^\vee : (M_\sigma/M_1)^\vee \rightarrow \mathfrak{a}_M^*. \quad (4.0.4)$$

For $m \in M$, let b_m be the element of $\mathbb{C}[\mathfrak{X}_{\text{nr}}(M)]$ defined by $b_m(\chi) := \chi(m)$ for any $\chi \in \mathfrak{X}_{\text{nr}}(M)$. Let h_α be the unique generator of M_σ/M_1 such that $\text{val}_F(\alpha(h_\alpha)) > 0$. We define $X_\alpha \in \mathbb{C}(\mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma))$ by

$$X_\alpha(\chi) := \chi(h_\alpha), \quad (4.0.5)$$

for $\chi \in \mathfrak{X}_{\text{nr}}(M)/\mathfrak{X}_{\text{nr}}(M, \sigma)$.

4.0.6. For a complex number s , let χ_s be the character defined by

$$\chi_s(m) := |\det(m)|_F^s \quad \text{for any } m \in M. \quad (4.0.7)$$

In particular, $\chi_s \in \mathfrak{X}_{\text{nr}}(M)$, and we have

$$X_\alpha(\chi_s) = |\det(h_\alpha)|_F^s. \quad (4.0.8)$$

Let $\tilde{\alpha}$ be the element of $\mathfrak{a}_M^*$ defined by $\tilde{\alpha} := \langle \rho_P, \alpha^\vee \rangle^{-1} \rho_P$, where ρ_P is half the sum of the roots of A_M in $\text{Lie } N$, with $P = MN$. Then $s\tilde{\alpha} \in \mathfrak{a}_M^* \otimes \mathbb{R} \subset \mathbb{C}$.

We recall the description of the Plancherel measure from [52] (see also [53] or [26] for the notations used here): for $\alpha \in \Sigma_{\mathcal{O}, \mu}$, where $\Sigma_{\mathcal{O}, \mu}$ is the root system defined in (2.1.13), there exist $q_\alpha, q_{\alpha^*} \in \mathbb{R}_{\geq 1}, c'_{s_\alpha} \in \mathbb{R}_{>0}$ for $\alpha \in \Sigma_{\mathcal{O}, \mu}$, such that

$$\mu^{M_\alpha}(\sigma \otimes \cdot) = c'_{s_\alpha} \frac{(1 - X_\alpha)(1 - X_\alpha^{-1})}{(1 - q_\alpha^{-1} X_\alpha)(1 - q_\alpha^{-1} X_\alpha^{-1})} \cdot \frac{(1 + X_\alpha)(1 + X_\alpha^{-1})}{(1 + q_{\alpha^*}^{-1} X_\alpha)(1 + q_{\alpha^*}^{-1} X_\alpha^{-1})}. \quad (4.0.9)$$

4.0.10. For $\alpha \in \Sigma_{\mathcal{O}, \mu}$, by [54, Proposition 3.1] there is a unique $\alpha^\sharp \in (M_\sigma/M_1)^\vee$ such that $H_M^\vee(\alpha^\sharp) \in \mathbb{R}\alpha$ and $\langle h_\alpha, \alpha^\sharp \rangle = 2$. We set

$$\Sigma_{\mathcal{O}} := \{\alpha^\sharp : \alpha \in \Sigma_{\mathcal{O}, \mu}\} \quad \text{and} \quad \Sigma_{\mathcal{O}}^\vee := \{\alpha^\sharp : h_\alpha \in \Sigma_{\mathcal{O}, \mu}\}.$$

The quadruple $(\Sigma_{\mathcal{O}}^\vee, M_\sigma/M_1, \Sigma_{\mathcal{O}}, (M_\sigma/M_1)^\vee)$ is a root datum with Weyl group $W_{\mathcal{O}}$. It has a natural action of the group $W(M, \mathcal{O})$, and $R(\mathcal{O})$ is the stabilizer of its basis determined by P (see [54]). We endow this based root datum with the parameter q_F and the labels

$$\lambda(\alpha) := \log(q_\alpha q_{\alpha^*}) / \log(q_F) \quad \text{and} \quad \lambda^*(\alpha) := \log(q_\alpha q_{\alpha^*}^{-1}) / \log(q_F). \quad (4.0.11)$$

To the above data we associate the affine Hecke algebra

$$\mathcal{H}_{\text{aff}}^s(\lambda, \lambda^*, q_F) := \mathcal{H}_{\text{aff}}(\Sigma_{\mathcal{O}}^\vee, M_\sigma/M_1, \Sigma_{\mathcal{O}}, (M_\sigma/M_1)^\vee, \lambda, \lambda^*, q_F). \quad (4.0.12)$$

It is defined as the vector space $\mathbb{C}[W_{\mathcal{O}}] \otimes_{\mathbb{C}} \mathbb{C}[M_\sigma/M_1]$ with the following multiplication rules:

- $\mathbb{C}[W_{\mathcal{O}}] = \text{span}\{T_w : w \in W_{\mathcal{O}}\}$ is embedded as $\mathcal{H}(W_{\mathcal{O}}, q_F^\lambda)$, the *Iwahori-Hecke algebra* of $W_{\mathcal{O}}$, i.e.

$$\begin{aligned} T_w T_v &= T_{wv} \quad \text{if } \ell(w) + \ell(v) = \ell(wv), \\ (T_{s_\alpha} + 1)(T_{s_\alpha} - q_F^{\lambda(\alpha)}) &= 0 \quad \text{if } \alpha \in \Delta_{\mathcal{O}, \mu}, \end{aligned} \quad (4.0.13)$$

where $\ell(w)$ is the word length of w ¹;

- $\mathbb{C}[M_\sigma/M_1] \simeq \mathbb{C}[\mathcal{O}]$ is embedded as a subalgebra;
- for $\alpha \in \Delta_{\mathcal{O}, \mu}$ and $x \in M_\sigma/M_1$ (corresponding to $\theta_x \in \mathbb{C}[M_\sigma/M_1]$):

$$\theta_x T_{s_\alpha} - T_{s_\alpha} \theta_{s_\alpha(x)} = \left(q_F^{\lambda(\alpha)} - 1 + X_\alpha^{-1} (q^{\frac{\lambda(\alpha) + \lambda^*(\alpha)}{2}} - q^{\frac{\lambda(\alpha) - \lambda^*(\alpha)}{2}}) \right) \frac{\theta_x - \theta_{s_\alpha(x)}}{1 - X_\alpha^{-2}}.$$

4.0.14. We set

$$W_{\text{aff}}^s := W_{\mathcal{O}} \ltimes \mathbb{Z} \Sigma_{\mathcal{O}}^\vee. \quad (4.0.15)$$

From now on we assume that the parabolic subgroup P is maximal. Then we have $M_\alpha = G$, and $W(M)$ is either trivial or of order 2.

Remark 4.0.16.

- (1) The groups $W(M, \mathcal{O})$, $W_{\mathcal{O}}$, and $R(\mathcal{O})$ are either trivial or of order 2. In particular, $\Sigma_{\mathcal{O}, \mu}$ is either empty or $\{\alpha, -\alpha\}$.
- (2) For $G = G_2$, if $\sigma \not\simeq \sigma^\vee$, then $W(M, \mathcal{O}) = 1$. It suffices to only check the case where $\sigma \simeq \sigma^\vee$.

In general, if $W(M, \mathcal{O}) = 1$, then the parabolically induced representation is irreducible, so we do not need to work with the case. In the case of G_2 , the condition $W(M, \mathcal{O}) \neq 1$ happens to be characterized by the condition that σ is self-dual. See [50] for more details.

¹ i.e., $\ell(w)$ is the smallest integer $\ell \geq 0$ such that w is a product of ℓ generators s_α .

4.0.17. If $\Sigma_{\mathcal{O},\mu} \neq \emptyset$, then $W(M) \neq \{1\}$ and the group $W_{\mathcal{O}}$ is generated by the unique non-trivial element of $W(M)$, say s_M . Then we have $W_{\mathcal{O}} = W(M, \mathcal{O}) = W(M)$. In particular, if $\Sigma_{\mathcal{O},\mu} \neq \emptyset$, we have $R(\mathcal{O}) = \{1\}$.

The condition $\Sigma_{\mathcal{O},\mu} = \emptyset$ is equivalent to the following

$$\mu^G(\chi \otimes \sigma) \neq 0 \quad \text{for any } \chi \in X_{\text{nr}}(M). \quad (4.0.18)$$

We recall the following theorem of Harish-Chandra.

Theorem 4.0.19. (Harish-Chandra) [52, 5.4.2.2 and 5.4.2.3] *Let M be a Levi subgroup of a maximal parabolic subgroup of G and let σ be a unitary irreducible supercuspidal representation of M .*

- (a) *If $\mu^G(\sigma) = 0$, then $W(M) = \{1, s_M\} \neq \{1\}$, and $s_M\sigma \simeq \sigma$.*
- (b) *Suppose $W(M) \neq \{1\}$. Then $\mu^G(\sigma) \neq 0$ if and only if the representation $i_P^G(\sigma)$ is reducible. In this case, the representation $i_P^G(\sigma)$ is the direct sum of two non-isomorphic irreducible representations.*

4.0.2. Some background on G_2

In the case where G is the split G_2 , we obtain more precise results than in previous sections. Let T be a maximal split torus in G . Let R be the set of roots of G with respect to T . Let $(\epsilon_1, \epsilon_2, \epsilon_3)$ be the canonical basis of $\mathbb{R}^3$, equipped with the scalar product $(\mid)$ for which this basis is orthonormal. Then $\{\alpha := \epsilon_1 - \epsilon_2, \beta := -2\epsilon_1 + \epsilon_2 + \epsilon_3\}$ defines a basis of R , and

$$R^+ = \{\alpha, \beta, \alpha + \beta, 2\alpha + \beta, 3\alpha + \beta, 3\alpha + 2\beta\} \quad (4.0.20)$$

is a subset of positive roots in R . We have

$$(\alpha|\alpha) = 2, \quad (\beta|\beta) = 6 \quad \text{and} \quad (\alpha|\beta) = -3. \quad (4.0.21)$$

Hence α is a short root, while β is a long root.

4.0.22. As in [44], we fix an isomorphism:

$$\eta_{\alpha}: T \xrightarrow{\sim} F^{\times} \times F^{\times} \quad (4.0.23)$$

$$t \longmapsto ((2\alpha + \beta)(t), (\alpha + \beta)(t)). \quad (4.0.24)$$

Under this identification we have

$$\alpha^{\vee}(a) = \eta_{\alpha}^{-1}(a, a^{-1}) \quad \text{and} \quad \beta^{\vee}(a) = \eta_{\alpha}^{-1}(1, a) \quad \text{for any } a \in F^{\times}. \quad (4.0.25)$$

Let $G^{\vee}$ be the dual group of G over $\mathbb{C}$, obtained via an identification of the roots of $G^{\vee}$ with the coroots of G_2 and vice versa. Then $G^{\vee}$ is a complex reductive group of type

G_2 , with simple roots $\alpha^\vee$ and $\beta^\vee$. Note that $\alpha^\vee$ (resp. $\beta^\vee$) is the long (resp. short) root of $G^\vee$. Consider the torus $T^\vee$ dual to T . Then $T^\vee$ is a maximal torus of $G^\vee$. We fix an isomorphism:

$$\eta_{\beta^\vee}: T \xrightarrow{\sim} \mathbb{C}^\times \times \mathbb{C}^\times \quad (4.0.26)$$

$$t \mapsto ((\alpha^\vee + 2\beta^\vee)(t), (\alpha^\vee + \beta^\vee)(t)). \quad (4.0.27)$$

We have

$$\alpha^\vee(a) = \eta_{\beta^\vee}^{-1}(1, a) \quad \text{and} \quad \beta^\vee(a) = \eta_{\beta^\vee}^{-1}(a, a^{-1}) \quad \text{for any } a \in F^\times. \quad (4.0.28)$$

4.0.29. For each root $\gamma \in R(G)$, we fix root group homomorphisms $x_\gamma: F \rightarrow G$ and $\mathbb{Z}$ -homomorphisms $\zeta_\gamma: \mathrm{SL}_2(F) \rightarrow G$ as in [18, (6.1.3) (b)]. We have

$$x_\gamma(u) = \zeta_\gamma \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}, \quad x_{-\gamma}(u) = \zeta_{-\gamma} \begin{pmatrix} 1 & 0 \\ u & 1 \end{pmatrix} \quad \text{and} \quad \gamma^\vee(t) = \zeta_\gamma \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}. \quad (4.0.30)$$

For $\gamma \in \{\alpha, \beta\}$, let P_γ be the maximal standard parabolic subgroup of G generated by γ . Let M_γ be the centralizer of the image of $(\gamma')^\vee$ in G , where γ' is the unique positive root orthogonal to γ , i.e.

$$\gamma' := \begin{cases} 3\alpha + \beta & \text{if } \gamma = \alpha, \\ 3\alpha + 2\beta & \text{if } \gamma = \beta. \end{cases} \quad (4.0.31)$$

Then M_γ is a Levi factor for P_γ . Moreover, M_α and M_β are representatives of the two conjugacy classes of maximal Levi subgroups of G .

We extend $\zeta_\gamma: \mathrm{SL}_2(F) \rightarrow M_\gamma$ to an isomorphism $\zeta_\gamma: \mathrm{GL}_2(F) \rightarrow M_\gamma$ by

$$\zeta_\gamma \left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) := \zeta_{\gamma'} \left(\begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right), \quad \text{for } t \in F^\times. \quad (4.0.32)$$

Then the restriction of ζ_γ^{-1} to T coincides with the isomorphism

$$\eta_\gamma: T \rightarrow F^\times \times F^\times, \quad (4.0.33)$$

where η_α has been defined in (4.0.23), and

$$\eta_\beta: t \mapsto ((\alpha + \beta)(t), \alpha(t)). \quad (4.0.34)$$

4.1. Explicit Hecke algebra parameters

4.1.1. The long root case

Let ψ be a fixed nontrivial additive character of F , and $\overline{\psi}$ be the dual of ψ . Assume for now the Levi factor M of $P = MN$ is generated by the long root of G . Let σ be an

irreducible unitary supercuspidal representation of M . We denote by $\omega := \omega_\sigma$ the central character of σ . Let L/F be a quadratic extension. Let χ be a character of $L^\times$. Let χ' be the conjugate of χ , i.e. $\chi'(a) = \chi(\bar{a})$. Let $\Pi(\sigma)$ denote the Gelbart-Jacquet lift of σ as defined in [24]. Our notations follow [50]. The Plancherel measure $\mu(s\tilde{\alpha}, \sigma)$ has the following four possibilities ([4]).

4.1.1. Case I. If ω is unramified, and if $\sigma = \sigma(\tau)$ with $\tau = \text{Ind}_{W_L}^{W_F} \chi$, with $\chi^2 \chi'$ unramified, then

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q_F^{n(\omega) + n(\sigma \times \Pi(\sigma)) - n(\sigma)} \frac{(1 - \omega(\varpi) q_F^{-2s})(1 - \omega^{-1}(\varpi) q_F^{2s})}{(1 - \omega^{-1}(\varpi) q_F^{-1+2s})(1 - \omega(\varpi) q_F^{-1-2s})} \quad (4.1.2)$$

$$\cdot \frac{(1 - \chi^2 \chi'^{-1}(\varpi_L) q_L^{-s})(1 - \chi^{-2} \chi'(\varpi_L) q_L^s)}{(1 - \chi^2 \chi'^{-1}(\varpi_L) q_L^{-1-s})(1 - \chi^{-2} \chi'(\varpi_L) q_L^{-1+s})} \quad (4.1.3)$$

Comparing (4.1.2) with the Plancherel formula in (4.0.9), we have

$$\begin{cases} X_\alpha(s) &= \omega(\varpi_F) q_F^{-2s} \\ X_\alpha(s) &= -\chi^2 \chi'^{-1}(\varpi_L) q_L^{-s}, \end{cases} \quad (4.1.4)$$

which implies that

$$\omega(\varpi_F) q_F^{-2s} + \chi^2 \chi'^{-1}(\varpi_L) q_L^{-s} = 0. \quad (4.1.5)$$

Since $q_L = q^{f(L/F)}$, (4.1.5) only has a solution when $f(L/F) = 2$ and

$$\omega(\varpi_F) + \chi^2 \chi'^{-1}(\varpi_L) = 0, \quad (4.1.6)$$

which is satisfied in our case. In particular, we have

$$q_\alpha = q_F, \quad q_{\alpha^*} = q_L = q_F^{f(L/F)}. \quad (4.1.7)$$

Therefore we have

$$\begin{aligned} \lambda(\alpha) &= \log(q_\alpha q_{\alpha^*}) / \log(q_F) = 1 + f(L/F), \\ \lambda^*(\alpha) &= |\log(q_\alpha q_{\alpha^*}^{-1}) / \log(q_F)| = |1 - f(L/F)|. \end{aligned} \quad (4.1.8)$$

Hence the parameters for the extended affine Hecke algebra in this case are $q_F^{1+f(L/F)}$ and $q_F^{|1-f(L/F)|}$.

4.1.9. Case II. If ω_σ is ramified and $\sigma = \sigma(\tau)$ with $\tau = \text{Ind}_{W_L}^{W_F} \chi$, and $\chi^2 \chi'$ unramified,

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q_F^{n(\sigma \times \Pi(\sigma)) - n(\sigma)} \frac{(1 - \chi^2 \chi'^{-1}(\varpi_L) q_L^{-s})(1 - \chi^{-2} \chi'(\varpi_L) q_L^s)}{(1 - \chi^2 \chi'^{-1}(\varpi_L) q_L^{-1-s})(1 - \chi^{-2} \chi'(\varpi_L) q_L^{-1+s})} \quad (4.1.10)$$

We compare (4.1.10) to the Plancherel formula (4.0.9) and obtain

$$q_{\alpha^*} = 1, \quad q_{\alpha} = q_L = q_F^{f(L/F)} \quad (4.1.11)$$

Recall the definition of X_{α} as

$$X_{\alpha}(\chi) := \chi(h_{\alpha}^{\vee}) \quad (4.1.12)$$

where $\chi \in \mathfrak{X}^{\text{nr}}(M)/\mathfrak{X}^{\text{nr}}(M, \sigma)$. Since the map $\psi_s: m \mapsto |\det(m)|_F^s$ is an unramified character of M , we have

$$X_{\alpha}(\psi_s) = (\chi^2 \chi'^{-1})(\varpi_L) q_L^{-s} \quad (4.1.13)$$

Recall from (4.0.11) that

$$q_F^{\lambda(\alpha)} = q_{\alpha} q_{\alpha^*} \in \mathbb{R}_{>1}. \quad (4.1.14)$$

Thus by (4.1.11), we have $q_F^{\lambda(\alpha)} = q_L = q_F^{f(L/F)}$, where $f(L/F)$ is the residue degree and is thus 1 if L/F is ramified, and 2 if L/F is unramified. In particular

$$\lambda(\alpha) = f(L/F), \quad \lambda^*(\alpha) = f(L/F). \quad (4.1.15)$$

Note that for $w \in W(M, \mathcal{O})$, one may check that

$$w(X_{\alpha}) = X_{w(\alpha)} \quad (4.1.16)$$

Since $w(\alpha) = \alpha$ for $w \in W(M, \mathcal{O})$ when $G = G_2$, (4.1.16) is simply $w(X_{\alpha}) = X_{\alpha}$. On the other hand, by [53, Prop 1.1] we have

$$wX_{\alpha}(\chi) = w(X_{\alpha}(\chi)) = w(\chi(h_{\alpha}^{\vee})) = \chi(w(h_{\alpha}^{\vee})) = \chi(h_{w(\alpha)}^{\vee}) = \chi(h_{\alpha}^{\vee}) = X_{\alpha}(\chi)$$

Thus $wX_{\alpha} = X_{\alpha} = X_{w(\alpha)}$. This reduces to check, in the long root case, that

$$s_{2\alpha+\beta}(\chi^2 \chi'^{-1}(\varpi_L) q_L^{-s}) = \chi^2 \chi'^{-1}(\varpi_L) q_L^{-s} \quad (4.1.17)$$

Since $\Sigma_{\mathcal{O}}^{\vee} = \{1, 2\alpha + \beta\}$ in the long root $M = M^{\beta}$ case, we have

$$W(\Sigma_{\mathcal{O}}^{\vee}) = \{1, s_{2\alpha+\beta}\}. \quad (4.1.18)$$

Thus the Iwahori-Hecke algebra of $W(\Sigma_{\mathcal{O}}^{\vee})$, as defined in (4.0.13), is given by

$$\mathcal{H}(W(\Sigma_{\mathcal{O}}^{\vee}), q_F^{\lambda}) = \begin{cases} \mathcal{H}(\{1, s_{2\alpha+\beta}\}, q_F), & L/F \text{ is ramified} \\ \mathcal{H}(\{1, s_{2\alpha+\beta}\}, q_F^2), & L/F \text{ is unramified} \end{cases} \quad (4.1.19)$$

Therefore, the affine Hecke algebra in this case is given by

$$\mathcal{H}_{\text{aff}}(M^\beta) = \mathcal{H}(\{1, s_{2\alpha+\beta}\}, q_F^{f(L/F)}) \ltimes \mathbb{C}[\mathcal{O}] \quad (4.1.20)$$

4.1.21. Case III. If ω is unramified and $\sigma \neq \sigma(\tau)$ or $\chi^2\chi'$ is ramified,

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q_F^{n(\omega)+n(\sigma \times \Pi(\sigma))-n(\sigma)} \frac{(1 - \omega(\varpi)q_F^{-2s})(1 - \omega^{-1}(\varpi)q_F^{2s})}{(1 - \omega^{-1}(\varpi)q_F^{-1+2s})(1 - \omega(\varpi)q_F^{-1-2s})} \quad (4.1.22)$$

In this case, we have

$$\begin{aligned} X_\alpha(s) &= \omega(\varpi)q_F^{-2s} \\ q_{\alpha^*} &= 1, \quad q_\alpha = q_F \end{aligned} \quad (4.1.23)$$

Thus $\lambda(\alpha) = 1$ and $\lambda^*(\alpha) = 1$. The parameters in this case are simply q_F .

4.1.24. Case IV. If ω ramified and $\sigma \neq \sigma(\tau)$ or $\chi^2\chi'$ is ramified,

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q_F^{n(\sigma \times \Pi(\sigma))-n(\sigma)} \quad (4.1.25)$$

In this case, we have

$$q_\alpha = 1, \quad q_{\alpha^*} = 1 \quad (4.1.26)$$

Thus $\lambda(\alpha) = 0$ and $\lambda^*(\alpha) = 0$. Thus the parameters in this case are trivial.

4.1.2. The short root case

Now we give the explicit computation in the short root case. Assume the Levi factor M of $P = MN$ is generated by the short root of split G_2 . Let σ be an irreducible unitary supercuspidal representation of M . Let $\omega = \omega_\sigma$ be its central character. Then by [51, Proposition 6.2] the Plancherel measure $\mu(s\tilde{\alpha}, \sigma)$ is given by the formula

$$\mu(s\tilde{\alpha}, \sigma) = \begin{cases} \gamma(G/P)^2 q_F^{n(\sigma)+n(\sigma \otimes \omega)} \frac{(1 - \omega(\varpi)q_F^{-2s})(1 - \omega(\varpi)^{-1}q_F^{2s})}{(1 - \omega(\varpi)q_F^{-1-2s})(1 - \omega(\varpi)^{-1}q_F^{-1+2s})} & \text{if } \omega \text{ is unramified} \\ \gamma(G/P)^2 q_F^{n(\sigma)+n(\omega)+n(\sigma \otimes \omega)} & \text{otherwise} \end{cases} \quad (4.1.27)$$

Here $n(\sigma)$, $n(\omega)$ and $n(\sigma \otimes \omega)$ are the corresponding conductors.

4.1.28. Case I. If ω is unramified,

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q_F^{n(\sigma)+n(\sigma \otimes \omega)} \frac{(1 - \omega(\varpi)q_F^{-2s})(1 - \omega(\varpi)^{-1}q_F^{2s})}{(1 - \omega(\varpi)q_F^{-1-2s})(1 - \omega(\varpi)^{-1}q_F^{-1+2s})} \quad (4.1.29)$$

Comparing (4.1.29) with (4.0.9) implies that

$$q_\alpha = q_F, \quad q_{\alpha^*} = 1 \quad (4.1.30)$$

Since χ_s is an unramified character of M , we have

$$X_\alpha(\chi_s) = \omega(\varpi)q_F^{-2s}. \quad (4.1.31)$$

Recall from (4.0.11) that $q_F^{\lambda(\alpha)} = q_\alpha q_{\alpha^*} \in \mathbb{R}_{>1}$. Thus by (4.1.30), we have $q_F^{\lambda(\alpha)} = q_F$ and thus $\lambda(\alpha) = 1$ and $\lambda^*(\alpha) = 1$. Note that for $w \in W(M, \mathcal{O})$, one may check that

$$w(X_\alpha) = X_{w(\alpha)}. \quad (4.1.32)$$

Since $w(\alpha) = \alpha$ for $w \in W(M, \mathcal{O})$ for $G = G_2$, (4.1.32) is simply $w(X_\alpha) = X_\alpha$. On the other hand, by [53, Prop 1.1] we have

$$wX_\alpha(\chi) = w(X_\alpha(\chi)) = w(\chi(h_\alpha)) = \chi(w(h_\alpha)) = \chi(h_{w(\alpha)}) = \chi(h_\alpha) = X_\alpha(\chi). \quad (4.1.33)$$

Thus $wX_\alpha = X_\alpha = X_{w(\alpha)}$. Since $\Sigma_{\mathcal{O}}^\vee = \{3\alpha + 2\beta\}$ in the short root $M = M_\alpha$ case (see [51, p. 389]), we have

$$W(\Sigma_{\mathcal{O}}^\vee) = \{1, s_{3\alpha+2\beta}\} \quad (4.1.34)$$

Therefore we have the affine Hecke algebra

$$\mathcal{H}_{\text{aff}}(M_\alpha) = \mathcal{H}(\{1, s_{3\alpha+2\beta}\}, q_F) \ltimes \mathbb{C}[\mathcal{O}]. \quad (4.1.35)$$

4.1.36. Case II. In the other case,

$$\mu(s\tilde{\alpha}, \sigma) = \gamma(G/P)^2 q^{n(\sigma)+n(\omega)+n(\sigma \otimes \omega)} \quad (4.1.37)$$

Comparing (4.1.37) and (4.0.9) gives us

$$q_\alpha = 1, \quad q_{\alpha^*} = 1. \quad (4.1.38)$$

Therefore we have $q_F^\lambda = 1$ and thus $\lambda(\alpha) = 0$ and $\lambda^*(\alpha) = 0$. Therefore the affine Hecke algebra in this case is given by

$$\mathcal{H}_{\text{aff}}(M_\alpha) = \mathcal{H}(\{1, s_{3\alpha+2\beta}\}, 1) \ltimes \mathbb{C}[\mathcal{O}]. \quad (4.1.39)$$

Remark 4.1.40. The computations of Hecke algebras with explicit parameters in this section will be collected into tables in § 4.2.2.

4.2. Intertwining algebras

4.2.1. For $b \in F^\times / F^{\times 2}$, let $U_b(1, 1)$ be the quasi-split unitary group, and $U_b(2)$ the compact unitary group in two variables in $F(\sqrt{b})$. We write $F^\times / F^{\times 2} = \{1, \varepsilon, \varpi, \varepsilon\varpi\}$, and the possible unitary groups in 2 variables are:

$$U_\varepsilon(1, 1), U_\varepsilon(2), U_\varpi(1, 1), U_\varpi(2), U_{\varepsilon\varpi}(1, 1), U_{\varepsilon\varpi}(2).$$

The group $U_\varepsilon(1, 1)$ is an unramified group. The group $U_{\varpi'}(1, 1)$ is ramified, where $\varpi' \in \{\varpi, \varepsilon\varpi\}$.

4.2.2. We now classify the twisted Levi sequences in G_2 (up to conjugacy) for $\mathbf{M} = \mathbf{M}_\gamma$ with $\gamma \in \{\alpha, \beta\}$:

- (1) Essentially depth zero case: If ρ_{Σ_M} is an essentially depth-zero supercuspidal type on M , then Σ_M is of the form (M, y, ϕ, r, ρ_M) (hence in particular $M^0 = M$), where $K_{\Sigma_M} = M_{y,0} \simeq \mathrm{GL}_2(\mathfrak{o}_F)$ is a maximal compact subgroup of $\mathrm{GL}_2(F)$ and $r = \mathrm{depth}(\rho_{\Sigma_M})$ is an integer. If $r = 0$, we may assume that $\phi = 1$ without loss of generality.
 - (a) $\vec{\mathbf{G}} = (\mathbf{G})$ (here $\mathbf{G}^0 = \mathbf{G}$, it is a depth zero case: $r = 0$),
 - (b) $\vec{\mathbf{G}} = (\mathbf{M}^0, \mathbf{G})$ (here $\mathbf{G}^0 = \mathbf{M} = \mathbf{M}^0$ and $r \neq 0$).
- (2) Positive depth cases [4]:
 - (a) $\vec{\mathbf{G}} = (U_\varepsilon(1, 1), \mathbf{G})$,
 - (b) $\vec{\mathbf{G}} = (U_{\varpi'}(1, 1), \mathbf{G})$, with $\varpi' \in \{\varpi, \varepsilon\varpi\}$,
 - (c) $\vec{\mathbf{G}} = (\mathbf{M}^0, \mathbf{G})$,
 - (d) $\vec{\mathbf{G}} = (\mathbf{M}^0, \mathbf{M}, \mathbf{G})$,
 where $\mathbf{M}^0$ is a torus in $\mathbf{G}^0$.

When $\mathbf{M} = \mathbf{M}_\gamma$, we have three possibilities for $\mathbf{M}^0$, denoted $T_{\gamma,\varepsilon}$, $T_{\gamma,\varpi}$ and $T_{\gamma,\varpi}$. If ϕ_0 has trivial restriction to Z_M° , then it can be extended to a character of U_b and we use the same notation ϕ_0 to denote the extended character.

Let $\mathbb{G}_y^0$ denote the reductive quotient of G_y^0 . Let $\varpi' \in \{\varpi, \varepsilon\varpi\}$. We have

$$\mathbb{G}_y^0 = \begin{cases} U(1, 1) & \text{if } G^0 = U_\varepsilon(1, 1), \\ \mathrm{SO}_2 & \text{if } G^0 = U_{\varpi'}(1, 1). \end{cases} \quad (4.2.3)$$

Remark 4.2.4. The central character ω_σ of σ can be either ramified or unramified. It is unramified if and only if ω_{σ^0} is trivial. When ω_σ is ramified, ω_{σ^0} is quadratic.

Lemma 4.2.5. *We have*

$$W_G^s \simeq W_{G^0}^{s^0}. \quad (4.2.6)$$

Proof. The representation σ is regular and p is good for G , i.e. $p \neq 2, 3$ and does not divide the order ($= 1$) of the fundamental group of $\mathbf{G}_{\text{der}}$. Hence Lemma 2.2.9 applies and gives the desired isomorphism. $\square$

4.2.1. The intertwining algebras of types attached to G^0

4.2.7. The case $G^0 = M^0$. It occurs in both the essentially depth-zero case with $r \neq 0$ and in the positive depth cases. We have two possibilities for M^0 : either $M^0 \simeq \text{GL}_2(F)$ or M^0 is a torus. In both cases, the algebra $\mathcal{H}(G^0, \rho_{\mathcal{D}^0}) = \mathcal{H}(M^0, \rho_{\mathcal{D}^0})$ is commutative by [13, 5.5, 5.6].

4.2.8. The case $G^0 = \text{U}_\varepsilon(1, 1)$. If $W_{G^0}^{\mathfrak{s}^0} = \{1\}$, then $\mathcal{H}(G^0, \rho_{\mathcal{D}^0})$ is commutative, as seen in Remark 2.1.60. From now on we suppose that $W_{G^0}^{\mathfrak{s}^0} \neq \{1\}$. Let $a \mapsto \bar{a}$ be the non-trivial element of $\text{Gal}(L/F)$. Set

$$w^0 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad w_1 := \begin{pmatrix} 0 & \overline{\varpi}_L^{-1} \\ \varpi_L & 0 \end{pmatrix}, \quad \text{and} \quad \mathfrak{P} := \begin{pmatrix} \mathfrak{o}_L^\times & \mathfrak{o}_L \\ \mathfrak{p}_L & \mathfrak{o}_L^\times \end{pmatrix} \cap G^0. \quad (4.2.9)$$

Recall that $\widetilde{\rho_{\mathcal{D}^0}}$ denotes the contragredient representation of ρ^0 . By [10, §3.1], the Iwahori-Matsumoto presentation of $\mathcal{H}(\text{U}_\varepsilon(1, 1), \rho_{\mathcal{D}^0})$ is given by: $\mathcal{H}(\text{U}_\varepsilon(1, 1), \rho_{\mathcal{D}^0})$ is the space spanned by functions

$$T_{w_i} : G^0 \rightarrow \text{End}_G(V_{\widetilde{\rho_{\mathcal{D}^0}}}), \quad \text{for } i \in \{0, 1\}, \quad (4.2.10)$$

satisfying

$$T_{w_i}(pgp') = \widetilde{\rho_{\mathcal{D}^0}}(p)T_{w_i}(g)\widetilde{\rho_{\mathcal{D}^0}}(p'), \quad \text{where } p, p' \in \mathfrak{P} \text{ and } g \in G^0. \quad (4.2.11)$$

Here T_{w_i} is supported on $\mathfrak{P}w_i\mathfrak{P}$, and satisfies the quadratic relation

$$(T_{w_i} - q_F)(T_{w_i} + 1) = 0. \quad (4.2.12)$$

One can then deduce the Bernstein presentation of $\mathcal{H}(\text{U}_\varepsilon(1, 1), \rho_{\mathcal{D}^0})$ using [36, §3]. In particular, we have $q_F^{\lambda(\alpha)} = q_F$.

4.2.13. The case $G^0 = \text{U}_{\varpi'}(1, 1)$. Let $\varpi' \in \{\varpi, \varepsilon\varpi\}$. Since $\text{U}_{\varpi'}(1, 1)$ is ramified, by [10, §5.1.1], the algebra $\mathcal{H}(\text{U}_{\varpi'}(1, 1), \rho_{\mathcal{D}^0})$ has trivial parameters with $R(\mathcal{O}^0) \neq 1$ and $W_{\mathcal{O}^0} = 1$ if $\omega_\sigma|_{\mathfrak{o}_F^\times} \neq 1$; and the Hecke algebra has parameter q_F otherwise, in which case $W_{\mathcal{O}^0} \neq 1$ and $R(\mathcal{O}^0) = 1$.

4.2.2. The intertwining algebras of types attached to G

4.2.14. Long root essentially depth zero case.

(a) $r = 0$, $\chi^3 = 1$ case and $\sigma = \sigma(\tau)$ for $\tau = \text{Ind}_{W_L}^{W_F} \chi$:

We have ρ_M self-dual, σ and τ correspond via LLC for $\text{GL}_2(F)$. Since σ has depth zero, L/F is unramified (so $e(L/F) = 1$ and $f(L/F) = 2$). We have the following four cases:

- The central character $\omega_\sigma = 1$ and $\chi^2 \chi'^{-1}$ unramified. This corresponds to Case 4.1.1, in which case the Plancherel formula has a zero, and the Hecke algebra is affine non-commutative, with parameters q_F^3 and q_F . We have $W_{\mathcal{O}} \neq 1$ and $R(\mathcal{O}) = 1$. Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.
- The central character $\omega_\sigma \neq 1$ is ramified, and $\chi^2 \chi'^{-1}$ is unramified. This corresponds to Case 4.1.9, in which case the Plancherel formula has a zero, and the Hecke algebra is affine non-commutative, with parameters q_F^2 . We have $W_{\mathcal{O}} \neq 1$ and $R(\mathcal{O}) = 1$. Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.
- The central character $\omega_\sigma = 1$ and $\chi^2 \chi'^{-1}$ ramified. This corresponds to Case 4.1.21, in which case the Plancherel formula has a zero, and the Hecke algebra is affine non-commutative, with parameter q_F . We have $W_{\mathcal{O}} \neq 1$ and $R(\mathcal{O}) = 1$. Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.
- The central character $\omega_\sigma \neq 1$ is ramified, and $\chi^2 \chi'^{-1}$ is ramified. This corresponds to Case 4.1.24, in which case the Plancherel formula has no zero, and the Hecke algebra is affine commutative of the form $\mathbb{C}[R(\mathcal{O})]$ plus the translation part $\mathbb{C}[\mathcal{O}]$. We have $W_{\mathcal{O}} = 1$ (and we don't know what $R(\mathcal{O})$ is in this case). Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.

(b) $r = 0$ and $\sigma \neq \sigma(\tau)$: We have $\sigma = \sigma(\tau')$ where $\tau' = \text{Ind}_{W_L}^{W_F} \zeta$ for ζ such that $\zeta^{-1} = \bar{\zeta}$ (the Galois conjugate). Since σ is still depth zero, we still have L/F unramified.

- The central character $\omega_\sigma = 1$. This corresponds to Case 4.1.21, in which case the Plancherel formula has a zero, and the Hecke algebra is affine non-commutative, with parameters q_F . We have $W_{\mathcal{O}} \neq \{1\}$ and $R(\mathcal{O}) = \{1\}$. Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.
- The central character $\omega_\sigma \neq 1$ ramified. This corresponds to Case 4.1.24, in which case the Plancherel formula has no zero, and the Hecke algebra is affine commutative of the form $\mathbb{C}[R(\mathcal{O})]$ plus the translation part $\mathbb{C}[\mathcal{O}]$, we have $W_{\mathcal{O}} = \{1\}$ (and we don't know what $R(\mathcal{O})$ is in this case). Since $G = G^0$ in this case, $W_{\mathcal{O}} = W_{\mathcal{O}^0}$ and $R(\mathcal{O}) = R(\mathcal{O}^0)$.

(c) $r \neq 0$ essentially depth zero case: Recall from §4.2.7 that $G^0 = M = M^0$. Thus we have

$$W_{G^0}^s \subset N_{G^0}(M^0)/M^0 = N_M(M)/M = \{1\}.$$

By Lemma 4.2.5, we get $W_G^s = \{1\}$. In this case, $W(M, \mathcal{O}) = W(M^0, \mathcal{O}^0) = 1$. Thus the algebras $\mathcal{H}(G, \rho)$ and $\mathcal{H}(G^0, \rho^0)$ are both of the form $\mathbb{C}[\mathcal{O}]$, and they are isomorphic.

4.2.15. Table for long root essentially depth zero cases.

Table 4.2.15
Long root essentially depth zero cases.

r	$\mathcal{D}$	ω_σ	$\chi^2 \chi'^{-1}$	$R(\mathcal{O})$	$R(\mathcal{O}^0)$	L/F	$\#X_{\text{nr}}(M, \sigma)$	$W_{\mathcal{O}}$	$W_{\mathcal{O}^0}$	$\mathcal{H}(G, \rho)$	$\mathcal{H}(G^0, \rho^0)$
$r=0$	$((G, M), (y, \iota), (M_{y,0}, \rho_M))$	$\neq 1$	unramified χ cubic	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm, q_F^3, q_F	non-comm, q_F^3, q_F
		$\neq 1$	unramified χ cubic	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm, q_F^2, q_F^2	non-comm, q_F^2, q_F^2
		$\neq 1$	ramified χ cubic	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm, q_F, q_F	non-comm, q_F, q_F
		$\neq 1$	ramified χ cubic	*	*	unramified	2	$=1$	$=1$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$
		$=1$	χ not cubic	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm, q_F, q_F	non-comm, q_F, q_F
		$\neq 1$	N/A	*	*	unramified	2	$=1$	$=1$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$
$r \neq 0$	$((M, G), M), (y, \iota), (r, 0), (\phi, 1), (M_{y,0}, \rho_M)$	$\neq 1$	N/A	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}]$

4.2.16. Long root positive depth case

- (a) $U_{\varpi'}(1, 1)$ case: $\sigma = \sigma(\tau') \neq \sigma(\tau)$, where τ' is induction of some quadratic character. (Note that the cubic character only occurs in depth zero, because we are assuming $p \neq 2, 3$. There are two possibilities, $\phi_0|_{Z_M^0}$ could be either trivial or non-trivial:
- When $\phi_0|_{Z_M^0} = 1$ unramified, since $\sigma = \sigma(\tau') \neq \sigma(\tau)$, this corresponds to Case 4.1.21, in which case the Plancherel formula has a zero, and the Hecke algebra is affine non-commutative, with parameters q_F . We have $W_{\mathcal{O}} \neq \{1\}$ and $R(\mathcal{O}) = 1$. By 4.2.13, the Hecke algebra for G^0 also has q_F parameter. Thus we have $W_{\mathcal{O}^0} \neq \{1\}$ and $R(\mathcal{O}^0) = 1$.
 - When $\phi_0|_{Z_M^0} = \text{sign character ramified}$, since $\sigma = \sigma(\tau') \neq \sigma(\tau)$, this corresponds to Case 4.1.24, in which case the Plancherel formula has no zero, and the Hecke algebra is of the form $\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$. We have $W_{\mathcal{O}} = \{1\}$. By 4.2.13, we have $W_{\mathcal{O}^0} = \{1\}$ and $R(\mathcal{O}^0) \neq 1$. Thus $R(\mathcal{O}) \cong R(\mathcal{O}^0) \neq 1$ by Lemma 2.2.9.
- (b) $U_\epsilon(1, 1)$ case: $\sigma = \sigma(\tau') \neq \sigma(\tau)$, where τ' is the induction of some quadratic character.
- When $\phi_0|_{Z_M^0} = 1$ unramified, since $\sigma = \sigma(\tau') \neq \sigma(\tau)$, this corresponds to 4.1.21, in which case the Plancherel formula has a zero, and the Hecke algebra for G is affine non-commutative, with parameters q . We have $W_{\mathcal{O}} \neq \{1\}$ and $R(\mathcal{O}) = \{1\}$. From 4.2.8, we have $W_{\mathcal{O}^0} \neq \{1\}$ and $R(\mathcal{O}^0) = \{1\}$. Note that the cardinality of $\mathfrak{X}_{\text{nr}}(M, \sigma(\tau'))$ is 2 (see Remark 2.1.10).

4.2.17. Table for long root positive depth cases. We summarize the above in the following table:

Table 4.2.17
Long root positive depth cases.

M^0	$\phi_0 _{Z_M^0}$	ϕ_1	$\vec{G}$	$R(\mathcal{O})$	$R(\mathcal{O}^0)$	L/F	$\#X_{\text{nr}}(M, \mathcal{D})$	$W_{\mathcal{O}}$	$W_{\mathcal{O}^0}$	$\mathcal{H}(G, \rho)$	$\mathcal{H}(G^0, \rho^0)$
$T_{\beta, \varpi'}$	$= \text{sign character}$	$\neq 1$	$(U_{\varpi'}(1, 1), G)$	$\neq 1$	$\neq 1$	ramified	1	$=1$	$=1$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$
	$\neq 1 \neq \text{sign character}$	$=1$	(M^0, G)	$=1$	$=1$	ramified	1	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
	both	$\neq 1$	(M^0, M, G)	$=1$	$=1$	ramified	1	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
$T_{\beta, \epsilon}$	$=1$	$=1$	$(U_\epsilon(1, 1), G)$	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm. q_F, q_F	non-comm. q_F, q_F
	$\neq 1$	$=1$	(M^0, G)	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
	both	$\neq 1$	(M^0, M, G)	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$

4.2.18. Short root essentially depth zero case.

(a) $r = 0$, there are only two cases:

- When $\rho_M|_{Z_M^\circ} = 1$, this corresponds to the central character being unramified case, and in this case the Plancherel formula in 4.1.28 has a zero. Thus $W_{\mathcal{O}} \neq 1$ and thus $R(\mathcal{O}) = 1$. In this case the Hecke algebra is non-commutative, and the q -parameter is just $q = q_F$. The case for G^0 again follows from 4.2.13.
- When $\rho_M|_{Z_M^\circ} \neq 1$, this corresponds to the central character being ramified case, and in this case the Plancherel formula in 4.1.36 has no zero, and thus $W_{\mathcal{O}} = \{1\}$. In this case the Hecke algebra is commutative, and the q -parameter is trivial.

(b) $r \neq 0$ essentially depth zero case. The same argument as in §4.2.14(c) applies.

4.2.19. Table for short root essentially depth zero cases.

Table 4.2.19

Short root essentially depth zero cases.

r	$\mathcal{D}$	ω_σ	$R(\mathcal{O})$	$R(\mathcal{O}^0)$	L/F	$\#X_{\text{ss}}(M, \sigma)$	$W_{\mathcal{O}}$	$W_{\mathcal{O}^0}$	$\mathcal{H}(G, \rho)$	$\mathcal{H}(G^0, \rho^0)$
$r=0$	$((G, M), (y, \iota), (M_{y,0}, \rho_M))$	$=1$	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm, q_F, q_F	non-comm, q_F, q_F
		$\neq 1$	*	*	unramified	2	$=1$	$=1$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$
$r \neq 0$	$((M, G), M), (y, \iota), (r, 0), (\phi, 1), (M_{y,0}, \rho_M))$	$=1$ $\neq 1$	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$

4.2.20. Short root positive depth case.

(a) $G^0 = U_{\varpi'}(1, 1)$ case:

- When $\phi_0|_{Z_M^0} = 1$, the Plancherel formula on the G_2 side in 4.1.28 has a zero, and thus $W_{\mathcal{O}} \neq \{1\}$ and thus $R(\mathcal{O}) = \{1\}$. In this case the Hecke algebra $\mathcal{H}(G, \rho)$ is non-commutative, and the q -parameter is just $q = q_F$. By Lemma 2.2.9, we have $W(M^0, \mathcal{O}^0) \neq 1$, and since $W_{\mathcal{O}^0} \neq 1$ by 4.2.13, we have $R(\mathcal{O}^0) = 1$. Moreover, the Hecke algebra $\mathcal{H}(G^0, \rho^0)$ has parameter q_F by 4.2.13.
- When the central character (of $\text{GL}_2^{\text{short}}$) $\phi_0|_{Z_M^0} = \text{sign character} \neq 1$ is ramified, the Plancherel formula on the G_2 side in 4.1.36 has no zero, and thus $W_{\mathcal{O}} = \{1\}$. In this case the Hecke algebra $\mathcal{H}(G, \rho) = \mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$ has trivial q -parameter. On the other hand, since $I(\sigma)$ is reducible by [51, Proposition 6.2], we have $R_\sigma \neq 1$. Since $W_\sigma \rtimes R_\sigma = W(M, \sigma) \subseteq W(M, \mathcal{O}) = W_{\mathcal{O}} \rtimes R(\mathcal{O})$ and $W_{\mathcal{O}} = 1$, we have $R_\sigma \subseteq R(\mathcal{O})$ and thus $R(\mathcal{O}) \neq 1$. By Lemma 2.2.9, we also have $R(\mathcal{O}^0) \neq 1$ since $W_{\mathcal{O}^0} = 1$ by 4.2.13.

(b) $G^0 = U_\epsilon(1, 1)$ case: When $\phi_0|_{Z_M^0} = 1$ the Plancherel formula on the G_2 side in 4.1.28 has a zero, and thus $W_{\mathcal{O}} \neq \{1\}$ and thus $R(\mathcal{O}) = \{1\}$. In this case the Hecke algebra $\mathcal{H}(G, \rho)$ is non-commutative, and the q -parameter is just $q = q_F$. From 4.2.8, we have $W_{\mathcal{O}^0} \neq \{1\}$ and $R(\mathcal{O}^0) = \{1\}$.

4.2.21. Table for short root positive depth cases. We summarize the above in the following table:

Table 4.2.21
Short root positive depth cases.

M^0	$\phi_0 _{\mathbb{Z}_\ell^\times}$	ϕ_1	$\bar{G}$	$R(\mathcal{O})$	$R(\mathcal{O}^0)$	L/F	$\mathfrak{X}_{\text{nr}}(M, \sigma)$	$W_{\mathcal{O}}$	$W_{\mathcal{O}^0}$	$\mathcal{H}(G, \rho)$	$\mathcal{H}(G^0, \rho^0)$
$T_{\alpha, \varpi'}$	$=1$	$=1$	$(U_{\varpi'}(1, 1), G)$	$=1$	$=1$	ramified	1	$\neq 1$	$\neq 1$	non-comm. q_F	non-comm. q_F
	$= \text{sign character}$	$\neq 1$		$\neq 1$	$\neq 1$	ramified	1	$=1$	$=1$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}]$	$\mathbb{C}[R(\mathcal{O})] \ltimes \mathbb{C}[\mathcal{O}^0]$
	$\neq 1 \neq \text{sign character}$	$=1$	(M^0, G)	$=1$	$=1$	ramified	1	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
	both	$\neq 1$	(M^0, M, G)	$=1$	$=1$	ramified	1	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
$T_{\alpha, \varepsilon}$	$=1$	$=1$	$(U_{\varepsilon}(1, 1), G)$	$=1$	$=1$	unramified	2	$\neq 1$	$\neq 1$	non-comm. q_F	non-comm. q_F
	$\neq 1$	$=1$	(M^0, G)	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$
	both	$\neq 1$	(M^0, M, G)	$=1$	$=1$	unramified	2	$=1$	$=1$	$\mathbb{C}[\mathcal{O}]$	$\mathbb{C}[\mathcal{O}^0]$

4.2.22. We keep the notations of §2.1. The following theorem establishes the validity, for G_2 , of a generalization of a conjecture of Yu’s [56, Conjecture 0.2] for supercuspidal types, which was proved by Ohara in [45]. The following result shows that a stronger version of Theorem 2.2.14(2) holds for the group G_2 .

Theorem 4.2.23. *Let $p \neq 2, 3$. The algebras $\mathcal{H}^s(G) := \text{End}_G(\Pi_G^s)$ and $\mathcal{H}^s(G^0) := \text{End}_{G^0}(\Pi_{G^0}^s)$ are isomorphic.*

Proof. By Proposition 2.1.58, it is equivalent to show that the algebras $\mathcal{H}(G, \rho_{\mathcal{D}})$ and $\mathcal{H}(G^0, \rho_{\mathcal{D}^0})$ are isomorphic. The latter can be read directly from the Tables 4.2.15, 4.2.17, 4.2.19 and 4.2.21. $\square$

The following corollary is a stronger version of Lemma 4.2.5 for $G = G_2$.

Corollary 4.2.24. *The groups $R(\mathcal{O}) \simeq R(\mathcal{O}^0)$ and $W_{\mathcal{O}} \simeq W_{\mathcal{O}^0}$.*

Proof. This can be read directly from our Tables 4.2.15, 4.2.17, 4.2.19 and 4.2.21, with explanations given in the sections immediately preceding the tables. $\square$

4.2.25. On Lusztig’s conjecture. Let $L^s: W_{\text{aff}}^s \rightarrow \mathbb{N}$ be the weight function² on W_{aff}^s defined by

$$L^s(s_{\alpha}) := \lambda(\alpha) \quad \text{and} \quad L^s(s'_{\alpha}) := \lambda^*(\alpha). \tag{4.2.26}$$

In [40, §1.a], Lusztig made the following conjecture.

Conjecture 4.2.27. (Lusztig) The function L^s on the affine Weyl group W_{aff}^s is in the collection of weight functions described in [37–39].

Many cases of Conjecture 4.2.27 have been proved in [53], e.g. for principal series representations of G .

² i.e., $L^s(w) > 0$ for all $w \in W_{\text{aff}}^s - \{1\}$, and $L^s(w w') = L^s(w) + L^s(w')$ for any $w, w' \in W_{\text{aff}}^s$ such that $\ell(w w') = \ell(w) + \ell(w')$.

Theorem 4.2.28. Conjecture 4.2.27 holds for the group G_2 .

Proof. It follows from Tables 4.2.15, 4.2.17, 4.2.19 and 4.2.21. $\square$

5. Applications to other groups

Let N be a positive integer. Let J'_N denote the $N \times N$ -matrix $\begin{pmatrix} & & & 1 \\ & & 1 & \\ & \cdot & & \\ & \cdot & & \\ 1 & & & \end{pmatrix}$. When $N = 2n$, let $J_N := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$.

5.1. Symplectic group

The F -rational points of the symplectic group Sp_{2n} are given by

$$\mathrm{Sp}_{2n}(F) = \{g \in \mathrm{GL}_{2n}(F) : {}^t g J_{2n} g = J_{2n}\}. \quad (5.1.1)$$

Let P be the Siegel parabolic subgroup of $\mathrm{Sp}_{2n}(F)$, i.e. the maximal parabolic subgroup consisting of matrices whose lower left $n \times n$ -block is zero. The standard Levi factor of P is isomorphic to $\mathrm{GL}_n(F)$.

5.2. General symplectic group

The F -rational points of the algebraic group GSp_{2n} are given by

$$\mathrm{GSp}_{2n}(F) = \{g \in \mathrm{GL}_{2n}(F) : {}^t g J_{2n} g = \mu_n(g) J_{2n}, \mu_n(g) \in F^\times\}. \quad (5.2.1)$$

Let P be the Siegel parabolic subgroup of $\mathrm{GSp}_{2n}(F)$. The standard Levi factor of P is isomorphic to $\mathrm{GL}_n(F) \times \mathrm{GL}_1(F)$.

5.3. Special orthogonal group

The F -rational points of the algebraic group SO_N are given by

$$\mathrm{SO}_N(F) = \{g \in \mathrm{GL}_N(F) : {}^t g J'_N g = J'_N, \det(g) = 1\}. \quad (5.3.1)$$

Let P be the Siegel parabolic subgroup of $\mathrm{SO}_N(F)$. The standard Levi factor of P is isomorphic to $\mathrm{GL}_n(F)$, where $N = 2n + 1$ or $N = 2n$.

The LLC for $\mathrm{GL}_n(F)$, established in [29,27,49], shows that the L -packets are always singletons in this case. Thus, by Proposition 1.2.5, the properties (1) and (2) are satisfied in the three cases above.

References

- [1] A.-M. Aubert, P. Baum, R. Plymen, M. Solleveld, Conjectures about p -adic groups and their non-commutative geometry, in: *Around Langlands Correspondences*, Orsay, 2015, in: *Contemp. Math.*, vol. 691, Amer. Math. Soc., 2017, pp. 15–51, MR 3666049.
- [2] A.-M. Aubert, P. Baum, R. Plymen, M. Solleveld, Hecke algebras for inner forms of p -adic special linear groups, *J. Inst. Math. Jussieu* 16 (2) (2017) 351–419, MR 3615412.
- [3] A.-M. Aubert, P. Baum, R. Plymen, M. Solleveld, The principal series of p -adic groups with disconnected center, *Proc. Lond. Math. Soc.* 114 (2017) 798–854, MR 3653247.
- [4] A.-M. Aubert, M. Emory, M. Fox, J.-L. Kim, Y. Xu, Notes from WIN5.
- [5] J. Adler, M. Mishra, Regular Bernstein blocks, *J. Reine Angew. Math.* (775) (2021) 71–86.
- [6] A.-M. Aubert, A. Moussaoui, M. Solleveld, Affine Hecke algebras for Langlands parameters, arXiv: 1701.03593, 2017.
- [7] A.-M. Aubert, A. Moussaoui, M. Solleveld, Generalizations of Springer correspondence and cuspidal Langlands parameters, *Manuscr. Math.* 157 (2018) 121–192, MR 3845761.
- [8] J. Arthur, The Endoscopic Classification of Representations: Orthogonal and Symplectic Groups, *Colloquium Publications*, vol. 61, American Mathematical Society, 2013.
- [9] A.-M. Aubert, Y. Xu, The explicit local Langlands correspondence for G_2 , arXiv:2208.12391, 2022.
- [10] P. Badea, Hecke algebras for covers of principal series Bernstein components in quasisplit unitary groups over local fields, PhD. Thesis, Radboud Universiteit Nijmegen, 2020.
- [11] J.N. Bernstein, Le “centre” de Bernstein, in: P. Deligne (Ed.), *Representations of Reductive Groups over a Local Field*, in: *Travaux en Cours*, Hermann, Paris, 1984, pp. 1–32, MR 771671.
- [12] C. Bushnell, P. Kutzko, The Admissible Dual of $GL(N)$ via Compact Open Subgroups, *Annals of Mathematics Studies*, vol. 129, Princeton University Press, Princeton, N.J., 1993, MR 1204652.
- [13] C.J. Bushnell, P.C. Kutzko, Smooth representations of reductive p -adic groups: structure theory via types, *Proc. Lond. Math. Soc.* (3) 77 (3) (1998) 582–634, MR 1643417.
- [14] C. Blondel, Une méthode de construction de types induits et son application à G_2 , *J. Algebra* (213) (1999) 231–271, MR 1674685.
- [15] A. Bourgeois, P. Mezo, Functoriality for supercuspidal L -packets, arXiv:2109.09552, 2021.
- [16] A. Borel, Automorphic L -Functions, *Proc. Symp. Pure Math.*, vol. 33, Amer. Math. Soc., 1979, pp. 27–61.
- [17] P. Bakić, G. Savin, The Gelfand–Graev representation of $SO(2n + 1)$ in terms of Hecke algebras, arXiv:2011.02456, 2020.
- [18] F. Bruhat, J. Tits, Groupes réductifs sur un corps local, I: Données radicielles valuées, *Publ. Math. IHÉS* 41 (1972) 1–251.
- [19] R.W. Carter, *Finite Groups of Lie Type: Conjugacy Classes and Complex Characters*, Wiley Classics Library, 1993.
- [20] R. Deseine, *Autour des représentations complexes et modulaires des groupes réductifs p -adiques*, Thèse Université Paris-Saclay, 2021.
- [21] P. Deligne, G. Lusztig, Representations of reductive groups over finite fields, *Ann. Math.* (103) (1976) 103–161.
- [22] J. Fintzen, Types for tame p -adic groups, *Ann. Math.* 193 (2) (2021) 303–346, MR 4199732.
- [23] J. Fintzen, T. Kaletha, L. Spice, A twisted Yu construction, Harish-Chandra characters, and endoscopy, arXiv:2106.09120, 2021.
- [24] S. Gelbart, G. Jacquet, A relation between automorphic representations of $GL(2)$ and $GL(3)$, *Ann. Sci. Éc. Norm. Supér.* (4) 11 (4) (1978) 471–542, MR 533066.
- [25] T. Haines, The stable Bernstein center and test functions for Shimura varieties, in: *Automorphic Forms and Galois Representations*, in: *London Mathematical Society Lecture Note Series*, vol. 415, Cambridge University Press, 2014, pp. 118–186.
- [26] V. Heiermann, Opérateurs d’entrelacement et algèbres de Hecke avec paramètres d’un groupe réductif p -adique: le cas des groupes classiques, *Sel. Math. New Ser.* 17 (3) (2011) 713–756, MR 2827179.
- [27] G. Henniart, Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique, *Invent. Math.* 139 (2) (2000) 439–455, MR 1738446.
- [28] J. Hakim, F. Murnaghan, Distinguished tame supercuspidal representations, *Int. Math. Res. Pap.* (2) (2008) rpn005.
- [29] M. Harris, R. Taylor, *The Geometry and Cohomology of Some Simple Shimura Varieties*, *Annals of Mathematics Studies*, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich, MR 1876802.

- [30] T. Kaletha, Epipelagic L -packets and rectifying characters, *Invent. Math.* 202 (1) (2015) 1–89, MR 3402796.
- [31] T. Kaletha, Regular supercuspidal representations, *J. Am. Math. Soc.* 32 (4) (2019) 1071–1170, MR 4013740.
- [32] T. Kaletha, Supercuspidal L -packets, arXiv:1912.03274, 2019.
- [33] T. Kaletha, Representations of Reductive Groups over Local Fields, 2022.
- [34] J.-L. Kim, J.-K. Yu, Construction of tame types, in: Representation Theory, Number Theory, and Invariant Theory, in: *Progr. Math.*, vol. 323, Birkhäuser/Springer, Cham, 2017, pp. 337–357.
- [35] G. Lusztig, Intersection cohomology complexes on a reductive group, *Invent. Math.* (75) (1984) 205–272.
- [36] G. Lusztig, Affine Hecke algebras and their graded version, *J. Am. Math. Soc.* 2 (3) (1989) 599–635, MR 991016.
- [37] G. Lusztig, Intersection cohomology methods in representation theory, in: *Proc. Int. Congr. Math.*, Kyoto, 1990, in: *Math. Soc. Japan*, Springer Verlag, 1991, pp. 155–174, MR732546.
- [38] G. Lusztig, Classification of unipotent representations of simple p -adic groups, *Int. Math. Res. Not.* (11) (1995) 517–589, MR 1927955.
- [39] G. Lusztig, Classification of unipotent representations of simple p -adic groups. II, *Represent. Theory* (6) (2002) 243–289, MR 1927955.
- [40] G. Lusztig, Open problems on Iwahori-Hecke algebras, *News. - Eur. Math. Soc.* (2023), in press, arXiv:2006.08535.
- [41] M. Mishra, Bernstein center of supercuspidal blocks, *J. Reine Angew. Math.* (748) (2019) 297–304.
- [42] A. Moussaoui, Centre de Bernstein dual pour les groupes classiques, *Represent. Theory* (21) (2017) 172–246, MR 3694312.
- [43] A. Moussaoui, Proof of the Aubert-Baum-Plymen-Solleveld conjecture for split classical groups, in: *Around Langlands Correspondences*, Orsay, 2015, in: *Contemp. Math.*, vol. 691, Amer. Math. Soc., 2017, pp. 257–281, MR 3618046.
- [44] G. Muić, The unitary dual of p -adic G_2 , *Duke Math. J.* 90 (3) (1997) 465–493, MR 1480543.
- [45] K. Ohara, Hecke algebras for tame supercuspidal types, arXiv:2101.01873, 2021.
- [46] A. Roche, Types and Hecke algebras for principal series representations of split reductive p -adic groups, *Ann. Sci. Éc. Norm. Supér.* (4) 31 (3) (1998) 361–413, MR 1621409.
- [47] A. Roche, Parabolic induction and the Bernstein decomposition, *Compos. Math.* 134 (2) (2002) 113–133, MR 1934305.
- [48] A. Roche, The Bernstein decomposition and the Bernstein centre, in: *Ottawa Lectures on Admissible Representations of Reductive p -Adic Groups*, in: *Fields Inst. Monogr.*, vol. 26, Amer. Math. Soc., Providence, RI, 2009, pp. 3–52, MR 2499587.
- [49] P. Scholze, The local Langlands correspondence for GL_n over p -adic fields, *Invent. Math.* 192 (3) (2013) 663–715, MR 3049932.
- [50] F. Shahidi, Third symmetric power L -functions for $GL(2)$, *Compos. Math.* 70 (3) (1989) 245–273, MR 1002045.
- [51] F. Shahidi, Langlands’ conjecture on Plancherel measures for p -adic groups, in: *Harmonic Analysis on Reductive Groups*, Brunswick, ME, 1989, in: *Progr. Math.*, vol. 101, Birkhäuser Boston, Boston, MA, 1991, pp. 277–295, MR 1168488.
- [52] A.J. Silberger, Introduction to Harmonic Analysis on Reductive p -Adic Groups, *Mathematical Notes*, vol. 23, Princeton University Press, Princeton, N.J., 1979, pp. 1971–1973, University of Tokyo Press, Tokyo, Based on lectures by Harish-Chandra at the Institute for Advanced Study, MR 544991.
- [53] M. Solleveld, Parameters of Hecke algebras for Bernstein components of p -adic groups, arXiv:2103.13113, 2021.
- [54] M. Solleveld, Endomorphism algebras and Hecke algebras for reductive p -adic groups, *J. Algebra* (606) (2022) 371–470.
- [55] J.-L. Waldspurger, La formule de Plancherel pour les groupes p -adiques (d’après Harish-Chandra), *J. Inst. Math. Jussieu* 2 (2) (2003) 235–333, MR 1989693.
- [56] J.-K. Yu, Construction of tame supercuspidal representations, *J. Am. Math. Soc.* (3) (2001) 579–622, MR 1824988.